Association of expression of mRNA encoding the PGF_{2α} receptor with luteal cell apoptosis in ovaries of pseudopregnant mice

K. Hasumoto¹, Y. Sugimoto¹, A. Yamasaki¹, K. Morimoto¹,
A. Kakizuka², M. Negishi¹ and A. Ichikawa^{1*}

¹Department of Physiological Chemistry, Faculty of Pharmaceutical Sciences, and

²Department of Pharmacology, Faculty of Medicine, Kyoto University, Sakyo-ku, Kyoto 606, Japan

The relationship between the expression of mRNA encoding the prostaglandin F_{2α} (PGF_{2α}) receptor and luteal cell apoptosis as determined by oligonucleosome formation was determined in mouse corpora lutea on days 2 (early phase), 6 (mid-phase), and 11 and 13 (late phase) of pseudopregnancy. No signals for mRNA encoding the PGF_{2α} receptor were detected in the ovary as shown by RNA blot analysis of pregnant mares' serum gonadotrophin (PMSG)-treated mice. The expression of the mRNA that could be detected in the corpora lutea on day 2 after hCG treatment was low, but it was increased on day 6 and reached a plateau on days 11 and 13. On days 11 and 13, *in situ* hybridization signals for mRNA encoding the PGF_{2α} receptor were localized to large luteal cells of the corpora lutea, especially the cells in the superficial layer. However, ethidium bromide staining revealed marked oligonucleosome formation in the corpora lutea on days 11 and 13. Similarly, positive signals of *in situ* nick-DNA-end labelling were also detected in the corpora lutea on days 11 and 13. Analysis of the adjacent sections of the corpora lutea on day 13 showed that both signals for DNA strand breaks and mRNA encoding the PGF_{2α} receptor were co-expressed in the corpus luteum. These results suggest that apoptosis during structural luteolysis closely associates with the increased expression of mRNA encoding the PGF_{2α} receptor in luteal cells of pseudopregnant mice.

Introduction

Prostaglandin F_{2α} (PGF_{2α}) is one of the major arachidonate metabolites produced by the cyclooxygenase system in the mammalian female reproductive tissues (Coleman *et al.*, 1990; Boone *et al.*, 1993). Accumulating evidence has suggested that PGF_{2α} is of uterine or ovarian origin, depending on animal species, and that it is responsible for luteal regression in the ovary (Horton and Poyser, 1976; Rothchild, 1981; Hansel and Dowd, 1986; McCracken and Schramm, 1988). As an initiator of luteolysis, PGF_{2α} is known to inhibit steroidogenesis through PGF_{2α} receptors by increasing the intracellular Ca²⁺ concentration or inhibiting cAMP synthesis. Recent evidence indicates that the disruption of the corpus luteum during luteolysis involves apoptosis on the basis of pharmacological experiments in rats and cattle (Zelevnik *et al.*, 1989; Sawyer *et al.*, 1990; Juengel *et al.*, 1993). However, little is known about on temporal relationship between luteal cell apoptosis and the expression of the PGF_{2α} receptor during the oestrous cycle. Recently, mouse cDNA clone for the PGF_{2α} receptor was isolated, and abundant expression of the receptor mRNA was demonstrated in the corpus luteum in the pregnant mouse ovary (Sugimoto *et al.*, 1994). The objective of this study was to investigate the possible association between the expression

of mRNA encoding the PGF_{2α} receptor and apoptosis in corpora lutea in pseudopregnant mice.

Materials and Methods

Animals

Immature 21-day-old female ddY mice were obtained from Shimizu Experimental Materials Co. (Kyoto). Animals were housed under standard conditions of light and temperature. Immature female mice were injected intraperitoneally with 5 iu pregnant mares' serum gonadotrophin (PMSG, Teikoku Hormone MFG, Tokyo), which was followed 48 h later by a injection of 5 iu hCG to obtain luteinized ovaries. These mice were then mated with castrated male mice. Ovaries ($n \geq 4$) were collected at specific time points, frozen in liquid nitrogen and stored at -80°C until assayed. Ovaries were categorized as early (2 days after hCG treatment), mid- (6 days after hCG treatment) or late (11 and 13 days after hCG treatment) luteal phase.

RNA blot and *in situ* hybridization

Total RNA was extracted from ovaries using the procedures of Chomczynski and Sacchi (1987). Total RNA (10 μg per lane)

*Correspondence.
Received 29 May 1996.

was separated by electrophoresis on a 1.2% (w/v) agarose-formaldehyde gel, transferred onto nylon membranes (Hybond-N, Amersham Corp.), and hybridized with a ^{32}P -labelled *NotI/HindIII* fragment of MC205, which is a functional cDNA clone for mouse $\text{PGF}_{2\alpha}$ receptor (Sugimoto *et al.*, 1994). At least three independent experiments were carried out.

The linearized MC205 was used as template DNA and an antisense riboprobe was transcribed by T3 RNA polymerase (Stratagene, La Jolla) in the presence of [^{35}S]CTPaS. Unlabelled antisense riboprobe was synthesized similarly with unlabelled nucleotides. The resultant riboprobes were degraded to about 150 bases by alkaline hydrolysis. The ovaries were removed from each animal ($n \geq 4$) at each specific time phase. Three or four ovaries were frozen together in OCT compound (Miles, Elkhart, IN), cut out, and sections 10 μm in thickness were mounted on poly L-lysine-coated slides. The sections were fixed with 4% (w/v) formaldehyde, treated with 0.25% (v/v) acetic anhydride, and dehydrated. The slides were hybridized with a ^{35}S -labelled antisense RNA probe at 57°C for 4 h in the presence or absence of a 100-fold excess of unlabelled probe (Sugimoto *et al.*, 1994). Hybridized sections were washed, treated with 20 μg RNase A ml^{-1} , and washed again. The slides were air-dried and dipped in NTB2 emulsion (Kodak) diluted 1:1 with distilled water. After exposure for 4 weeks at 4°C, the dipped slides were developed in Kodak D-19 developer, fixed and counter-stained with haematoxylin and eosin.

Measurement of serum progesterone

Blood samples were collected from each animal ($n \geq 4$) when the ovaries were removed. Measurement of progesterone was performed using a radioimmunoassay kit manufactured by Dai-ichi Radioisotope Laboratory (Tokyo). Serum concentrations of progesterone on days 8, 10, and 12 of pseudopregnancy was 295 ± 38.1 , 31.9 ± 4.8 , and 29.8 ± 6.3 (means $\pm$ SEM) nmol l^{-1} , respectively. The coefficients of variation of analysis of progesterone were 5.8% intra-assay and 7.8% interassay.

Detection of oligonucleosome formation by electrophoresis and *in situ* detection of the cells undergoing DNA fragmentation (TUNEL method)

The genomic DNA was isolated using a standard procedure and separated according to size in a 2% (w/v) agarose gel by

Fig. 1. RNA blot analysis of mRNA encoding the prostaglandin $\text{F}_{2\alpha}$ ($\text{PGF}_{2\alpha}$) receptor expressed in mouse ovaries during gonadotrophin-primed pseudopregnancy. Total RNA (10 μg of each sample) isolated from ovaries at different phases of the pseudopregnancy were separated, transferred onto a nylon membrane, and then hybridized with a ^{32}P -labelled $\text{PGF}_{2\alpha}$ receptor-specific probe. Hybridization signals for mRNA encoding glyceraldehyde-3-phosphate dehydrogenase are shown in the lower panel. Arrowheads, three major bands of the $\text{PGF}_{2\alpha}$ receptor transcripts; P represents ovaries treated with pregnant mares' serum gonadotrophin for 48 h; 2, 6, 11, and 13 represent ovaries obtained 2, 6, 11 and 13 days after hCG treatment, respectively.

electrophoresis. The gels were stained with ethidium bromide and viewed with a UV lightbox. The extent of DNA breakdown in fixed ovarian tissue sections was assessed using an *in situ* terminal transferase reaction to the nonisotopically labelled free 3'-ends of the DNA (Schmitz *et al.*, 1991; Thiry, 1992). *In situ* DNA labelling analysis was performed using ApopTag (Oncor, Gaithersburg). The ovary sections described above ($n \geq 4$) were fixed in 4% (w/v) neutral buffered formaldehyde, and then post-fixed in ethanol:acetic acid (2:1) at -20°C . Endogenous peroxidase was quenched in 2.0% (w/v) hydrogen peroxide in PBS, washed with PBS, and subsequently labelled with digoxigenin-11-dUTP using a terminal transferase enzyme at 37°C for 1 h. In the negative control sections, the procedure was performed without the terminal transferase enzyme. The labelled sections were washed and subsequently incubated with anti-digoxigenin-peroxidase at 37°C for 30 min. The localization of digoxigenin-anti-digoxigenin-peroxidase complexes was detected by incubation of the slides

Fig. 2. Expression and localization of the mRNA encoding the prostaglandin $\text{F}_{2\alpha}$ ($\text{PGF}_{2\alpha}$) receptor in pseudopregnant mouse ovaries primed with gonadotrophin. A series of sections 10 μm in thickness were hybridized with ^{35}S -labelled antisense RNA probe for the $\text{PGF}_{2\alpha}$ receptor in the absence (a–l) or presence (m) of a 100-fold excess of unlabelled probe. After being exposed to the emulsion, the sections were developed and stained with haematoxylin and eosin. Figures show bright (a–d, i, k and l) and dark field (e–h, j and m) photomicrographs of the sections obtained from the ovaries treated with pregnant mares' serum gonadotrophin (PMSG) for 48 h (a and e), ovaries treated with PMSG followed by hCG for 2 days (b, f, and i–k), and ovaries obtained 6 (c, g, and l) or 11 days (d, h, and m) after hCG treatment. The first and second horizontal rows are bright and dark field micrographs of the same sections, respectively (a and e; b and f; c and g; d and h, respectively). Since the ovaries shown in a–h are on the same slide, the intensities of the signals are comparable. In micrographs b–d and i, the corpora lutea are seen in dark grey because of the dense hybridization signals. The magnifications of micrographs a–h and m are the same. Photomicrographs i–k show the hybridization signals in the corpus luteum. Arrows (k) or arrowheads (l) indicate the large luteal cells showing strong signals. Scale bars represent 500 μm (a), 100 μm (i), and 50 μm (k and l), respectively.

with diaminobenzidine at room temperature, and the sections were counterstained with methylgreen.

Results

Expression of mRNA encoding the PGF_{2α} receptor in the corpus luteum of pseudopregnant mice

The relative abundance of mRNA encoding the PGF_{2α} receptor in the ovary during the oestrous cycle was investigated by performing RNA blot hybridization in the ovaries at different stages of pseudopregnancy (Fig. 1). Expression of the receptor mRNA could not be detected in untreated immature ovaries (data not shown) or ovaries obtained 48 h after PMSG treatment, while three major species of the mRNA with estimated sizes of 2.0, 2.3 and 6.0 kilobases (kb) were detected in the hCG-treated ovaries. Significant expression of mRNA encoding the PGF_{2α} receptor was consistently observed in the ovaries of mice on day 2; higher expression was also detected on day 6 and it reached maximum values on days 11 and 13 of pseudopregnancy.

Cellular localization of mRNA encoding the PGF_{2α} receptor was examined by *in situ* hybridization (Fig. 2). The hybridization signals of the receptor mRNA were specifically present in corpora lutea, but no signals were detected in the unovulated developing follicles (Fig. 2, a and e). On day 2, significant numbers of signals were detectable in all corpora lutea (Fig. 2, b and f), and large luteal cells localized near the surface of the corpus showed strong signals (Fig. 2, i–k). On day 6 of pseudopregnancy, the expression increased in most of the corpora lutea (Fig. 2, c and g), and strong signals were localized in the large cells, especially the cells in the superficial layers (Fig. 2l). On day 11 of pseudopregnancy, the highest expression was observed (Fig. 2, d and h).

Oligonucleosome formation in corpus luteum of pseudopregnancy

In our pseudopregnant model, the serum concentrations of progesterone declined on day 10 (see Materials and Methods), which indicated that the onset of luteal regression (functional luteolysis) occurred on day 10. We, therefore, electrophoretically examined the ethidium bromide-stained DNA isolated from the ovaries during the mid- to late phases (Fig. 3). Slight oligonucleosome formation was seen on day 8, which reflects apoptosis involved in follicular atresia (see below). Marked oligonucleosome formation was noted on days 10 and 12. An *in situ* nick-DNA-end labelling technique was used to examine the cells that underwent apoptosis in the ovary during pseudopregnancy. Typical sections of ovaries at each stage are

Fig. 3. Ethidium bromide-stained genomic DNA isolated from mouse ovaries during the mid- to late phase of pseudopregnancy (20 µg per lane). The day of pseudopregnancy is given at the top of the figure.

shown (Fig. 4). The sections shown in Fig. 4 a–c, are on the same slide. Positively-stained nuclei appeared dark brown. Slightly stained regions can be seen around positively stained nuclei but they were thought to be artefacts from the fixation. In every section during the different phases, positive staining of the atretic follicles was observed (Tilly *et al.*, 1991), and the staining intensities were constant during the different phases. In corpora lutea, no specific signals were observed in the sections on days 2 and 6 (Fig. 4a and b, respectively). In contrast,

Fig. 4. Appearance of apoptotic cells and the expression of mRNA encoding the PGF_{2α} receptor in the corpora lutea during the late phases of pseudopregnancy in mice. A series of sections 10 µm in thickness was used for *in situ* analysis of DNA fragmentation of the apoptotic cells (a–d and h). Positively-stained nuclei appear dark brown. Slightly stained regions are seen around the positively stained nuclei, but they were thought to be artefacts from the fixation. After staining for the DNA strand breaks, the sections were counterstained by methyl green. The adjacent sections were hybridized with a ³⁵S-labelled antisense RNA probe for PGF_{2α} receptor (e and i, darkfield). The sections obtained 2 (a), 6 (b), 11 (c–f), or 13 days (g–i) after hCG treatment are shown. Photomicrographs of c–f or g–i are the continuous adjacent sections of the same ovary, respectively. Micrographs (f) and (g) show the sections counterstained by haematoxylin and eosin. The micrograph of d shows the section of a negative control for DNA end staining. The magnifications of micrographs are the same. Scale bars represent 250 µm.

degenerating cells with DNA breakdown signals were seen in the corpora lutea of the ovaries on days 11 and 13 (Fig. 4c and h). On day 13, the number of degenerating luteal cells increased compared with day 11 (Fig. 4h). Furthermore, both signals for DNA strand breaks and for mRNA encoding the PGF_{2α} receptor were co-expressed in the same corpus luteum on days 11 and 13 (Fig. 4, c and e; h and i). Thus, expression of mRNA encoding the PGF_{2α} receptor in the luteal cells remained high until the cells exhibited DNA fragmentation.

Discussion

The maintenance of the female reproductive cycle is achieved by several processes including cell proliferation, differentiation and cell death. Kerr *et al.* (1972) and Wyllie *et al.* (1980) indicated on the basis of histological observations that luteal regression is elicited by a highly specific programmed cell death called apoptosis. Sawyer *et al.* (1990) confirmed the histological appearance of luteal regression as apoptosis, and Zeleznik *et al.* (1989) demonstrated the activity of a Ca²⁺-Mg²⁺-dependent endonuclease, one of the DNases known to associate with programmed cell death, in rat luteal cells. Considering the known role of PGF_{2α} in luteal regression, we have suggested that PGF_{2α} receptor-mediated intracellular signalling directly influences the apoptosis of rodent corpora lutea. However, the temporal relationship between the expression of mRNA encoding the PGF_{2α} receptor and luteal cell apoptosis has not been established. In the present study, we demonstrated that the high expression of the receptor mRNA in the luteal cells on day 6 was followed by a plateau on day 11. In our pseudopregnant model, functional luteolysis, which is a decline of steroidogenesis, occurred on day 10. Similarly, electrophoretic or *in situ* analyses revealed that slight DNA fragmentation occurred in the luteal cells on days 10 and 11, and marked oligonucleosome formation was observed on days 12 and 13. Since nuclease activation is thought to be a relatively late event in apoptosis (Gavrieli *et al.*, 1992), the luteal cells may receive 'death signals' at the beginning of the late phase before day 10 when the highest expression of mRNA encoding the PGF_{2α} receptor was observed. Furthermore, the increased expression was maintained until most of the luteal cells underwent apoptosis. Thus, the present study has demonstrated that the expression of the PGF_{2α} receptor closely associates with apoptosis in mouse luteal cells.

Although luteolytic PGF_{2α} originates from the uterus in sheep and cattle, recent reports have suggested that a significant amount of PGF_{2α} is synthesized locally in the corpora lutea in rats (Boone *et al.*, 1993; Olofsson and Leung, 1994). In fact, using tissue samples of pseudopregnant rats, Olofsson and Selstam reported that the PGF_{2α} content in the corpus luteum was increased on day 11, and then increased further on day 13 of pseudopregnancy (Olofsson and Selstam, 1988). If the PGF_{2α} content in the corpora lutea is markedly increased at the beginning of the late phase, then it must become higher at the end of the late phase. Furthermore, it has also been suggested that intraluteal PG production is essential for structural luteolysis induced by prolactin in rats (Sánchez-Criado *et al.*, 1987). The present study together with these previous reports strongly suggests that PGF_{2α} and its receptor signalling are involved in luteal cell apoptosis.

A number of experiments showed that programmed cell death associates with endogenous endonuclease activity (Wyllie *et al.*, 1980; McConkey *et al.*, 1990). Mg²⁺-Ca²⁺-dependent endonuclease is known to be one of these DNases, and its activation was reported in regressing rat luteal cells (Zeleznik *et al.*, 1989). It is possible in this respect that PGF_{2α} and its receptor stimulate this enzyme, since the PGF_{2α} receptor increases the intracellular Ca²⁺ concentration via Gq-phospholipase C (Leung *et al.*, 1986; Ito *et al.*, 1994; Sugimoto *et al.*, 1994). However, this activity does not fully account for the actions of PGF_{2α} (Boone *et al.*, 1993). Recently, it was demonstrated that treatment of luteal tissue with PGF_{2α} caused a rapid but transient increase in superoxide radicals in the plasma membrane (Sawada and Carlson, 1991). Superoxide radicals activate phospholipase A₂, and inhibit the production of cAMP and progesterone (Gatzuli *et al.*, 1991; Ikebuchi *et al.*, 1991). Thus, at least two pathways may be involved in PGF_{2α} and its receptor signalling.

PGF_{2α} may be a physiological mediator in the induction of follicular rupture (Carvalho *et al.*, 1989). PGF_{2α} concentrations increase in the preovulatory fluid of several species (Priddy *et al.*, 1989). If PGF_{2α} induces follicular rupture, expression of mRNA encoding the PGF_{2α} receptor can be detected in preovulatory follicles. In the present study, by the administration of PMSG, follicles developed and became enriched in the ovaries, but no significant expression of the PGF_{2α} receptor was detected. However, 2 days after hCG injection, the signals were detected in the large luteal cells in the corpora lutea. In the study reported here the expression of the PGF_{2α} receptor in the corpora lutea was detected even in the ovaries obtained 24 h after hCG injection (data not shown). Since ovulation usually occurred about 12 h after hCG treatment, the mRNA encoding the PGF_{2α} receptor should have been induced soon after ovulation. Therefore, it is possible that the transcription of mRNA encoding the PGF_{2α} receptor starts when granulosa cells differentiate into luteal cells. However, a more detailed analysis is required to elucidate the transcriptional regulation of the PGF_{2α} receptor during follicular maturation, ovulation and luteinization of ruptured follicles.

This work was supported in part by grants for scientific research from the Ministry of Education, Science and Culture of Japan, the Kato Memorial Bioscience Foundation, the Sankyo Life Science Research Foundation, and the Kanae Foundation for Research on New Medicine. The authors thank J. A. McCracken, H. Fujiwara and T. Endo for helpful advice, and also thank A. Uesugi for photographic assistance.

References

- Boone DL, Currie WD and Leung PCK (1993) Arachidonic acid and cell signalling in the ovary and placenta *Prostaglandins, Leukotrienes and Essential Fatty Acids* **48** 79–87
- Carvalho CB, Yeik BS and Murdoch WJ (1989) Significance of follicular cyclooxygenase and lipoxygenase pathways of metabolism of arachidonate in sheep *Prostaglandins* **37** 553–558
- Chomczynski P and Sacchi N (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate–phenol–chloroform extraction *Analytical Biochemistry* **162** 156–159
- Coleman RA, Kennedy I, Humphrey PPA, Bruce K and Lumley P (1990) Prostanoids and their receptors. In *Comprehensive Medicinal Chemistry* Vol. 3 pp 643–714 Eds C Hansch, PG Sammes, JB Taylor and JC Emmett. Pergamon Press, Oxford

- Gatzuli E, Aten R and Behrman HR (1991) Inhibition of gonadotropin action and progesterone synthesis by xanthine oxidase in rat luteal cells *Endocrinology* **128** 2253–2258
- Gavrieli Y, Sherman Y and Ben-Sasson SA (1992) Identification of programmed cell death *in situ* via specific labeling of nuclear DNA fragmentation *Journal of Cell Biology* **119** 493–501
- Hansel W and Dowd JP (1986) New concepts of the control of corpus luteum function *Journal of Reproduction and Fertility* **78** 755–768
- Horton EW and Poyser NL (1976) Uterine luteolytic hormone: a physiological role for prostaglandin F_{2α} *Physiological Review* **56** 595–651
- Ikebuchi Y, Masumoto N, Tasaka K, Koike K, Kasahara K, Miyake A and Tanizawa O (1991) Superoxide anion increases intracellular pH, intracellular free calcium, and arachidonate release in human amnion cells *Journal of Biological Chemistry* **266** 13 233–13 237
- Ito S, Sakamoto K, Mochizuki-Oda N, Ezashi T, Miwa K, Okuda-Ashitaka E, Shevchenko VI, Kiso Y and Hayaishi O (1994) Prostaglandin F_{2α} receptor is coupled to Gq in cDNA-transfected Chinese hamster ovary cells *Biochemical and Biophysical Research Communications* **200** 756–762
- Juengel JL, Garverick HA, Johnson AI, Youngquist RS and Smith MF (1993) Apoptosis during luteal regression in cattle *Endocrinology* **132** 249–254
- Kerr JFR, Wyllie AH and Currie AR (1972) Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics *British Journal of Cancer* **26** 239–257
- Leung PCK, Minegishi T, Ma F, Zhou F and Ho-Yuen B (1986) Induction of polyphosphoinositide breakdown in rat corpus luteum by prostaglandin F_{2α} *Endocrinology* **119** 12–18
- McConkey DJ, Orrenius S and Jodal M (1990) Cellular signalling in programmed cell death (apoptosis) *Immunology Today* **11** 120–121
- McCracken JA and Schramm W (1988) Prostaglandins and corpus luteum regression. In *Prostaglandins: Biology and Chemistry of Prostaglandins and Related Eicosanoids* pp 425–462 Ed PB Curtis-Prior. Churchill Livingstone, London
- Olofsson J and Leung PCK (1994) Auto/paracrine role of prostaglandins in corpus luteum function *Molecular and Cellular Endocrinology* **100** 87–91
- Olofsson J and Selstam G (1988) Changes in corpus luteum content of prostaglandin F_{2α} and E in the adult pseudopregnant rat *Prostaglandins* **35** 31–40
- Priddy AR, Killick SR, Elstein M, Morris J, Sullivan M, Patel L and Elder MG (1989) Ovarian follicular fluid eicosanoid concentrations during the pre-ovulatory period in humans *Prostaglandins* **38** 197–202
- Rothchild I (1981) The regulation of the mammalian corpus luteum *Recent Progress in Hormone Research* **37** 183–298
- Sánchez-Criado JE, Ochiai K and Rothchild I (1987) Indomethacin treatment prevents prolactin-induced luteolysis in the rat *Journal of Endocrinology* **112** 317–322
- Sawada M and Carlson JC (1991) Rapid plasma membrane changes in superoxide radical formation, fluidity, and phospholipase A₂ activity in the corpus luteum of the rat during induction of luteolysis *Endocrinology* **128** 2992–2997
- Sawyer HR, Niswender KD, Braden TD and Niswender GD (1990) Nuclear changes in ovine luteal cells in response to PGF_{2α} *Domestic Animal Endocrinology* **7** 229–238
- Schmitz GG, Walter T, Seibl R and Kessler C (1991) Nonradioactive labeling of oligonucleotides *in vitro* with the hapten digoxigenin by tailing with terminal transferase *Analytical Biochemistry* **192** 222–231
- Sugimoto Y, Hasumoto K, Namba T, Irie A, Katsuyama M, Negishi M, Kakizuka A, Narumiya S and Ichikawa A (1994) Cloning and expression of a cDNA for mouse prostaglandin F receptor *Journal of Biological Chemistry* **269** 1356–1360
- Thiry M (1992) Highly sensitive immunodetection of DNA on sections with exogenous terminal deoxynucleotidyl transferase and non-isotopic nucleotide analogs *Journal of Histochemistry and Cytochemistry* **40** 411–419
- Tilly JL, Kowalski KI, Johnson AL and Hsueh AJW (1991) Involvement of apoptosis in ovarian follicular atresia and postovulatory regression *Endocrinology* **129** 2799–2801
- Wyllie AH, Kerr JFR and Currie AR (1980) Cell death: the significance of apoptosis *International Review of Cytology* **68** 251–305
- Zelevnik AJ, Ihrig LL and Bassett SG (1989) Developmental expression of Ca⁺⁺/Mg⁺⁺-dependent endonuclease activity in rat granulosa and luteal cells *Endocrinology* **125** 2218–2220