***In vitro* maturation, fertilization and embryo development after ultrarapid freezing of immature human oocytes**

J. Wu^{1*}, L. Zhang¹ and X. Wang¹

¹Department of Obstetrics and Gynecology, The Third Clinical Medical College, Beijing Medical University, Beijing 100083, China

The purpose of this study was to determine the rates of maturation, fertilization and embryo development of ultrarapidly frozen immature oocytes (immature cumulus–oocyte complexes; COCs) obtained from antral follicles in ovaries of patients with chocolate ovarian cysts. The COCs were cryopreserved by a vitrification method using 5.5 mol ethylene glycol l⁻¹ plus 1.0 mol sucrose l⁻¹ in Dulbecco's PBS (DPBS). The survival, maturation and fertilization rates, and the percentage of embryos developing to the two-cell stage were 59, 64, 70 and 71%, respectively. No significant differences were noted in the

rates of maturation, fertilization and embryo development between control and cryopreserved oocytes. Two embryos that developed from cryopreserved oocytes of the oocyte donor programme were selected for transfer into the uterus of a recipient with premature ovarian failure, after the recipient had received steroid replacement. A biochemical pregnancy occurred in the recipient after embryo transfer. These results indicate that immature oocytes can survive after cryopreservation and subsequently can be cultured to mature oocytes that are capable of undergoing fertilization *in vitro* and developing into embryos.

Introduction

The first mammalian oocytes to be frozen and thawed successfully were those of mice and hamsters (Parkening *et al.*, 1976; Tsunoda *et al.*, 1976; Whittingham, 1977). Successful procedures for freezing human oocytes (Chen, 1986; Van Uem *et al.*, 1978) were reported about 10 years later. The cryopreservation of human oocytes is an attractive alternative to embryo preservation in human *in vitro* fertilization (IVF) programmes and overcomes many of the ethical and legal objections to storage of human embryos (Capron *et al.*, 1992; Perry and Schneider, 1992). Cryopreservation could be used to circumvent iatrogenic sterility after chemo- or radiotherapy. Moreover, this technique would allow potential fertility in women suffering from pathologies of the reproductive system that compromise the function of the ovaries, such as premature ovarian failure, endometriosis, cysts and pelvic infections. The use of frozen oocytes would allow the maintenance of fertility in such patients. The cryopreservation of oocytes may allow for family planning for women who delay maternity due to career demands, the absence of a partner or because of pathologies that prevent pregnancy temporarily. Finally, the use of frozen oocytes might be included in a programme of oocyte donation.

Although there are case reports of successful pregnancies after cryopreservation of unfertilized oocytes (Chen, 1986; Van Uem *et al.*, 1978), cryopreservation of human oocytes is

still considered experimental, especially for immature human oocytes (Mandelbaum *et al.*, 1988; Toth *et al.*, 1994a,b; Son *et al.*, 1996). The purpose of this study was to determine rates of maturation, fertilization and embryo development after ultrarapid freezing of human immature oocytes.

Material and Methods

Collection of immature oocyte

The Third Clinical Medical College Beijing Medical University, Beijing, China, granted ethical approval of this study. Informed consent was obtained from each patient. Ovarian tissue was obtained from 36 patients undergoing gynaecological surgery for chocolate ovarian cysts on days 9–12 after their last menstrual period. Patients ranged in age from 20 to 45 years (mean 32.0 ± 5.8 years) and had regular menstrual cycles ranging from 27 to 33 days. These patients had not been exposed to exogenous hormones for 3 months before the surgery. Excised ovarian tissue was washed with saline at 35°C to remove adhering blood clots. Immature oocytes were aspirated by puncturing the ovarian follicles (4–10 mm in diameter) with a 21-gauge needle attached to a syringe filled with Dulbecco's PBS (DPBS, Gibco, Beijing). Only cumulus–oocyte complexes (COCs) with a compact cumulus cell mass were included in the study. COCs were randomized and either cryopreserved or cultured as controls.

Cryopreservation of immature oocyte

Immature oocytes were frozen using a vitrification method modified from Steponkus *et al.* (1990) and Martino *et al.* (1996). The vitrification solution used was composed

Email: ji.wu@hsc.utah.edu

*Present address: Division of Urology, University of Utah School of Medicine, 50 No. Medical Drive, Salt Lake City, UT 84132, USA.

of 5.5 mol ethylene glycol l⁻¹ and 1.0 mol sucrose l⁻¹ in DPBS (EG 5.5). Two or three COCs were exposed to 2 ml vitrification solution for 20 s and, using a pipette, were transferred in a very small volume of fluid on to electron microscope copper grids (3.05 mm in diameter; Electron Microscopy Sciences, Fort Washington, PA). The grids were plunged immediately into liquid nitrogen. Immature oocytes were stored in liquid nitrogen for 20–105 days.

Thawing of immature oocytes

Grids with vitrified oocytes were transferred as rapidly as possible from the liquid nitrogen into a culture dish containing 4 ml of 0.5 mol sucrose l⁻¹ at 37°C. After 10 s, the grids were transferred at intervals of 1 min into culture dishes containing sucrose solutions of 0.5, 0.25, 0.125 and 0.0625 mol l⁻¹, and finally into DPBS for 5 min. COCs were recovered from the grids in the DPBS solution using a pipette and were washed (3 × 5 min) in holding medium, Ham's F10 (Flow Laboratories, Irvine) with 10% fetal cord serum.

Oocyte maturation and fertilization, and embryo development

Oocytes that showed clear bright and homogeneous cytoplasm and an intact plasma membrane and zona pellucida were considered to have survived the freezing process. These oocytes were transferred to the maturation medium after thawing. The maturation medium was Ham's F10 supplemented with 0.75 iu human menopausal gonadotrophin ml⁻¹ (HMG; Organon, Oss) and 40% human follicular fluid (hFF). Vitrified and control oocytes were cultured at 37°C in 5% CO₂ in air for 32–56 h in four-well culture dishes (Nunc, Roskilde) containing the maturation medium. Control and frozen–thawed oocytes that extruded the first polar body were transferred to Ham's F10 with 10% fetal cord serum and inseminated with spermatozoa (0.1 ml spermatozoa at a concentration of 1 × 10⁶ ml⁻¹), which had been prepared using routine washing, centrifugation and swim-up techniques (Carrell *et al.*, 1998). The presence of two pronuclei at 16–19 h after insemination was used as an indication that fertilization had occurred. The ova were returned to the culture dishes for up to 144 h and the embryos were examined every 24 h. The stage of embryo development was determined on the basis of the number of blastomeres. When the blastomeres of a morula began to rearrange and become organized about a central cavity (containing fluid), inner cell mass and outer cell mass, the morula was considered to be a blastocyst.

Oocyte donor programme and embryo transfer

After the pre-clinical study, an oocyte donor programme was carried out. Immature oocytes were obtained on day 12 of the menstrual cycle from a patient with chocolate ovarian cysts who was 27 years old and had a regular menstrual cycle. The oocytes were cryopreserved (sterile electron

microscope copper grids, fresh liquid nitrogen and a sterile separate storage vessel were used) and fertilized with spermatozoa from the recipient's partner. Two embryos were transferred on day 18 of steroid replacement into the uterus of a 29-year-old recipient with premature ovarian failure. The exogenous steroid replacement was formulated to mimic the ovarian steroid profile during a normal 28 day menstrual cycle, and was the same as that described by Cha (1991). Urinary concentrations of hCG in the recipient were measured using an enzyme immunoassay, 16, 18 and 20 days after embryo transfer (Kobayashi *et al.*, 1983).

Statistical analysis

Chi-squared test was used for this statistical analysis.

Results

In the pre-clinical experiment, 157 immature oocytes were obtained from 36 patients. The mean number of immature oocytes collected per patient was 4.0 ± 1.9. At the time of cryopreservation, 61% of immature oocytes were at the germinal vesicle (GV) stage. Surviving oocytes had distinct intact plasma membranes (Fig. 1a)

Oocyte survival, maturation, fertilization and embryo development rates are presented (Table 1). No significant differences between control and cryopreserved oocytes were noted in rates of maturation, fertilization or embryo development (Fig. 1b,c,d). Overall, only one of 79 cryopreserved oocytes and two of 78 control oocytes reached the blastocyst stage. In the oocyte donor programme eight oocytes were collected, cryopreserved and subsequently inseminated (two-cell stage and four-cell stage). Four embryos were obtained and two were selected for transfer. A biochemical pregnancy occurred in the recipient after embryo transfer as determined by hCG in urine at days 16, 18 and 20.

Discussion

Most attempts to freeze human oocytes have involved conventional procedures similar to those developed for freezing embryos. These protocols include slow or direct cooling of the oocytes to the seeding temperature of about -7°C, holding the oocytes at this temperature for a few minutes before or after seeding, and cooling them at < 1°C min⁻¹ to an intermediate temperature of -30°C to -80°C before plunging them into liquid nitrogen (Son *et al.*, 1996; Trounson and Kirby, 1989; Toth *et al.*, 1994a,b). Vitrification or ultrarapid freezing of oocytes is a rapid simple technique compared with these conventional protocols. Vitrification circumvents the formation of potentially damaging intra- and extracellular ice, but cryoprotectant toxicity and osmotic injury to oocytes can occur. These effects depend on the choice and concentration of the cryoprotectant and the choice of protocol. Martino *et al.* (1996) succeeded in cryopreserving bovine

Table 1. Comparison of maturation, fertilization and cleavage rates between control and cryopreserved immature human oocytes

	Control	Cryopreserved
Number (n)	78	79
Morphological survival (%)	—	47 (59)
Maturation (%)	54 (69)	30 (64)
Fertilization (%)	43 (80)	21 (70)
Two-cell embryo (percentage of fertilization)	33 (77)	15 (71)
Three- to eight-cell embryo (percentage of two-cell embryos)	20 (61)	7 (47)
Early blastocyst (percentage of two-cell embryos)	2 (6)	1 (7)

Fig. 1. Development of immature human oocytes after ultrarapid cryopreservation. (a) Immature oocyte after thawing; (b) mature oocyte after thawing and *in vitro* maturation; (c) Two-cell embryo and (d) early blastocyst 144 h, respectively, after insemination of immature oocytes. Scale bars represent 25 μm.

oocytes using a vitrification solution containing only ethylene glycol and sucrose as cryoprotectants, and ultrarapid cooling. A significant improvement in oocyte survival was found when electron microscope grids were used as a physical support and oocytes were plunged

directly into liquid nitrogen, compared with cryopreserving oocytes in straws (Martino *et al.* 1996). Very high cooling rates can be achieved using these grids. The estimated cooling rate for *Drosophila* embryos cooled on copper grids and plunged directly into liquid nitrogen was 3000°C min⁻¹,

a rate about threefold higher than that obtained using straws (Steponkus *et al.*, 1990). Furthermore, the use of electron microscope grids for this protocol may help rapid heat conduction from the outside to prevent the oocyte from being damaged by the solution effect of the cryoprotectant.

In the present study, 59% of oocytes survived, 64% of the oocytes matured, 70% of the mature oocytes underwent fertilization and 71% of the fertilized oocytes cleaved. No significant differences in the rates of maturation, fertilization and embryo development were noted between control and cryopreserved oocytes.

In previous studies, 16–43% of oocytes collected from unstimulated ovaries survived freezing, and between 20 and 58% of these oocytes subsequently matured *in vitro* (Mandelbaum *et al.*, 1988; Toth *et al.*, 1994b). Son *et al.* (1996) reported that the survival rate of immature human oocytes after freezing and thawing was 55.1%; and maturation, fertilization and cleavage rates were 59.3, 42.9 and 16.7, respectively. The survival and maturation rates of immature oocytes in the present study are in agreement with the results of Son *et al.* (1996), but higher fertilization and embryo development rates were obtained in the present study compared with that of Son *et al.* (1996). This difference may be due to the use of very high cooling rates to circumvent the extreme chilling sensitivity of the oocyte.

Toth *et al.* (1994a) reported a maturation rate of 83.3%, a fertilization rate of 56.5%, and a blastocyst formation rate of 3.3% for cryopreserved immature human oocytes from stimulated IVF cycles. They suggested that the developmental capacity of immature oocytes from stimulated ovaries was higher than that of unstimulated ovaries. However, the rates of fertilization and embryo development of immature oocytes from unstimulated ovaries in the present study were similar to those from stimulated ovaries in the study of Toth *et al.* (1994a). In addition, it has been assumed that oocytes at prophase I collected after stimulation may be of reduced quality because immature oocytes collected after hyperstimulation demonstrate low rates of pregnancy after IVF and embryo transfer. Thus, additional work is required before conclusions can be reached regarding the developmental capacity of immature oocytes obtained from unstimulated and stimulated ovaries.

There is a risk of contamination when oocytes are brought into direct contact with liquid nitrogen. Tedder *et al.* (1995) reported that bone marrow samples were contaminated with hepatitis B virus during storage in liquid nitrogen. The virus was introduced into the refrigerator from a stored sample and was transferred to other samples either by failure of the integrity of the plastic sample containers used or by inadvertent contamination of the thawed sample contents from the surface of its storage bag. Procedures that avoid the risk of infection must be developed before ultrarapid oocyte cryopreservation is applied clinically.

The microtubular spindle of the mature oocyte is sensitive to temperature changes, and a potential benefit of cryopreserving oocytes at prophase I is that damage to the microtubule spindle may be avoided. In mice, oocytes at

prophase I did not demonstrate any major abnormalities in the meiotic spindle at the second metaphase after cryopreservation and maturation (Van der Elst *et al.*, 1992). Cryopreservation of immature oocytes at prophase I has been performed in mice (Schroeder *et al.*, 1990; Van der Elst *et al.*, 1992; Candy *et al.*, 1994), rats (Pellicer *et al.*, 1988) and humans (Mandelbaum *et al.*, 1988; Toth *et al.*, 1994a,b; Son *et al.*, 1996). Candy *et al.* (1994) obtained live normal offspring from cryopreserved immature mouse oocytes. There are few reports of pregnancy after cryopreservation of human oocytes at prophase I (Tucker *et al.*, 1998). Results from the current study confirm that immature human oocytes are able to survive cryopreservation and that surviving oocytes are able to mature to metaphase II, undergo fertilization *in vitro* and develop into early blastocysts. A biochemical pregnancy occurred in a recipient after embryo transfer. However, more detailed genetic screening and cytological analysis of the embryos obtained after maturation and fertilization of vitrified oocytes need to be undertaken before this method is applied clinically.

The authors wish to thank J. Riesen for the careful reading of the manuscript.

References

- Candy CJ, Wood MJ, Whittingham DG, Merriman JA and Choudhury N (1994) Cryopreservation of immature mouse oocytes *Human Reproduction* **9** 1738–1742
- Capron AM (1992) Parenthood and frozen embryos: more than property and privacy *Hastings Centre Reproduction* **22** 32–33
- Carrell DT, Kuneck PH, Peterson CM, Hatasaka HH, Jones KP and Campbell BF (1998) A randomized, prospective analysis of five sperm preparation techniques before intrauterine insemination of husband sperm *Fertility and Sterility* **69** 122–126
- Cha KY, Koo JJ, Ko JJ, Choi DH, Han SY and Yoon TK (1991) Pregnancy after *in vitro* fertilization of human follicular oocytes collected from nonstimulated cycles, their culture *in vitro* and their transfer in a donor oocyte program *Fertility and Sterility* **55** 109–113
- Chen C (1986) Pregnancy after human oocyte cryopreservation *Lancet* **1** 884–886
- Kobayashi O, Shirotake S and Takamizawa H (1983) Studies on enzymeimmunoassay for human chorionic gonadotrophin and its clinical application *Nippon Sanka Fujinka Gakkai Zasshi* **35** 764–772
- Mandelbaum J, Junca AM, Plachot M, Alnot MO, Salar Barolx J and Alvarez A (1988) Cryopreservation of human embryos and oocytes *Human Reproduction* **3** 117–119
- Martino A, Songsasen N and Leibo SP (1996) Development into blastocysts of bovine oocytes cryopreserved by ultra-rapid cooling *Biology of Reproduction* **54** 1059–1069
- Parkening TA, Tsunoda Y and Chang MC (1976) Effects of various low temperatures, cryoprotective agents and cooling rates on the survival, fertilize ability and development of frozen-thawed mouse eggs *Journal of Experimental Zoology* **197** 369–374
- Pellicer A, Lightman A, Parmer TG, Behrman HR and DeCherney AH (1988) Morphologic and functional studies of immature rat oocyte–cumulus complexes after cryopreservation *Fertility and Sterility* **50** 805–810
- Perry C and Schneider LK (1992) Cryopreserved embryos: who shall decide their fate? *Journal of Legal Medicine* **13** 463–500
- Schroeder AC, Champlin AK, Mabraaten LE and Eppig JJ (1990) Developmental capacity of mouse oocytes cryopreserved before and after maturation *in vitro*. *Journal of Reproduction and Fertility* **89** 43–50

- Son WY, Park SE, Lee KA, Lee WS, Ko JJ, Yoon TK and Cha KY (1996) Effects of 1,2-propanediol and freezing-thawing on the *in vitro* developmental capacity of human immature oocytes *Fertility and Sterility* **66** 995–999
- Steponkus PL, Myers SP, Lynch DC *et al.* (1990) Cryopreservation of *Drosophila melanogaster* embryos *Nature* **345** 170–172
- Tedder RS, Zuckerman MA, Goldstone AH *et al.* (1995) Hepatitis B transmission from contaminated cryopreservation tank *Lancet* **346** 137–140
- Toth TL, Baka SG, Veeck LL, Jones HW, Jr, Muasher S and Lanzendorf SE (1994a) Fertilization and *in vitro* development of cryopreserved human prophase I oocytes *Fertility and Sterility* **61** 891–894
- Toth TL, Lanzendorf SE, Sandow BA, Veeck LL, Hassen WA, Hansen K and Hodgen GD (1994b) Cryopreservation of human prophase I oocytes collected from unstimulated follicles *Fertility and Sterility* **61** 1077–1082
- Trounson A and Kirby C (1989) Problems in the cryopreservation of unfertilized eggs by slow cooling in dimethyl sulfoxide *Fertility and Sterility* **52** 778–786
- Tsunoda Y, Parkening TA and Chang MC (1976) *In vitro* fertilization of mouse and hamster eggs after freezing and thawing *Experientia* **32** 223–224
- Tucker MJ, Wright G, Morton PC and Massey JB (1998) Birth after cryopreservation of immature oocytes with subsequent *in vitro* maturation *Fertility and Sterility* **70** 578–579
- Van der Elst J, Nerinckx S and Van Steirteghem AC (1992) *In vitro* maturation of mouse germinal vesicle-stage oocytes following cooling, exposure to cryoprotectants and ultrarapid freezing: limited effect on the morphology of the second meiotic spindle *Human Reproduction* **7** 1440–1446
- Van Uem JF, Siebzehnubel ER, Schuh B, Koch R, Trotnow S and Lang N (1978) Birth after cryopreservation of unfertilized oocytes *Lancet* **1** 752–753
- Whittingham DG (1977) Fertilization *in vitro* and development to term of unfertilized mouse oocyte previously stored at –196°C *Journal of Reproduction Fertility* **49** 89–94

Resubmitted manuscript received 18 September 2000.

Accepted 9 October 2000.