Intraovarian actions of oestrogen

Cheryl S. Rosenfeld¹, Jessica S. Wagner², R. Michael Roberts^{1,3,4}
and Dennis B. Lubahn^{1,2,3,5}

Departments of ¹Animal Sciences, ²Molecular Biology Program, ³Biochemistry, ⁴Veterinary Pathobiology, ⁵Child Health, University of Missouri, Columbia, MO 65211, USA

Oestrogen regulates several hypothalamic and pituitary hormones, which in turn control ovarian functions. Oestrogen and its metabolites, such as catecholestrogens, also have direct effects within the ovary. This review examines the roles of oestrogen in regulating ovarian folliculogenesis, ovulation and corpus luteum formation. Oestrogen promotes follicular development, which culminates in ovulation, by potentiating follicular development, granulosa cell expression of gonadotrophin receptors, steroidogenesis, and gap junction formation by granulosa cells, and by inhibiting granulosa cell apoptosis. In addition, oestrogen may be needed for corpus luteum formation and maintenance. Studies on mutant mice that either lack one or both of the known oestrogen receptors or are unable to synthesize oestrogen support some but not all of these prior inferences of the roles of oestrogen within the ovary. Although these transgenic mice have proved useful in determining some of the intraovarian actions of oestrogen, they present confounding problems, including hormonal imbalances, that hinder interpretation. Transgenic mice with conditional or tissue-directed mutations in their oestrogen receptors are needed to dissect the ovarian actions of oestrogen further. In addition, microarray technologies, combined with specific hormone treatment regimens are likely to provide an attractive, alternative approach to using mutant mice in clarifying the direct actions of oestrogen in the ovaries of other species.

Since the discovery of oestrogens in the early 1920s (Allen and Doisy, 1923), their effects have intrigued reproductive biologists. Classically, two approaches have been used to study the role of oestrogen within the ovary: surgical removal of the pituitary gland to eliminate confounding gonadotrophins, and pharmacological blockage of either the synthesis of oestrogen or its receptors both *in vivo* and *in vitro*. Although some studies have implicated a role for oestrogen within the ovary (Goldenberg *et al.*, 1972; Nakayama *et al.*, 1981; Nakano *et al.*, 1982; Gore-Langton and Daniel, 1990; Nayudu and Osborn, 1992; Hulshof *et al.*, 1995), others have not (Coney *et al.*, 1987; Spears *et al.*, 1998). Transgenic gene knockout technology has been used to investigate the intraovarian effects of oestrogen. Mice that lack either the P450aromatase gene (Fisher *et al.*, 1998; Honda *et al.*, 1998) or one or both of the known oestrogen receptors have been produced (Lubahn *et al.*, 1993; Krege *et al.*, 1998; Couse *et al.*, 1999; Dupont *et al.*, 2000; for review, see Rosenfeld *et al.*, 2001). The main ovarian events studied for oestrogen involvement have been folliculogenesis, steroidogenesis, ovulation and corpus luteum formation.

Oestrogen synthesis, sites of action and metabolism

Androgens are produced by the thecal cells and, in general, they are taken up by the granulosa cells and converted by P450aromatase to various oestrogens with the main one being oestradiol (Fig. 1). In some species, such as pigs (Lautincik *et al.*, 1994; Shores and Hunter, 1999), chickens (Kato *et al.*, 1995) and tree shrews (Kimura *et al.*, 2000), the thecal cells can also synthesize oestrogen. This steroid hormone is then released into the vasculature and circulated to the uterus, hypothalamus, mammary gland and various other reproductive and non-reproductive organs. Oestrogens may also exert local effects within the ovary. Some oestrogens appear also to be metabolized into presumed inactive as well as active products, such as the catecholestrogens (Zhu and Conney, 1998; Fig. 1).

Catecholestrogens

The enzymes, 2-hydroxylase (cyp1a1) and 4-hydroxylase (cyp1b1) catalyse the formation of catecholestrogens from oestradiol (Fig. 1). Both of these catecholestrogens can bind to the two known oestrogen receptors, ER α and ER β (Kuiper *et al.*, 1997). In addition, it has been proposed that they bind to a novel oestrogen receptor (Das *et al.*, 1997) and possibly

Email: RosenfeldC@missouri.edu

Fig. 1. Synthesis of oestradiol and catechoestrogens. Testosterone is converted by P450aromatase into oestrogen (oestradiol). Oestradiol can then be metabolized into inactive and active metabolites, such as 2-hydroxyoestradiol and 4-hydroxyoestradiol. The enzymes 2-hydroxylase and 4-hydroxylase catalyse the conversion of oestradiol to 2-hydroxyoestradiol and 4-hydroxyoestradiol, respectively. All of these enzymes are expressed by the granulosa cells.

even androgen receptor (Hudson and Hillier, 1985). Pig and human ovaries contain high concentrations of 2-hydroxylase and 4-hydroxylase (Hammond *et al.*, 1986; Muskhelishvili *et al.*, 2001), and certain ovarian responses appear to be induced directly by these steroid metabolites, as discussed further below. The pattern of expression of these enzymes has not been examined throughout the oestrous cycle.

Oestrogen receptors

Oestrogen was the first steroid hormone demonstrated to activate transcription by binding to a receptor, the oestrogen receptor (for review, see Jensen and DeSombre, 1973). The receptors with their bound hormone modulate transcription by binding to oestrogen response elements (ERE) present on oestrogen-inducible genes (Kumar *et al.*, 1987). The receptors can also act as co-regulators through binding to other transcription factors already attached to gene regulatory regions (Adler *et al.*, 1988; Feng *et al.*, 1998), and by ligand-independent mechanisms (Ignar-Trowbridge *et al.*, 1992). Although it was first believed that oestrogen receptors were predominantly cytoplasmic, it was later proposed that they are located exclusively within the nucleus (Welshons *et al.*, 1984; Greene *et al.*, 1986). It is now clear that these receptors can be found in both the cytoplasm and nucleus, with the ratio dependent upon the cell type and physiological conditions (Yamashita, 1998; Zieba *et al.*, 2000). Some studies also indicate that oestrogen receptors reside on the plasma membrane and modulate cellular activity without directly associating with DNA (Morley *et al.*, 1992; Levin, 1999; Pietras and Szego, 1999; Nadal *et al.*, 2000).

The oestrogen receptors are members of the superfamily of steroid nuclear receptors (Parker, 1995; Baker, 1997; White and Parker, 1998). All steroid receptors have a DNA-binding domain that is composed of two zinc fingers followed by a C-terminal ligand-binding domain (Parker, 1995; White and Parker, 1998). Currently, two oestrogen receptors, ERα (Green *et al.*, 1986; Greene *et al.*, 1986) and ERβ (Kuiper *et al.*, 1996; Mosselman *et al.*, 1996) have been cloned. These two receptors have considerable sequence identity in the DNA-binding domains, which permit both receptor types to interact with EREs of various genes. Sequence differences between the two receptors occur primarily in the N- and C-terminal regions (Kuiper *et al.*, 1996; Mosselman *et al.*, 1996).

There are many alternatively spliced forms of both ERα and ERβ (Chu and Fuller, 1997; Petersen *et al.*, 1997; Kos *et al.*, 2000; Poola *et al.*, 2000). The alternatively spliced forms of ERα have been identified predominantly in human breast cancer and other tumour cell lines (Kos *et al.*, 2000; Poola *et al.*, 2000) but not in normal ovarian cells. One alternative form of ERβ identified in rats lacks exon 4, which forms the second zinc finger of the DNA-binding domain, whereas another form has 18 additional codons in the ligand-binding domain, which results in a protein with only 10% of the oestradiol-binding affinity and 100- to 1000-fold lower transcriptional activation potential compared with the full length receptor (Petersen *et al.*, 1997). This form with 18 additional amino acids appears to be the predominant ERβ in various mouse organs, including the ovary (Lu *et al.*, 2000). A third form of ERβ described in rats carries a deletion of the second zinc finger, as well as the 18 additional codons in the ligand-binding domain (Petersen *et al.*, 1997) (Fig. 2). All three of these aforementioned forms have been identified in

Fig. 2. Alternatively spliced forms of oestrogen receptor β (ER β) identified in the rat ovary. Spliced forms that have a deletion of the second finger, which contributes to binding of the oestrogen responsive element on DNA, are presumably unable to induce transcription through the oestrogen response element. Other forms that have insertion of amino acids within the ligand-binding domain may have different binding affinities to the various endogenous oestrogens.

the rat ovary (Petersen *et al.*, 1997), and thus, each may regulate oestrogenic activity in the rodent ovary. In addition, other novel human ER β forms, which differ in their C-terminal sequences, have been identified, and some of these are expressed in the human ovary (Moore *et al.*, 1998).

The discovery of ER β raises the question as to whether mammals possess additional oestrogen receptors that have yet to be identified. No such genes have been identified to date in the human genome, although there is some evidence for at least one additional oestrogen receptor in the mouse ovary and uterus (Kudolo *et al.*, 1984a,b; Hillier *et al.*, 1989; Das *et al.*, 1997; Ghosh *et al.*, 1999; for review, see Rosenfeld *et al.*, 2001). Moreover, a third oestrogen receptor, ER γ , has been identified in fish (Hawkins *et al.*, 2000). ER γ is distinct from other fish ER α s and ER β s, but is quite closely related to ER β . In mammals, alternatively spliced forms of ER α and ER β that differ in their ligand-binding domains may represent novel oestrogen receptors

because these mutations might alter their ability to interact with various endogenous oestrogens, their metabolites and antagonists.

For oestrogens to act on specific cell types in the ovary, ovarian oestrogen receptors must be present, and oestrogen-induced gene activation generally occurs. The ovarian cell types in which ER α and ER β have been identified are listed (Table 1). In all species studied to date, ER β is the most abundant ovarian oestrogen receptor. This receptor is especially prominent in granulosa cells, but it is also expressed in luteal cells and, to a lesser extent, in thecal cells (see references in Table 1).

Regulation of ovarian oestrogen receptor expression

As in other oestrogen-responsive organs, such as the uterus, oestrogen downregulates granulosa cell expression of ER β protein (Sharma *et al.*, 1999). In rodents, treatment

Table 1. Expression of oestrogen receptor α and β in ovarian cells of various species, as determined by *in situ* hybridization or immunohistochemistry

Ovarian cells	Oestrogen receptor α			Oestrogen receptor β		
	Expression	Species	Reference	Expression	Species	Reference
Granulosa cells of pre-antral follicles	Not detected	Marmoset	Saunders <i>et al.</i> , 2000	High	Human	Pelletier and El-Alfy, 2000;
		Human	Pelletier and El-Alfy, 2000;			Saunders <i>et al.</i> , 2000;
		Rat	Saunders <i>et al.</i> , 2000			Taylor and Al-Azzawi, 2000
Granulosa cells of antral follicles	Low	Human	Sar and Welsch, 1999	High	Rat	Byers <i>et al.</i> , 1997;
						Sar and Welsch, 1999;
		Marmoset	Pelletier <i>et al.</i> , 2000		Marmoset	Pelletier <i>et al.</i> , 2000
			Saunders <i>et al.</i> , 2000			Saunders <i>et al.</i> , 2000
			Taylor and Al-Azzawi, 2000			Pelletier and El-Alfy, 2000;
Theca cells of antral follicles	Low	Human	Saunders <i>et al.</i> , 2000;	Low	Human	Saunders <i>et al.</i> , 2000;
			Taylor and Al-Azzawi, 2000			Taylor and Al-Azzawi, 2000
		Rat	Saunders <i>et al.</i> , 2000		Marmoset	Taylor and Al-Azzawi, 2000
			Pelletier and El-Alfy, 2000			Saunders <i>et al.</i> , 2000
			Saunders <i>et al.</i> , 2000			Rosenfeld <i>et al.</i> , 1999
Luteal cells	Low-moderate	Human	Sar and Welsch, 1999	Not detected	Rat	Sar and Welsch, 1999
			Pelletier and El-Alfy, 2000			Saunders <i>et al.</i> , 2000;
		Marmoset	Saunders <i>et al.</i> , 2000;		Human	Taylor and Al-Azzawi, 2000;
			Taylor and Al-Azzawi, 2000			Saunders <i>et al.</i> , 2000;
			Saunders <i>et al.</i> , 2000			Hosokawa <i>et al.</i> , 2001
Germinal epithelium	Low	Human	Taylor and Al-Azzawi, 2000	Not detected	Rat	Saunders <i>et al.</i> , 2000
			Telleria <i>et al.</i> , 1998			Pelletier and El-Alfy, 2000;
		Marmoset	Zieba <i>et al.</i> , 2000		Human	Sar and Welsch, 1999
			Saunders <i>et al.</i> , 2000			Pelletier and El-Alfy, 2000;
			Saunders <i>et al.</i> , 2000			Saunders <i>et al.</i> , 2000

with gonadotrophins, in particular LH, results in a marked decrease of ER β in the ovary (Byers *et al.*, 1997; Fitzpatrick *et al.*, 1999). This effect is mediated through a cAMP-dependent pathway. Not surprisingly, addition of cAMP to cultured rat granulosa cells also decreases ER β expression (O'Brien *et al.*, 1999). In contrast, when cattle are infused with ovulatory dosages of LH, ovarian ER β concentrations are not altered (Manikkam *et al.*, 1999). Why this species difference occurs is not clear, but it may relate to the manner in which follicular maturation and selection occurs in the two species, that is, polyovulatory rodents versus generally monovulatory cattle. ER β may regulate early follicular growth in rodents, whereas it may exert control over both early and late follicular development in cattle.

Mice that lack LH-hCG receptors have a marked decrease in ovarian ER α mRNA, but an increase in ovarian ER β mRNA expression, as determined by northern blot analysis (Lei *et al.*, 2001). Oestrogen and progesterone treatment of these mice restores ER α mRNA to the amounts found in wild-type mice, but ER β mRNA remains increased. These findings support the contention that LH signalling is needed in mice to maintain proper ovarian ER β mRNA concentrations. Concomitantly, the absence of ER α results in increased serum concentrations of LH (Rissman *et al.*, 1997; Couse and Korach, 1999), and thus, ER α is needed to maintain normal LH concentrations.

Intraovarian effects of oestrogen

Folliculogenesis

As the ovarian follicles grow and differentiate, increasing amounts of oestrogen are produced (Fortune, 1994), which, in turn, upregulate the synthesis and release of the pituitary gonadotrophins, FSH and LH, thereby promoting ovarian follicular growth. On the basis of some reports, oestrogen directly increases the number and size of ovarian follicles *in vivo* and the size of rat, mouse and bovine follicles in culture (Goldenberg *et al.*, 1972; Nakayama *et al.*, 1981; Nakano *et al.*, 1982; Gore-Langton and Daniel, 1990; Nayudu and Osborn, 1992; Hulshof *et al.*, 1995). However, other studies on rat and ovarian follicles have not supported this finding (Coney *et al.*, 1987; Spears *et al.*, 1998).

If oestrogen is indeed directly promoting ovarian follicular growth, it is uncertain which oestrogen receptors are involved. Both the ER α knockout (ER α KO) (Rosenfeld *et al.*, 2000) and ER β KO (Krege *et al.*, 1998) mice exhibit follicular development to the Graafian stage, albeit to a lesser extent than in wild-type sibling mice. However, ER $\alpha\beta$ double KO mice are infertile because of follicular arrest (Couse *et al.*, 1999; Dupont *et al.*, 2000). Nevertheless, early follicular growth and development occurs in these mice, even though mature Graafian follicles do not form. Collectively, these results indicate that, in mice, ER α or ER β can compensate, at least partially, for the absence of the other to provide mature ovulatory follicles, but may not be necessary for the earlier stages of follicular growth.

Aromatase knockout (ArKO) mice can complete folliculogenesis without exogenous oestrogens, but the process is impaired, with fewer follicles reaching maturity than in wild-type controls (Fisher *et al.*, 1998; Britt *et al.*, 2000). ArKO mice that are treated with oestradiol every 4 days from 4 weeks of age for 1 month have increased numbers of follicles compared with untreated ArKO mice. However, these treated mice still do not develop corpora lutea (Toda *et al.*, in press). On the basis of these data, folliculogenesis can occur, albeit sub-normally, in the absence of either oestrogen or one of its two known receptors.

Gonadotrophin receptor expression

Growing follicles express gonadotrophin receptors. Before selection, follicles express FSH receptors, but LH receptors emerge later in the dominant follicles. In cattle, when the dominant follicles are first selected, FSH reaches its nadir (Fortune, 1994) while serum concentrations of LH begin to increase. It is presumed that only those follicles that express LH receptors in granulosa cells can be rescued and, in time, ovulated.

Oestrogen increases follicular expression of both FSH and LH receptors in rat granulosa cells (Richards *et al.*, 1976, 1979). Those follicles that first begin to produce significant amounts of oestrogen are more likely to possess more gonadotrophin receptors. Therefore, whether or not a particular follicle expresses oestrogen receptors may control its fate. However, ER α KO mice have increased expression of granulosa and thecal LH receptor (Schomberg *et al.*, 1999), again indicating a lack of ER α involvement in control of follicular growth. These mice also have increased serum oestrogen concentrations, which may increase granulosa cell expression of LH receptor by acting through ER β or one of the putative additional receptor types. There is currently no information on the ovarian expression of FSH and LH receptors in ER β KO and ER $\alpha\beta$ double KO mice. If oestrogen acts through ER β to upregulate gonadotrophin receptor expression, ER β mutant mice presumably would have decreased expression of FSH and LH receptors.

Steroid production

Ovarian steroid hormones are produced throughout the oestrous cycle. During folliculogenesis, the thecal cells produce androgens, which are converted into oestrogens by P450aromatase in the granulosa cells. Both oestradiol (Fortune and Hansel, 1979; Leung and Armstrong, 1980; Welsh *et al.*, 1983; Roberts and Skinner, 1990) and catechol-oestrogens (Spicer and Hammond, 1987; Tekpetey and Armstrong, 1994) can regulate the production of androgen and progesterone within bovine, rat and pig ovaries. Oestrogen is known to increase CYP17 expression by rat theca cells (Johnson and Crane, 1995). The oestrogen-driven increase in progesterone presumably promotes corpus luteum formation and maintenance through luteal progesterone receptors (Duffy and Stouffer, 1995; Smith *et al.*, 1995).

Promotes folliculogenesis

Increases expression of gonadotrophin receptors by granulosa cells

Increases gap junction formation by granulosa cells

Increases steroid synthesis

Inhibits granulosa cell apoptosis

Gap junctions

As the follicles continue to grow and proliferate, the avascular granulosa cell lining becomes removed from the nutrient-providing interstitial blood vessels. Gap junctions permit transfer of nutrients and cytokines to and from the granulosa cells and developing oocytes (Albertini and Anderson, 1974; Anderson and Albertini, 1976). Oestrogen controls granulosa cell gap junction formation (Merk *et al.*, 1972; Burghardt and Anderson, 1981). Gap junctions are composed of various connexin proteins: connexin 43 is a major granulosa and luteal gap junction protein in bovine (Nuttinck *et al.*, 2000), ovine (Grazul-Bilska *et al.*, 1998), pig (Lenhart *et al.*, 1998) and rat (Mayerhofer and Garfield, 1995) ovaries, and is oestrogen-regulated (Yu *et al.*, 1994). One method whereby oestrogen might potentiate gap junction formation between granulosa cells is through induction of this protein. However, connexin 43 protein expression has not been reported in any of the oestrogen receptor or P450arom mutant female mice.

Apoptosis

Most follicles do not reach the ovulatory stage but instead become atretic owing to apoptosis of the granulosa cell lining (Kaipia and Hsueh, 1997; Amsterdam *et al.*, 1999). This process is normally tightly regulated with anti- and pro-apoptotic factors balancing each other, and with crosstalk among multiple intracellular pathways determining the final outcome of the cell (Kaipia and Hsueh, 1997; Amsterdam *et al.*, 1999). Oestrogen inhibits granulosa cell apoptosis (Billig *et al.*, 1993). In contrast, androgens promote apoptosis (Billig *et al.*, 1993).

As with other ovarian processes, it is uncertain which oestrogen receptors are mediating this oestrogenic anti-apoptotic effect. In addition, it is not clear whether there are physiological repercussions if the anti-apoptotic mechanism is perturbed. In 10–12-week-old ArKO mice, many apoptotic granulosa cells are present in large antral follicles (Britt *et al.*, 2000), and by 21–23 weeks of age, these mice have fewer antral follicles than do wild-type controls. Predictably, ArKO mice have increased ovarian expression of pro-apoptotic genes, such as *p53* and *Bax*, compared with wild-type mice (Toda *et al.*, in press). At 1 year of age, the ArKO mice have a reduced number of primary follicles, all of which contain numerous apoptotic cells (Britt *et al.*, 2000). By contrast, ER α KO mice have similar numbers of apoptotic granulosa cells when compared with wild-type control mice (Schomberg *et al.*, 1999). Although oestrogen may be essential for preventing the demise of ovarian follicles, this protection is clearly not conferred through ER α .

Corpus luteum formation and maintenance

Early studies with hypophysectomized or X-irradiated rabbits and rats with hypothalamic lesions indicated that oestrogen directly regulates corpus luteum formation and controls luteal maintenance (Robson, 1937; Bogdanove, 1966; Keyes and Nalbandov, 1967). Unexpectedly, ER β KO mice demonstrate normal corpus luteum development, but have an overall reduced fertility (Krege *et al.*, 1998). Pre-pubertal ER α KO mice treated with superovulatory dosages of gonadotrophins also develop steroidogenically functional corpora lutea, but they remain infertile (Rosenfeld *et al.*, 2000). Thus, the phenotypes of these mutant mice do not corroborate earlier studies. In contrast, ArKO and ER $\alpha\beta$ double KO mice do not undergo luteinization and corpus luteum formation (Fisher *et al.*, 1998; Couse *et al.*, 1999; Britt *et al.*, 2000; Dupont *et al.*, 2000). Taken together, these findings indicate that oestrogen is indeed needed for corpus luteum formation and maintenance, and that it likely acts non-selectively through ER α or ER β with one oestrogen receptor compensating for the absence of the other. If both receptors are lacking, oestrogen cannot stimulate luteinization.

Non-genomic actions of oestrogen within the ovary

The complex physiological responses to oestrogen are generally presumed to involve transcriptional regulation of many genes. However, oestrogen-binding to putative membrane oestrogen receptors (Levin, 1999) may also induce non-genomic–non-transcriptionally mediated responses. In cultured granulosa cells, oestrogen has been implicated in causing a rapid increase in intracellular Ca²⁺ concentrations (Morley *et al.*, 1992). As this response occurs within seconds of treating the cells with oestrogen, a reasonable conclusion is that it is occurring without transcriptional activation, that is, it is elicited through some existing membrane-associated response. The demonstration that ER α associates with the regulatory subunit of phosphatidylinositol-3-OH kinase (PI(3)K) in the presence of ligand may provide at least a partial explanation for the non-genomic effects of oestrogen (Simoncini *et al.*, 2000), and thus provide another dimension to oestrogenic action within the ovary. Non-genomic responses have not yet been studied in transgenic mice.

Confounding problems in interpreting ovarian phenotypes in oestrogen- and oestrogen receptor-deficient mice

Although these knockout mice have proved invaluable in elucidating some of the roles of oestrogen within the ovary,

Fig. 3. Illustration of the proposed intraovarian actions of oestrogen. On the basis of past pharmacological, surgical, and culture approaches, oestrogen modulates many ovarian responses, including promoting folliculogenesis, increasing granulosa cell gonadotrophin receptor expression, increasing gap junction formation between granulosa cells, increasing steroid production by both thecal and luteal cells, and inhibiting granulosa cell apoptosis. E₂: oestrogen; LH-R: LH receptor, FSH-R: FSH receptor.

confounding factors hinder interpretation of the resulting ovarian phenotypes. In rodents, only one P450aromatase gene has been identified and knocked out (Youngblood *et al.*, 1989; Fisher *et al.*, 1998; Honda *et al.*, 1998). However, such mice are likely to have been exposed *in utero* to maternal oestrogen and postnatally to other sources of oestrogen, including compounds in the environment and diet, which would hinder interpreting the reproductive phenotypes of the adult animals. Furthermore, it is now apparent that there are non-aromatized steroid molecules, such as hermaphrodiol, that have both androgenic and oestrogenic activity (Lima *et al.*, 2000; Rosenberg-Zand *et al.*, 2000). Therefore, proper interpretation of ovarian phenotypes in ArKO mice requires exclusion of all sources of both aromatic and non-aromatic oestrogenic molecules. In addition, ArKO mice have secondary hormonal imbalances, such as increased serum concentrations of LH, FSH and androgen, which may account for some of the ovarian pathology (Fisher *et al.*, 1998).

The ERKO mice also have secondary hormonal imbalances, including an increase in serum LH, which is present in both ER α KO (Rissman *et al.*, 1997; Couse and Korach, 1999) and ER $\alpha\beta$ double KO (Couse *et al.*, 1999a) mice. In ERKO mice, all receptors that bind oestrogen have to be considered and deleted before it is possible to make accurate inferences about which oestrogen receptors mediate the intraovarian effects of oestrogen. Novel oestrogen receptors may be discovered once the mouse and human genomic sequences are fully annotated and the properties of forms arising by alternative splicing elucidated. It is interesting to note that the orphan oestrogen-related receptors (ERR α , β , and γ), the endogenous ligands of which are unknown, have been demonstrated, albeit with low affinity, to bind diethylstilboestrol (DES) but not oestradiol (Tremblay *et al.*, 2001). Thus, the ERRs may play a role in oestrogenic actions within the ovary, although as yet these receptors have not been examined fully in the ovary.

Ultimate resolution of these difficulties may depend on the creation of more sophisticated approaches to producing mutant mice that have ovarian cell-specific and conditional ablation of oestrogen receptors. Production of such mice would permit a thorough analysis of the effects of oestrogen-binding to oestrogen receptors throughout the oestrous cycle and pregnancy and would have minimal confounding hormonal imbalances. Transgenic mice with tamoxifen-dependent Cre recombinase fused to a mutated ligand-binding domain of human oestrogen receptor have been produced (Metzger and Chambon, 2001) and, when these mice are treated with tamoxifen, the chromosomally integrated gene, which is flanked by loxP sites, is excised.

Studying the intraovarian actions of oestrogen in other species

It is worth considering whether the mouse is the best model with which to study the intraovarian effects of oestrogen. Is it possible, for example, to extrapolate from responses seen in

rodents to other species? Although women, cows and mares are generally monovulatory, rodents are polyovulatory, and thus, the regulation of folliculogenesis may be different. How then can the *in vivo* effects of oestrogen in the ovaries of other species be studied, where the targeted deletion of genes is not currently possible? Microarray technology could be invaluable in this respect and provide comparison of oestrogen-regulated genes in various ovarian cells during puberty, the oestrous cycle, pregnancy and in response to oestrogenic treatment. This technique could be used in combination with pharmacological blockage of one or more oestrogen receptors to provide multiple snapshots of the intraovarian effects of oestrogen. An ovarian gene database (<http://ovary.stanford.edu>; Melner and Korach, 2000) has been established that may assist in the identification of all the oestrogen-regulated ovarian genes.

Regulation of oestrogen action within the ovary

Although there is ample evidence that oestrogen acts within the ovary, it also originates there, and is present in highest concentrations in this organ. Such quantities would in theory continually saturate and activate any cognate receptors, and potentially result in either continual transcriptional activation or other consequences such as desensitization. Some regulation of this system must occur to prevent such scenarios. One regulatory step could be at the receptor level. Oestrogen receptors may be required for only a limited period of follicular development but persist thereafter in a sub-functional state.

Co-activators may provide another level of regulation. These factors act in concert with oestrogen receptors to induce transcription of oestrogen-responsive genes. Various transcription factors, such as SRC-1, GRIP1, RAC3, p300, SPA AIB1, RIP140, SMRT, Fas-associated protein-tyrosine phosphatase-1 (FAP-1), and co-activator independent of AF-2 function (CIA) that regulate oestrogen-responsive genes have been identified (Rey *et al.*, 2000; Hlaing *et al.*, 2001; Sauve *et al.*, 2001). Possibly, it is their specific expression pattern and not that of the cognate receptors that governs the oestrogen responsiveness of granulosa and other ovarian cells. Differential expression of some of the oestrogen receptor co-activators supports this hypothesis (Hlaing *et al.*, 2001). Although granulosa, thecal and stromal cells in ovine ovaries express SRC-1, RIP140 and SPA mRNA, granulosa cells have the highest mRNA expression for these co-activators (Hlaing *et al.*, 2001). Similarly, the ovine corpus luteum expresses smaller amounts of SRC-1 and RIP140 mRNA than do follicular cells (Hlaing *et al.*, 2001).

A third form of regulation may be through metabolism of oestrogen by the *cyp* enzymes into various active and inactive metabolites. Compounds, such as the catechoestrogens, may bind oestrogen receptors (Kuiper *et al.*, 1997), androgen receptor (Hudson and Hillier, 1985) and other novel steroid receptors (Das *et al.*, 1997, 2000), resulting in responses both quantitatively and qualitatively distinct from the parental steroid. How efficiently the various ovarian cells convert

oestradiol into these metabolites might provide another level of control.

In summary, selective expression of oestrogen receptors, co-activational factors, and metabolic conversion of oestradiol may regulate the intraovarian effects of oestrogen. Similar processes could be invoked in other tissues in which a hormone acts within its originating tissue. For example, progesterone receptors are found in the bovine and primate corpus luteum (Smith *et al.*, 1995; Duffy and Stouffer, 1995).

Conclusions

The intraovarian effects of oestrogen have been examined. Early work indicated that oestrogen directly regulates ovarian folliculogenesis, corpus luteum formation, granulosa cell expression of the gonadotrophin receptors, gap junction formation between granulosa cells, granulosa cell apoptosis and steroid production by granulosa, thecal and luteal cells (Fig. 3). These actions are presumably regulated by specific oestrogen receptors. ER β is the predominant form of oestrogen receptor found in the ovary (Byers *et al.*, 1997; Couse *et al.*, 1997).

Aromatase-deficient and oestrogen receptor-deficient mice have been created to study the effects of oestrogen. Although results from these mice support a role for oestrogen in the ovary, some of the data are equivocal, such that the phenotypes cannot always be directly attributed to the absence of the gene, since secondary hormonal imbalances and other confounding factors invariably occur in these mice. More sophisticated, tissue-targeted knockout mice are needed to elucidate the roles of oestrogen within the ovary. The involvement of oestrogen in ovarian physiology needs to be examined in several species before any firm and general conclusions can be drawn about its intraovarian effects.

The authors wish to thank Alison Murray, Norah Spears, Michael F. Smith, Mohan Manikkam, Michele Calder, H. Allen Garverick, and members of Dennis Lubahn's and R. Michael Roberts' laboratories for their assistance. These studies were supported by the following grants: NIH ES 08272 (D. B. Lubahn), Army DAMD 17-97-1-7171 (D. B. Lubahn), EPA R825295 (D. B. Lubahn), and a USDA National Needs Fellowship (to C. S. Rosenfeld).

References

Key references are identified by asterisks.

- Adler S, Waterman ML, He X and Rosenfeld MG** (1988) Steroid receptor-mediated inhibition of rat prolactin gene expression does not require the receptor DNA-binding domain *Cell* **52** 685–695
- Albertini DF and Anderson E** (1974) The appearance and structure of intercellular connections during the ontogeny of the rabbit ovarian follicle with particular reference to gap junctions *Journal of Cell Biology* **63** 234–250
- Allen E and Doisy EA** (1923) An ovarian hormone. Preliminary report on its localization, extraction, and purification, and action in test animals *Journal of the American Medical Association* **81** 819–821
- Amsterdam A, Gold RS, Hosokawa K, Yoshida Y, Sasson R, Jung YS and Kotsuji F** (1999) Crosstalk among multiple signaling pathways controlling ovarian cell death *Trends in Endocrinology and Metabolism* **10** 255–262
- Anderson E and Albertini DF** (1976) Gap junctions between the oocytes and companion follicle cells in the mammalian ovary *Journal of Cell Biology* **71** 680–686
- Baker ME** (1997) Steroid receptor phylogeny and vertebrate origins *Molecular Cellular Endocrinology* **135** 101–107
- Billig H, Furuta I and Hsueh AJW** (1993) Estrogens inhibit and androgens enhance ovarian granulosa cell apoptosis *Endocrinology* **133** 2204–2212
- Bogdanove EM** (1966) Preservation of functional corpora lutea in the rat by estrogen treatment *Endocrinology* **79** 1011–1015
- *Britt KL, Drummond AE, Cox VA, Dyson M, Wreford NG, Jones MEE, Simpson ER and Findlay JK** (2000) An age-related ovarian phenotype in mice with targeted disruption of the *Cyp 19* (Aromatase) gene *Endocrinology* **141** 2614–2623
- Burghardt RC and Anderson E** (1981) Hormonal modulation of gap junctions in rat ovarian follicles *Cell Tissue Research* **214** 181–193
- Byers M, Kuiper GGJM, Gustafsson J-A and Park-Sarge O-K** (1997) Estrogen receptor- β mRNA expression in rat ovary: downregulation by gonadotropins *Molecular Endocrinology* **11** 172–182
- Chu S and Fuller PJ** (1997) Identification of a splice variant of the rat estrogen receptor beta gene *Molecular Cellular Endocrinology* **132** 195–199
- Coney P, Yoshimura Y, Hosoi Y, Bongiovanni A and Wallach E** (1987) Effect of aromatase inhibitors on the histology of cycling rat ovary *Gynecological Obstetrics Investigations* **23** 177–183
- Couse JF and Korach KS** (1999) Estrogen receptor null mice: what have we learned and where will they lead us? *Endocrine Reviews* **20** 358–417
- Couse JF, Lindzey J, Grandien K, Gustafsson J-A and Korach KS** (1997) Tissue distribution and quantitation analysis of estrogen receptor- α (ER α) and estrogen receptor- β (ER β) messenger ribonucleic acid in the wild-type and ER- α knockout mouse *Endocrinology* **138** 4613–4621
- *Couse JF, Curtis-Hewitt S, Bunch DO, Sar M, Walker VR, Davis BJ and Korach KS** (1999) Postnatal sex reversal of the ovaries in mice lacking estrogen receptors α and β *Science* **286** 2328–2331
- *Das SK, Taylor JA, Korach KS, Paria BC, Dey SK and Lubahn DB** (1997) Estrogenic responses in estrogen receptor- α deficient mice reveal a distinct estrogen signaling pathway *Proceedings of the National Academy of Sciences USA* **94** 12 786–12 791
- Das SK, Tan J, Raja S, Halder J, Paria BC and Dey SK** (2000) Estrogen targets genes involved in protein processing, calcium homeostasis, and Wnt signaling in the mouse uterus independent of estrogen receptor- α and - β *Journal of Biological Chemistry* **275** 28 834–28 842
- Duffy DM and Stouffer RL** (1995) Progesterone receptor messenger ribonucleic acid in the primate corpus luteum during the menstrual cycle: possible regulation by progesterone *Endocrinology* **136** 1869–1876
- *Dupont S, Krust A, Gansmuller A, Dierich A, Chambon P and Mark M** (2000) Effect of single and compound knockouts of estrogen receptors α (ER α) and β (ER β) on mouse reproductive phenotypes *Development* **127** 4277–4291
- Feng W, Ribeiro RC, Wagner RL, Nguyen H, Apriletti JW, Fletterick RJ, Baxter JD, Kushner PJ and West BL** (1998) Hormone-dependent coactivator binding to a hydrophobic cleft on nuclear receptors *Science* **280** 1747–1749
- *Fisher CR, Graves KH, Parlow AF and Simpson ER** (1998) Characterization of mice deficient in aromatase (ArKO) because of targeted disruption of the *cyp19* gene *Proceedings of the National Academy of Sciences USA* **95** 6965–6970
- Fitzpatrick SL, Funkhouser JM, Sindoni DM, Stevis PE, Deecher DC, Bapat AR, Merenthaler I and Frail DE** (1999) Expression of estrogen receptor-beta protein in rodent ovary *Endocrinology* **140** 2581–2591
- Fortune JE** (1994) Ovarian follicular growth and development in mammals *Biology of Reproduction* **50** 225–232
- Fortune JE and Hansel W** (1979) The effects of 17 β -estradiol on progesterone secretion by bovine theca and granulosa cells *Endocrinology* **104** 1834–1838
- Ghosh D, Taylor JA, Green JA and Lubahn DB** (1999) Methoxychlor

- stimulates estrogen-responsive messenger ribonucleic acids in mouse uterus through a non-estrogen receptor (Non-ER) α and Non-ER β mechanism *Endocrinology* **140** 3526–3533
- Goldenberg RL, Vaitukaitis JL and Ross GT (1972) Estrogen and follicle-stimulating hormone interactions on follicle growth in rats *Endocrinology* **90** 1492–1498
- Gore-Langton RE and Daniel SAJ (1990) Follicle-stimulating hormone and estradiol regulate antrum-like reorganization of granulosa cells in rat preantral follicle cultures *Biology of Reproduction* **43** 65–72
- Grazul-Bilska AT, Redmer DA, Bilski JJ, Jablonka-Shariff A, Doraiswamy V and Reynolds LP (1998) Gap junctional proteins, connexin 26, 32, and 43 in sheep ovaries throughout the estrous cycle *Endocrine* **8** 269–279
- Green S, Walter P, Kumar V, Krust A, Bornert J-M, Argos P and Chambon P (1986) Human oestrogen receptor cDNA: sequence, expression, and homology to v-erb-A *Nature* **320** 134–139
- Greene GL, Gilna P, Waterfield M, Baker A, Hort Y and Shine J (1986) Sequence and expression of human estrogen receptor complementary DNA *Science* **231** 1150–1154
- Hammond JM, Hersey RM, Walega MA and Weisz J (1986) Catecholestrogen production by porcine ovarian cells *Endocrinology* **118** 2292–2299
- Hawkins MB, Thornton JW, Crews D, Skipper JK, Dotte A and Thomas P (2000) Identification of a third distinct estrogen receptor and reclassification of estrogen receptors in teleosts *Proceedings of the National Academy of Sciences USA* **97** 10 751–10 756
- Hillier SG, Saunders PTK, White R and Parker MG (1989) Oestrogen receptor mRNA and a related transcript in mouse ovaries *Journal of Molecular Endocrinology* **2** 39–45
- Hlaing M, Nam K, Lou J, Pope WF and Nephew KP (2001) Evidence for expression of estrogen receptor cofactor messenger ribonucleic acid in the ovary and uterus of domesticated animals (sheep, cow and pig) *Life Sciences* **68** 1427–1438
- Honda S, Harada N, Ito S, Takagi Y and Maeda S (1998) Disruption of sexual behavior in male aromatase-deficient mice lacking exons 1 and 2 of the cyp19 gene *Biochemical Biophysical Research Communications* **252** 445–449
- Hosokawa K, Ottander U, Wahlberg P, Ny T, Cajander S and Olofsson IJ (2001) Dominant expression and distribution of oestrogen receptor β over oestrogen receptor α in the human corpus luteum *Molecular Human Reproduction* **7** 137–145
- Hudson KE and Hillier SG (1985) Catechol oestradiol control of FSH-stimulated granulosa cell steroidogenesis *Journal of Endocrinology* **106** R1–R4
- Hulshof SCJ, Figueiredo JR, Beckers JF, Beyers MM, van der Donk JA and van den Hurk R (1995) Effects of fetal bovine serum, FSH, and 17 β -estradiol on the culture of bovine preantral follicles *Theriogenology* **44** 217–226
- Ignar-Trowbridge DM, Nelson KG, Bidwell MC, Curtis SW, Washburn TF, McLachlan JA and Korach KS (1992) Coupling of dual signaling pathways: epidermal growth factor action involves the estrogen receptor *Proceedings of the National Academy of Sciences USA* **89** 4658–4662
- Jensen EV and DeSombre ER (1973) Estrogen–receptor interaction *Science* **182** 126–134
- Johnson DC and Crane LH (1995) Inhibitory and stimulatory effects of oestrogens upon 17 α -hydroxylase/C17,20-lyase in immature hypophysectomized rats treated with gonadotrophin *Journal of Endocrinology* **145** 59–67
- Kaipia A and Hsueh AJW (1997) Regulation of ovarian follicular atresia *Annual Review of Physiology* **59** 349–363
- Kato M, Shimada K, Saito N, Noda K and Ohta M (1995) Expression of P450 17 α -hydroxylase and P450aromatase genes in isolated granulosa, theca interna, and theca externa layers of chicken ovarian follicles during follicular growth *Biology of Reproduction* **52** 405–410
- Keyes PL and Nalbandov AV (1967) Maintenance and function of corpora lutea in rabbits depend on estrogen *Endocrinology* **80** 938–946
- Kimura J, Tsukise A, Watanabe G, Taya K, Rerkamnuaychoke W, Endo H, Kurohmaru M, Yamada J and Nishida T (2000) Immunohistochemical localization of inhibin and steroidogenic enzymes in ovary of common tree shrew (*Tupaia glis*) and northern smooth-tailed shrew (*Dendrogale murina*) *Anatomical Histological Embryology* **29** 267–271
- Kos M, O'Brien S, Flouriot G and Gannon F (2000) Tissue-specific expression of multiple mRNA variants of the mouse estrogen receptor α gene *FEBS Letters* **477** 15–20
- *Krege JH, Hodgin JB, Couse JF, Enmark E, Warner M, Mahler JF, Sar M, Korach KS, Gustafsson J-A and Smithies O (1998) Generation and reproductive phenotypes of mice lacking estrogen receptor β *Proceedings of the National Academy of Sciences USA* **95** 15 677–15 682
- Kudolo GB, Elder MG and Myatt L (1984a) A novel oestrogen-binding species in rat granulosa cells *Journal of Endocrinology* **102** 83–91
- Kudolo GB, Elder MG and Myatt L (1984b) Further characterization of the second oestrogen-binding species of the rat granulosa cell *Journal of Endocrinology* **102** 93–102
- *Kuiper GGJM, Enmark E, Peltö-Huikko M, Nilsson S and Gustafsson J-A (1996) Cloning of a novel estrogen receptor expressed in rat prostate and ovary *Proceedings of the National Academy of Sciences USA* **93** 5925–5930
- Kuiper GGJM, Carlsson B, Grandien K, Enmark E, Häggblad J, Nilsson S and Gustafsson J-Å (1997) Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors α and β *Endocrinology* **138** 863–870
- Kumar V, Green S, Stack G, Berry M, Jin JR and Chambon P (1987) Functional domains of the human estrogen receptor *Cell* **51** 941–951
- Lautincik J, Kolodzieyski L, Elias V, Hyttel P, Osawa Y and Sirotkin A (1994) Immunocytochemical localization of aromatase in the ovary of superovulated cattle, pigs, and sheep *Acta Veterinaria Scandinavica* **35** 185–191
- Lei ZM, Mishra S, Zou W, Xu B, Foltz M, Li X and Rao ChV (2001) Targeted disruption of luteinizing hormone/human chorionic gonadotropin receptor gene *Molecular Endocrinology* **15** 184–200
- Lenhart JA, Downer BR, Bagnell CA (1998) Connexin 43 gap junction protein expression during follicular development in the porcine ovary *Biology of Reproduction* **58** 583–590
- Leung PCK and Armstrong DT (1980) Further evidence in support of a short-loop feedback action of estrogen on ovarian androgen production *Life Sciences* **27** 415–420
- Levin ER (1999) Cellular functions of the plasma membrane estrogen receptor *Trends in Endocrinology and Metabolism* **10** 374–377
- Lima PH, Hotchkiss JR and Blomquist CH (2000) Characterization of 3 α -, 3 β -, and 17 β -hydroxysteroid oxidoreductase, 5 α -reductase, and aromatase activity in adult rabbit lung *Endocrinology Annual Meeting, Toronto Abstract* 1546
- Lu B, Leygue E, Dotzlaw H, Murphy LJ and Murphy LC (2000) Functional characteristics of a novel estrogen receptor- β isoform, estrogen receptor- β 2 *Journal of Endocrinology* **25** 229–242
- *Lubahn DB, Moyer JS, Golding TS, Couse JF, Korach KS and Smithies O (1993) Alteration of reproductive function but not prenatal sexual development after insertional disruption of the mouse estrogen receptor gene *Proceedings of the National Academy of Sciences USA* **90** 11 162–11 166
- Manikkam M, Bao B, Rosenfeld CS, Yuan X, Salfen BE, Calder MD, Youngquist RS, Keisler DH, Lubahn DB and Garverick HA (1999) Expression of bovine estrogen receptor- β (bER β) mRNA throughout the first follicular wave in the ovary and effect of LH infusion on bER β mRNA expression of second wave follicles *Society for the Study of Reproduction Annual Meeting, Pullman, WA Abstract* 52
- Mayerhofer A and Garfield RE (1995) Immunocytochemical analysis of the expression of gap junction protein connexin 43 in the rat ovary *Molecular Reproduction and Development* **41** 331–338
- Melner MH and Korach KS (2000) A new database for tissue-specific gene expression and function in the ovary *Endocrinology* **141** 3051
- Merk FB, Botticelli CR and Albright JT (1972) An intracellular response to estrogen by granulosa cells in the rat ovary; an electron microscopic study *Endocrinology* **99** 992–1007
- Metzger D and Chambon P (2001) Site- and time-specific gene targeting in the mouse *Methods* **24** 71–80
- Moore JT, McKee DD, Slentz-Kesler K, Moore LB, Jones SA, Horne EL, Su JL, Kliewer SA, Lehmann JM and Willson TM (1998) Cloning and characterization of human estrogen receptor beta isoforms *Biochemical Biophysical Research Communications* **247** 75–78

- Morley P, Whitfield JF, Vanderhyden BC, Tsang BK and Schwartz JL (1992) A new, nongenomic estrogen action: the rapid release of intracellular calcium *Endocrinology* **131** 1305–1312
- Mosselman S, Polman J and Dijkema R (1996) ER β : identification and characterization of a novel human estrogen receptor *FEBS Letters* **392** 49–53
- Muskhelishvili L, Thompson PA, Kusewitt DF, Wang C and Kadlubar FF (2001) *In situ* hybridization and immunohistochemical analysis of cytochrome P450 1B1 expression in human normal tissues *Journal of Histochemistry and Cytochemistry* **49** 229–236
- Nadal A, Ropero AB, Laribi O, Maillet M, Fuentes E and Soria B (2000) Nongenomic actions of estrogens and xenoestrogens by binding at a plasma membrane receptor unrelated to estrogen receptor alpha and estrogen receptor beta *Proceedings of the National Academy of Sciences USA* **97** 11 603–11 608
- Nakano R, Nakayama T and Iwao M (1982) Inhibition of ovarian follicle growth by a chemical antiestrogen *Hormone Research* **16** 230–236
- Nakayama T, Nakano R and Iwao M (1981) Evidence for a role of endogenous estrogen on follicular growth in immature female rats *Endocrinology Japan* **28** 715–723
- Nayudu PL and Osborn SM (1992) Factors influencing the rate of preantral and antral growth of mouse ovarian follicles *in vitro*. *Journal of Reproduction and Fertility* **95** 349–362
- Nuttinck F, Peynot N, Humblot P, Massip A, Dessy F and Flechon JE (2000) Comparative immunohistochemical distribution of connexin 37 and connexin 43 throughout folliculogenesis in the bovine ovary *Molecular Reproduction Development* **57** 60–66
- O'Brien ML, Park K, In Y and Park-Sarge OK (1999) Characterization of estrogen receptor- β (ER β) messenger ribonucleic acid and protein expression in rat granulosa cells *Endocrinology* **140** 4530–4541
- Parker MG (1995) Structure and function of estrogen receptors *Vitamins Hormones* **51** 267–287
- Peitras RJ and Szego CM (1999) Cell membrane estrogen receptors resurface *Nature Medicine* **5** 1130–1131
- Pelltier G and El-Alfy M (2000) Immunocytochemical localization of estrogen receptors α and β in the human reproductive organs *Journal of Endocrinology and Metabolism* **85** 4835–4840
- Pelletier G, Labrie C and Labrie F (2000) Localization of oestrogen receptor α , oestrogen receptor β and androgen receptors in the rat reproductive organs *Journal of Endocrinology* **165** 359–370
- Petersen DN, Tkalecic GT, Koza-Taylor PH, Turi TG and Brown TA (1997) Identification of estrogen receptor β 2, a functional variant of estrogen receptor β expressed in normal rat tissues *Endocrinology* **139** 1082–1092
- Poola I, Koduri S, Chatra S and Clarke R (2000) Identification of twenty alternatively spliced estrogen receptor α mRNAs in breast cancer cell lines and tumors using splice targeted primer approach *Journal of Steroid Biochemistry and Molecular Biology* **72** 249–258
- Rey JM, Pujol P, Callier P, Cavailles V, Freiss G, Maudelonde T and Brouillet JP (2000) Semiquantitative reverse transcription-polymerase chain reaction to evaluate the expression patterns of genes involved in the oestrogen pathway *Journal of Molecular Endocrinology* **24** 433–440
- Richards JS, Ireland JJ, Rao MC, Bernath GA, Midgley AR, Jr and Reichert LE, Jr (1976) Ovarian follicular development in the rat: hormone receptor regulation by estradiol, follicle stimulating hormone, and luteinizing hormone *Endocrinology* **99** 1562–1570
- Richards JS, Jonassen JA, Rolfes AI, Kersey K and Reichert LE, Jr (1979) Adenosine 3',5'-monophosphate, luteinizing hormone receptor, and progesterone during granulosa cell differentiation: effects of estradiol and follicle stimulating hormone *Endocrinology* **104** 765–773
- Rissman EF, Wersinger SR, Taylor JA and Lubahn DB (1997) Estrogen receptor function as revealed by knockout studies: neuroendocrine and behavioral aspects *Hormones and Behavior* **31** 232–243
- Roberts AJ and Skinner MK (1990) Estrogen regulation of theca cell steroidogenesis and differentiation: thecal cell-granulosa cell interactions *Endocrinology* **127** 2918–2929
- Robson JM (1937) Maintenance by oestrin of the luteal function in hypophysectomized rabbits *Journal of Physiology* **90** 435–439
- Rosenberg-Zand RS, Diamandis EP and Stanczyk FZ (2000) Hermaphrodite may contribute significantly to the estrogenic/androgenic hormonal milieu of breast and prostrate cancer cells *Endocrinology Annual Meeting, Toronto Abstract* 773
- Rosenfeld CS, Yuan X, Manikkam M, Calder MD, Garverick HA and Lubahn DB (1999) Cloning, sequencing, and localization of bovine estrogen receptor- β within the ovarian follicle *Biology of Reproduction* **60** 691–697
- *Rosenfeld CS, Murray AA, Simmer G, Hufford MG, Smith MF, Spears N and Lubahn DB (2000) Gonadotropin induction of ovulation and corpus luteum formation in young estrogen receptor- α knockout mice *Biology of Reproduction* **62** 599–605
- *Rosenfeld CS, Roberts RM and Lubahn DB (2001) Estrogen receptor- and aromatase-deficient mice provide insight into the roles of estrogen in the ovary and uterus *Molecular Reproduction and Development* **59** 336–346
- Sar M and Welsch F (1999) Differential expression of estrogen receptor- β and estrogen receptor- α in the rat ovary *Endocrinology* **140** 963–971
- Saunders PTK, Millar MR, Williams K, Macpherson S, Harkiss D, Anderson RA, Orr B, Groome NP, Scobie G and Fraser HM (2000) Differential expression of estrogen receptor- α and - β and androgen receptor in the ovaries of marmosets and humans *Biology of Reproduction* **63** 1098–1105
- Sauve F, McBroom LD, Gallant J, Moraitis AN, Labrie F and Giguere V (2001) CIA, a novel estrogen receptor coactivator with a bifunctional nuclear receptor interacting determinant *Molecular Cellular Biology* **21** 343–353
- Schomberg DW, Couse JF, Mukherjee A, Lubahn DB, Sar M, Mayo KE and Korach KS (1999) Targeted disruption of the estrogen receptor- α gene in female mice: characterization of ovarian responses and phenotype in the adult *Endocrinology* **140** 2733–2744
- Sharma SC, Clemens JW, Pisarska MD and Richards JS (1999) Expression and function of estrogen receptor subtypes in granulosa cells: regulation by estradiol and forskolin *Endocrinology* **140** 4320–4334
- Shores EM and Hunter MG (1999) Immunohistochemical localization of steroidogenic enzymes and comparison with hormone production during follicle development in the pig *Reproduction Fertility and Development* **11** 337–344
- Simoncini T, Hafezi-Moghadam A, Brazil DP, Ley K, Chin WW and Liao JK (2000) Interaction of oestrogen receptor with the regulatory subunit of phosphatidylinositol-3-OH kinase *Nature* **407** 538–541
- Smith GW, Gentry PC, Kong DK, Bao B, Roberts RM and Smith MF (1995) Expression of progesterone receptor mRNA within ovine post-surge follicles and corpora lutea *28th Annual Meeting of the Society for the Study of Reproduction, Davis, CA Abstract* 380
- Spears N, Murray AA, Allison V, Boland NI and Gosden RG (1998) Role of gonadotrophins and ovarian steroids in the development of mouse follicles *in vitro*. *Journal of Reproduction and Fertility* **113** 19–26
- Spicer LJ and Hammond JM (1987) Catecholestrogens stimulate progesterin secretion by cultured porcine granulosa cells *Molecular and Cellular Endocrinology* **50** 139–147
- Taylor AH and Al-Azzawi F (2000) Immunolocalisation of oestrogen receptor β in human tissues *Journal of Molecular Endocrinology* **24** 145–155
- Tekpetey FR and Armstrong DT (1994) Catecholestrogen modulation of steroid production by rat luteal cells: mechanism of action *Molecular and Cellular Endocrinology* **101** 49–57
- Telleria CM, Zhong L, Deb S, Srivastava RK, Park KS, Sugino N, Park-Sarge OK and Gibori G (1998) Differential expression of the estrogen receptors α and β in the rat corpus luteum of pregnancy: regulation by prolactin and placental lactogens *Endocrinology* **139** 2432–2442
- Toda K, Takeda K, Okada T, Akira S, Saibara T, Kaname T, Yamamura K, Onishi S and Shizuta Y Targeting disruption of the aromatase P450 gene (*Cyp19*) in mice and their ovarian and uterine responses to 17 β -oestradiol *Journal of Endocrinology* (in press)
- Tremblay GB, Kunath T, Bergeron D, Lapointe L, Champigny C, Bader JA, Rossant J and Giguere V (2001) Diethylstilbestrol regulates trophoblast stem cell differentiation as a ligand of orphan nuclear receptor ERR β *Genes and Development* **15** 833–838
- Welsh TH, Jr, Zhuang L-Z and Hsueh AJW (1983) Estrogen augmentation of gonadotropin-stimulated progesterin biosynthesis in cultured rat granulosa cells *Endocrinology* **112** 1916–1924

- Welshons WV, Lieberman ME and Gorski J** (1984) Nuclear localization of unoccupied oestrogen receptors *Nature* **307** 747–749
- *White R and Parker MG** (1998) Molecular mechanisms of steroid hormone action *Endocrine Related Cancer* **5** 1–14
- Yamashita S** (1998) Localization and functions of steroid hormone receptors *Histological Histopathology* **13** 255–270
- Youngblood GL, Nesbitt MN and Payne AH** (1989) The structural genes encoding P450scc and P450arom are closely linked on mouse chromosome 9 *Endocrinology* **125** 2784–2786
- Yu W, Dahl G and Werner R** (1994) The connexin43 gene is responsive to oestrogen *Proceedings of the Royal Society of London B Biological Sciences* **22** 125–132
- Zhu BT and Conney AH** (1998) Functional role of estrogen metabolism in target cells – review and perspectives *Carcinogenesis* **19** 1–27
- Zieba D, Bilinska B, Schmalz-Fraczek B and Murawski M** (2000) Immunohistochemical localization of estrogen receptors in the ovine corpus luteum throughout the estrous cycle *Folia Histochemistry Cytobiology* **38** 111–117