Dissociation of oocyte nuclear and cytoplasmic maturation by the addition of insulin in cultured bovine antral follicles

A A Fouladi-Nashta and K H S Campbell¹

Roslin Institute, Roslin, Midlothian, Edinburgh EH13 9HN, UK and ¹School of Biosciences, University of Nottingham, Sutton Bonington, Loughborough, Leicestershire LE12 5RD, UK

Correspondence should be addressed to A A Fouladi-Nashta; Email: ali.fouladi-nashta@nottingham.ac.uk

A A Fouladi-Nashta is now at School of Biosciences, University of Nottingham, Sutton Bonington, Loughborough, Leicestershire LE12 5RD, UK

Abstract

Follicles of 4–8 mm in diameter were dissected from ovaries and cultured in Waymouth culture medium in the presence or absence of insulin (5 µg/ml) at 39 °C in a humidified atmosphere of 45% O₂, 5% CO₂ and 50% N₂ for 24 h. Following follicle culture, the oocytes were collected and examined for developmental potential, total protein profile and ultrastructural aspects. Oocytes aspirated directly from follicles of the same size were used as controls. Addition of insulin to the follicle culture medium significantly reduced expression of the low molecular weight insulin-like growth factor-binding proteins (IGFBPs) in the follicular fluid, and significantly reduced the cleavage rate of subsequently matured and fertilised oocytes (0.52 vs 0.61). However, there were no differences in the proportion of cleaved embryos which developed to the blastocyst stage (0.30 vs 0.28), nor embryo quality as assessed by total cell number (137 ± 8.53 vs 124.6 ± 6.95). The total protein profiles of immature oocytes recovered after 24 h of follicle culture were compared by PAGE. There were marked differences between the two groups, unmaturing oocytes recovered from the insulin-positive follicle group showed a protein pattern similar to that of matured oocytes. In addition, examination of ultrastructural features by transmission electron microscopy indicated that oocytes from follicles cultured in the presence of insulin undergo many of the cytoplasmic changes associated with oocyte maturation. In conclusion, follicle culture in the presence of insulin is beneficial for follicular survival and significantly reduces cleavage but has no detrimental effects on the development of cultured embryos. However, many of the cytoplasmic changes associated with oocyte maturation occur prior to the induction of nuclear maturation.

Reproduction (2006) **131** 449–460

Introduction

The growth and development of ovarian follicles, oocytes and embryos depend upon the spatial and temporal expression of specific genes and on a continuous supply of energy, hormones and growth factors. Insulin, which is primarily involved in the regulation of glucose concentration in the circulation, also regulates cell growth and development in a wide variety of cell types (Starus 1981, Hsueh *et al.* 1984). The importance of insulin as a regulator of ovarian function was revealed in patients with altered insulin concentrations. Insulin-dependent diabetes mellitus in women is commonly accompanied by ovarian hypo-function and amenorrhoea, whilst hyper-insulinaemia is associated with ovarian hyperstimulation and hyperandrogenism (Poretsky & Kalin 1987, Poretsky 1991, Fulghesu *et al.* 1997). Mouse models of maternal hypoin-sulinaemia and hyperglycaemia produce oocytes and developing ovarian follicles of smaller sizes, reduced percentage of germinal vesicle breakdown and an increased

occurrence of apoptosis (Chang *et al.* 2005). In cattle, high insulin concentrations in non-lactating heifers were associated with increased numbers of small follicles (<4 mm) (Gong *et al.* 1993a) and an increased response to superovulatory treatment (Gong *et al.* 1993b).

In vitro, insulin has repeatedly been reported to stimulate proliferation and steroidogenesis of granulosa and theca cells (Campbell *et al.* 1995, Duleba *et al.* 1997) and is routinely included in tissue and cell culture media. In porcine (May *et al.* 1980) and human (Hill & Osteen 1992) ovarian tissue, insulin stimulates granulosa cell progesterone secretion and granulosa cell luteinisation (Channing *et al.* 1976) and induces granulosa cell luteinising hormone (LH)/human chorionic gonadotrophin (hCG) receptors (May *et al.* 1980). It also increases rat granulosa cell oestrogen and progestin production in response to follicle-stimulating hormone (FSH) (Davoren & Hsueh 1984).

A role for insulin in oocyte maturation has been described. Insulin induced meiotic maturation in *Xenopus*

laevis oocytes. Tsafiri & Channing (1975) demonstrated that insulin stimulates maturation of pig oocytes beyond first metaphase and extrusion of the first polar body. Subsequently, Lessman & Schuetz (1981) reported that insulin facilitates germinal vesicle breakdown in leopard frogs (*Rana pipiens*). In cattle, mRNA encoding receptor for insulin has been detected at all embryonic stages from the one-cell zygote to the blastocyst (Schultz *et al.* 1992). However, insulin does not improve bovine oocyte maturation (Betteridge *et al.* 1989) and has no effect on the fertilisation rate or subsequent development (Stubblings *et al.* 1990). In the mouse, protein synthesis increases in the presence of insulin at the compacted morulae stage of development (Rao *et al.* 1990, Kaye *et al.* 1992) when the insulin receptor appears (Harvey & Kaye 1988).

The insulin-like growth factors (IGFs) are small proteins, structurally related to proinsulin, that stimulate growth and differentiation of a wide variety of cell types (Jones & Clemmons 1995). In the circulation and body fluids, IGFs are complexed with specific binding proteins (IGFBPs) and association of IGFs with their binding proteins in the circulation prolongs their half-life by decreasing their clearance rate (Jones & Clemmons 1995).

The IGFBP content of ovarian follicular fluid from a number of species has been examined using Western ligand blots and immunoblots. For example, in the cow, binding proteins of molecular mass 40–44 kDa (IGFBP-3), 34 kDa (IGFBP-2), 29–27 kDa (IGFBP-5) and 22 kDa (IGFBP-4) have been identified (Echternkamp *et al.* 1994). The actions of IGFs in ovarian cell cultures have been shown to be inhibited by IGFBPs (Mason *et al.* 1992, Monget *et al.* 1993). A decrease in follicular IGFBP production would therefore be expected to increase the biological activity of locally produced IGF and thus increase the sensitivity of the follicle to gonadotrophins. The observed changes in the concentration of IGFBPs in follicular fluid during follicle development agree with this hypothesis and indicate that IGFBPs are key autocrine and paracrine regulators of ovarian function.

We have previously reported a system for the culture of bovine intact antral follicles *in vitro* (Fouladi-Nashta *et al.* 1998). Oocytes recovered following 24 h of follicle culture remained at the germinal vesicle (GV) stage and gave a significantly greater frequency of development compared with oocytes aspirated from fresh follicles, following subsequent maturation, fertilisation and culture. Insulin is generally added to tissue culture including ovarian follicles as a survival/mitogenic factor in high concentrations. Use of supra-physiological concentrations of insulin may be beneficial for the survival of ovarian follicular somatic cells, but may have a detrimental effect on oocyte quality and development. The aims of the experiments reported here were to elucidate the effects of addition of insulin to the follicle culture medium on follicular survival, via the evaluation of embryonic development and the quality of oocytes following 24 h of follicle culture. The effect of insulin on the ultrastructure of immature oocytes derived following

follicle culture in the presence of insulin was examined. In addition, the effects of insulin on IGFBP and total protein profiles in follicles and oocytes were investigated.

Materials and Methods

All chemicals and reagents were purchased from Sigma (Poole, Dorset, UK) unless otherwise stated.

Isolation and culture of antral follicles

Ovaries were obtained from local abattoirs and maintained at 28–35 °C during transport to the laboratory (1–2 h). Using a pair of scissors and forceps, follicles of 4–8 mm diameter were dissected from the surrounding connective tissues and cultured as previously described (Fouladi-Nashta *et al.* 1998). Briefly, non-atretic follicles were selected on the basis of morphological criteria, including translucency, lack of free particles in the antrum and the presence of blood vessels in the follicular wall (Kruip & Dieleman 1982).

Selected follicles were cultured in six-well culture dishes fitted with netwell inserts of 500 µm pore diameter containing Waymouth medium MB752/1 (Invitrogen, Paisley, Strathclyde, UK) supplemented with 0.23 mM pyruvic acid, 50 mg/l streptomycin sulphate, 75 mg/l penicillin G and 3 mg/ml BSA (pH 7.4, 280 mOsmol/kg H₂O) in the presence or absence (control) of 5 µg/ml insulin. A gaseous atmosphere of 45% O₂, 50% N₂ and 5% CO₂ was used according to Moor *et al.* (1973). Before use, the culture medium was placed in an anaerobic jar which was gassed for 5 min, sealed and warmed to 39 °C in an incubator. After transferring the follicles into the culture dishes, they were replaced in the anaerobic jar which was gassed for 5 min before being sealed and returned to the incubator.

Oocyte collection

Cumulus–oocyte complexes (COCs) were aspirated from follicles 4–8 mm in diameter using a 10 ml syringe fitted with a hypodermic needle (1.2 mm internal diameter). The aspirated follicular fluid was placed into sterile universal containers (Nunc, Roskilde, Denmark), in a warm air chamber (39 °C) and allowed to settle for 10–15 min. The majority of the follicular fluid was then removed by surface aspiration and the remaining follicular material diluted with an equal volume of dissection medium (TCM199 with Earl's salts (Invitrogen, Paisley, UK) 75.0 mg/l kanamycin, 7.08 g/l Hepes (pH 7.8, osmolarity 279 mOsmol/kg H₂O) supplemented with 10% fetal calf serum (FCS). The diluted follicular fluid was transferred into an 85 mm Petri dish (Bibby Sterilin, Stone, UK) and examined for COCs under a dissecting microscope.

Oocytes from follicle culture treatments were collected after opening the follicles with forceps. After 24 h of follicle culture, COCs from both groups ($n = 106$ in insulin-positive and $n = 98$ in insulin-negative groups) were

collected and washed in dissection medium as previously described.

Oocyte maturation

COCs isolated following 24 h of follicle culture were washed twice in dissection medium and transferred into 500 µl pre-gassed maturation medium (TCM199 with Earl's salts, 75 mg/l kanamycin, 4.75 g/l HEPES, 2.29 g/l NaHCO₃ (pH 7.8, osmolarity 280 mOsmol/kg H₂O) supplemented with 0.006 IU/ml FSH/LH (Pergonal; Serono Laboratories, Herts, UK) and 10% FCS in 35 mm tissue culture dishes. COCs were then incubated in a humidified atmosphere of 5% CO₂ in air at 39°C for 24 h.

In vitro fertilisation

In vitro matured oocytes were fertilised as previously described (Fouladi-Nashta *et al.* 1998). Briefly, COCs were gently pipetted in order to remove adhering granulosa cells and break up aggregated COCs. Disaggregated COCs were then washed once in oocyte wash medium containing 6.8 g/l NaCl, 0.23 g/l KCl, 168 mg/l NaHCO₃, 47 mg/l Na₂HPO₄, 4.8 g/l HEPES, 75 mg/l kanamycin monosulphate, 11 mg/l pyruvic acid, 6 g/l BSA, 1.86 ml/l 60% syrup lactic acid, 100 mg/l MgCl₂·6H₂O, 0.84 g/l CaCl₂·2H₂O (pH 7.4, osmolarity 282 mOsmol/kg H₂O) and transferred into 45 µl microdrops of fertilisation medium (five to ten oocytes/drop) containing sperm (1.5×10^6 /ml) and maintained for 24 h at 39°C in a humidified incubator in 5% CO₂ in air.

In vitro embryo culture

At 46–48 h after co-incubation of the spermatozoa and oocytes, cleaved embryos with at least four cells were selected, washed twice in a HEPES-buffered synthetic oviductal fluid (SOF) medium and transferred to 20 µl droplets (5 µl/embryo) of SOF medium (Thompson *et al.* 1991) supplemented with 4 mg/ml Pentex crystalline BSA (Bayer, Elkhart, Germany) overlaid with mineral oil. Embryo culture was carried out in 35 mm cell culture dishes at 39°C, in a humidified incubator with a gaseous atmosphere of 5% CO₂ and 5% O₂.

Total cell counting

Blastocyst stage embryos were incubated for 15 min in dissection medium containing 5 µg/ml bisbenzimidazole (Hoechst 33258; Sigma). The embryos were then placed onto clean glass slides in 3 µl drops of diazabicyclo (2.2.2) octane (DABCO; Sigma) and secured with 22 mm² coverslips. Counting took place using an inverted, differential interference contrast microscope fitted with epifluorescence (Nikon, Garden City, NY, USA).

Transmission electron microscopy of oocyte ultrastructure

COCs were collected following 24 h of follicle culture in the presence ($n = 18$) or absence ($n = 18$) of insulin. A proportion of oocytes from each group were matured *in vitro* ($n = 6$ in each treatment). Immature and *in vitro* matured oocytes were processed for electron microscopy using a modification of the protocol described by Hyttel & Madsen (1987). Briefly COCs were fixed in 3% glutaraldehyde in 0.1 M Millonig's buffer (Sigma; pH 7.4), for 1 h at 4°C, then washed three times each for 20 min in 0.2 M buffer and post-fixed in 1% OsO₄ in 0.1 M buffer for 1 h at 4°C. The oocytes were then washed in distilled water for 30 min, placed in 0.5% uranyl acetate (TAAB Laboratories, Reading, UK) for 2 h, and dehydrated in ascending concentrations of ethanol, cleared in inhibisol (1.1.1-trichloroethane), and impregnated and embedded in Araldite. Ultrathin sections (300 Å) of oocytes were cut perpendicular to the plating surface. Oocyte sections were stained for 3 min with uranyl acetate (25% uranyl acetate in 50% ethanol; Watson 1958) and 7 min with lead citrate (Reynolds 1963). Ultra-structural observations were made using a Philips EM300 electron microscope (Eindhoven, The Netherlands).

Total protein profile of oocytes

Oocyte preparation

Non-mature oocytes (total of 54 oocytes in three replicate cultures) were collected following 24 h of follicle culture in the presence ($n = 27$) or absence ($n = 27$) of insulin, transferred into 500 µl drops of dissection medium containing 300 units/ml hyaluronidase (Sigma) and incubated at 39°C for 15 min. After incubation, the oocytes were transferred into dissection medium and the cumulus cells removed by repeated pipetting. Denuded oocytes were placed into 1.5 ml Eppendorf tubes containing 15 µl sample buffer. The oocytes were then lysed by multiple rapid freeze–thaw cycles in liquid nitrogen and methanol respectively. The prepared samples were stored at –20°C until PAGE analysis.

One dimensional SDS-PAGE

A 6% acrylamide gel for protein separation and 4% stacking gel were freshly prepared in a mini-protein II (BioRad, Hemel Hempstead, UK). Prepared samples were thawed on ice, boiled for 3 min to denature the proteins and then stored on ice until loading onto the gel. Molecular weight markers (29 000–116 000) were run on each gel to ascertain the molecular weight in kDa of the separated proteins. A total of five oocytes were loaded/sample well. The gel was run for 6 h at a constant voltage of 200 V. After running, gels were fixed in a solution of 10% glacial acetic acid and 40% methanol in distilled water for at least 1 h.

Silver staining

After fixation, the gels were silver stained using a modification of the method described by Blum *et al.* (1987). Briefly, the gels were washed twice for 20 min in double-distilled water (DDW) followed by a 20-min dehydration in 50% ethanol and pretreated in a 0.2 mg/ml solution of Na₂S₂O₃.5H₂O (BDH, Poole, Dorset, UK) for 1 min. They were then passed through three quick washes in DDW and placed into a solution containing 2 µg/ml silver nitrate (BDH) and 0.75 µl/ml 36% formaldehyde (TAAB Laboratories) for 30 min to allow silver impregnation. After staining, the gels were given two quick washes in DDW and the protein bands were visualised in a development solution containing 60 mg/ml Na₂CO₃ (BDH), 0.5 µl/ml 36% HCHO and 0.5 µg/ml Na₂S₂O₃.5H₂O for 3–4 min. Development was monitored visually and stopped by placing the gel into a solution containing 18.6 mg/ml EDTA (Sigma) and 0.2 mg/ml thiomersol for 30 min. The gels were then washed in DDW for 30 min and equilibrated in a solution containing 33% ethanol and 3% glycerol at least for 1 h.

Analysis of IGFBP profiles

IGFBP profiles in both follicular fluid from individual follicles cultured in the presence (*n* = 14 in 2–4 mm and *n* = 8 in 4–8 mm) and absence of insulin (*n* = 12 in 2–4 mm and *n* = 8 in 4–8 mm) and culture media were analysed by Western ligand blot. Samples of follicular fluid were diluted (2 µl follicular fluid + 23 µl distilled H₂O) and the 25 µl follicular sample or 25 µl culture media were mixed with 25 µl Laemmli sample buffer (10 ml 0.5 M Tris–HCl, pH 6.8, 10 ml 10% SDS and 10 ml glycerol, containing bromophenol blue in 100 ml dH₂O).

Samples were solubilised and loaded onto 12% (w/v) non-reducing SDS-PAGE gels and electroblotted onto PVDF membrane. Non-specific binding sites on the membrane were blocked using 3% (w/v) BSA in PBS–0.1% (v/v) Tween 20 (blocking solution). The blot was then left overnight in 1% BSA and 0.1% Tween 20 in TS buffer (150 mM NaCl, 10 mM Tris–HCl pH 7.4) containing 3 × 10⁶ c.p.m. ¹²⁵I-labelled ligand (in 20 ml). After labelling, the blots were washed four times, each for 30 min, in 0.2% Nonidet P40 in TS buffer. All washing steps were performed at 4 °C. The nitrocellulose was then dried and exposed to a Cronez 4 film (Sterling Diagnostic Imaging

Inc, Newark, DE, USA) for 3 days. The films were developed using an autoradiograph developer.

Statistical analysis

The data from the proportion of oocytes which cleaved after fertilisation and the proportion of cleaved embryos which became blastocysts were analysed by fitting replicates as a source of random variation in a general linear mixed model (Breslow & Clayton 1993).

Results

The developmental competence of oocytes and embryo quality

The effects of addition of insulin to follicle culture media are summarised in Table 1. There was a significant reduction in the proportion of cleaved embryos in the insulin-treated group as compared with the control (means ± S.E.M.; 0.52 ± 0.029 vs 0.61 ± 0.032) (*P* < 0.05). However, there was no significant difference in the proportion of cleaved embryos that became blastocysts (0.30 ± 0.038 in the insulin-positive group vs 0.28 ± 0.047 in the control group; 0.15 ± 0.089 vs 0.18 ± 0.075 of oocytes) nor in the quality of embryos between the two groups as assessed by total cell number (137 ± 8.53 vs 124.6 ± 6.95).

Total protein profile in the oocytes

Total protein profiles of immature oocytes (groups of five oocytes in each category) collected either directly from follicular aspiration (non-mature oocytes aspirated from fresh follicles; NM) or following follicle culture in the presence of insulin (insulin-treated follicles; FNMI) as well as the *in vitro* matured oocytes from these categories (24 h of oocyte maturation (M) and 24 h of follicle culture + 24 h of oocyte maturation (FM)) were compared. A representative profile of total proteins in the oocytes is shown in Fig. 1. Non-matured oocytes recovered from follicle culture in the presence of insulin (immature oocytes from insulin-treated follicles; FNMI) had a remarkably different profile from both immature NM and non-mature oocytes collected after follicle culture (FNM) (insulin-negative) oocytes. This group of oocytes (insulin-positive group) showed a profile more similar to that of matured oocytes. The differences were observed either as reduction in the concentration or disappearance of protein bands such as

Table 1 Effect of addition of insulin during follicle culture on the developmental competence of the recovered oocytes. Insulin adversely reduced the cleavage (*P* < 0.05) rate but there were no differences in either the number or quality of blastocysts between the two groups. The values are presented as means ± S.E.M.

	No. of oocytes	Cleaved (%)*	Blastocysts/oocytes (%)	Blastocysts/cleaved (%)	Cell numbers
Control	98	61 ± 3.28	15 ± 8.9	28 ± 4.74	124.6 ± 6.95
Insulin	106	52 ± 2.92	18 ± 7.5	30 ± 3.81	137.4 ± 8.53

* *P* < 0.05.

Figure 1 Effects of addition of insulin during the culture of bovine large antral follicles on total protein profile of oocytes. Immature oocyte from insulin-treated follicles (FNMI) displayed a protein profile identical to matured oocytes. There was no difference in the protein profile of oocytes after maturation. NM, non-mature oocyte aspirated from fresh follicle; FNM, non-mature oocyte collected after follicle culture; FM, 24 h of follicle culture + 24 h of oocyte maturation; M, 24 h of oocyte maturation; FMI, 24 h of follicle culture in the presence of insulin + 24 h of oocyte maturation.

the 34, 39, 42, 50 and 55 kDa proteins that were also observed following maturation *in vitro*. There were no detectable differences in the total protein profile between the two groups following subsequent *in vitro* maturation (FM vs M). However, there was a significant reduction in the 50 kDa protein in follicles cultured for 24 h in the presence of insulin + 24 h of oocyte maturation (FMI) oocytes.

Ultrastructure of oocytes

A total of 36 oocytes were sectioned and analysed ($n = 24$ immature and 12 matured). Representative images of the oocytes from both treatments are shown in Figs 2–7. The ultrastructure of immature oocytes recovered following follicle culture in the absence of insulin and *in vitro* matured oocyte are shown in Figs 2 and 3 respectively. Immature oocytes were characterised by peripherally located clusters of mitochondria and cortical granules (Fig. 2). In matured oocytes, the mitochondria become evenly distributed throughout the cytoplasm and the cortical granules became individually located in the marginal area under the oocyte membrane (Fig. 3). The ultrastructure of immature oocytes recovered following 24 h of follicle culture in the presence of insulin is shown in Figs 4 and 5. In these oocytes, the cortical granules changed their

position from being in clusters to becoming individually located in the marginal area. The mitochondria became dispersed throughout the cytoplasm. The number of vesicles reduced and these moved towards the central area. The majority of these oocytes exhibited an organelle-free area under the zona pellucida. This feature was recorded in 75% of the oocytes examined ($n = 9$). The majority of the oocyte microvilli remained attached to the zona pellucida. However, some of them became terminated in the previtelline space. Similar features were observed in *in vitro* matured oocytes following follicle culture in the absence (Fig. 6) and presence of insulin (Fig. 7).

IGFBPs

The IGFBPs present in follicular fluid and the culture media were compared following culture of small (2–4 mm)- and medium (4–8 mm)-sized follicles in the presence or absence of insulin. The control contained samples from the pool of follicular fluid collected from the aspiration of oocytes that were used for subsequent maturation. Follicular fluid and culture media collected from the culture of individual follicles cultured in the presence ($n = 22$ follicles) and absence ($n = 20$ follicles) of insulin were analysed by Western blotting ($n = 5$ Western blots repeats). A representative profile of IGFBPs is shown in Fig. 8. Whilst the control group showed moderate amounts of IGFBPs, there were major qualitative differences in both culture medium and follicular fluid between the insulin-positive and control groups. In insulin-treated follicles two strong bands corresponding to IGFBP-4 (19–22 kDa) and -2 (29–34 kDa) disappeared in both the follicular fluid and the culture media of 4–8 mm diameter follicles. In contrast, smaller follicles retained the same pattern of IGFBPs.

Discussion

The results of the present studies have demonstrated that the presence of insulin has a number of effects on the immature or GV stage oocyte, including changes in protein profile and ultrastructure. In addition, following subsequent maturation and fertilisation, a decrease in cleavage rates was observed in insulin-treated oocytes. Transcripts for the receptors of insulin have been detected at all developmental stages of preimplantation bovine embryos (Watson *et al.* 1994). Insulin stimulates both amino acid uptake and protein synthesis (Harvey & Kaye 1988, Kaye *et al.* 1992) in mouse preimplantation embryos. We observed significant quantitative changes in the total protein profile of the immature oocytes following follicle culture in the presence of insulin. For example, proteins with molecular weights of 34, 39, 42 and 50 kDa were decreased or became undetectable. The involvement of proteins of these molecular weights has previously been documented in a number of different species (Cicirelli *et al.* 1988, Mutter *et al.* 1988, Naz *et al.* 2001). The 50 kDa protein that was reduced in FMI oocytes may be the oocyte

Figure 2 Electron micrograph showing part of an immature oocyte following 24 h of follicle culture (FNM) in the control group. Note the clusters of mitochondria (M) and cortical granules (CG) located peripherally and distribution of fat vesicles (V) throughout the cytoplasm. ZP, zona pellucida.

Figure 3 Electron micrograph showing part of an *in vitro* matured control oocyte. Note relocation of the mitochondria (M) throughout the cytoplasm and location of the cortical granules (CG) in the marginal area. The number of fat vesicles (V) has reduced. ZP, zona pellucida; Gr, granulosa cell.

Figure 4 Electron micrograph showing part of an immature oocyte following culture of large antral follicles in the presence of insulin. The oocyte gained the features of a matured oocyte. The mitochondria (M) became evenly distributed throughout the cytoplasm (see inset). Some of the small vesicles fused together and formed giant vesicles (V). Note the organelle-free area in the marginal area. The cortical granules (CG) became located individually in the margin of oocyte. The inset shows high magnification of a group of mitochondria. ZP, zona pellucida.

Figure 5 Electron micrograph showing part of an immature oocyte following culture of large antral follicles in the presence of insulin. Cortical granules (CG) are located individually in the margin of oocyte. Many of the microvilli are disconnected and terminate in the perivitelline space (see inset) but some of them still penetrated into the zona pellucida (ZP). MV, microvilli.

membrane antigen that is important for binding to sperm during fertilisation (Naz *et al.* 2001). Reduction in the expression of this protein may have contributed to the lower cleavage rate of the FMI oocytes.

There is a limited number of studies on the effects of insulin on oocyte maturation and preimplantation

bovine embryo development. Over-stimulation of IGF-I (Armstrong *et al.* 2001) and probably insulin (Armstrong *et al.* 2003) is detrimental to follicle and oocyte development. In mice, high concentrations of IGF-I and insulin induced a down-regulation of the IGF-I receptor in blastocysts, with a subsequent decrease in signalling of IGF-I

Figure 6 Electron micrograph showing part of an *in vitro* matured oocyte following follicle culture in the absence of insulin. Note the relocation of mitochondria (M) throughout the cytoplasm and relocation of the cortical granules (CG) in the marginal area (see inset). The number of fat vesicles (V) has reduced. ZP, zona pellucida; G, Golgi apparatus.

receptor-associated pathways (Chi *et al.* 2000). This decrease in IGF-I receptor reduced glucose uptake and triggered apoptosis. In addition, women with polycystic ovary syndrome exhibit elevated concentrations of insulin and IGF-I and experience increased pregnancy losses (Tulppala *et al.* 1993, Zephne *et al.* 2004, Dumesic *et al.*

2005). A threshold may exist at which the level of growth hormone, IGF-I or insulin goes from being beneficial to detrimental on oocyte and embryo development. Matsui *et al.* (1997) reported that insulin at concentrations of 0.5–10 $\mu\text{g/ml}$ had a beneficial effect on the rate of development to the morula stage on day 5 of embryo culture

Figure 7 Electron micrograph showing part of an *in vitro* matured oocyte following follicle culture in the presence of insulin. Cortical granules (CG) are located individually in the margin of oocyte. Microvilli are disconnected and terminate in the perivitelline space (see inset). V, fat vesicle; M, mitochondria; ZP, zona pellucida.

Figure 8 Effects of addition of insulin during culture of bovine antral follicle on the IGFBP profile. Follicles from two size ranges of small (2–4 mm) and medium (4–8 mm) were cultured in the presence (I^+) or absence of insulin (I^-). Follicular fluid (FF) and culture medium (CM) were collected from two size groups of follicle following 24 h of follicle culture. Controls, follicular fluid from directly aspirated follicles. a vs b, $P < 0.001$; b vs c, $P < 0.01$; a vs c, $P < 0.01$; d vs e, $P < 0.05$; f vs g, $P < 0.01$.

and that this stimulation of embryonic development was mediated through the IGF-I receptor. However, these authors did not report the frequency of development to the blastocyst stage. In the studies reported here, insulin at a concentration of 5 $\mu\text{g/ml}$ was added to the follicle culture medium. This level of insulin, which is routinely used for the culture of preantral follicles, is greater than the physiological concentrations of insulin reported in biological fluids (0.5–10 ng/ml) (Diamond *et al.* 1985). However, although this high concentration of insulin resulted in a decrease in oocyte cleavage, the frequency of blastocyst production was not affected. In addition, the quality of blastocysts produced (as assessed by total cell number) was not significantly different between the two groups. In the present study, we have used Waymouth medium to culture follicles. Waymouth medium is extensively used for culture of follicles in different species. However, it contains a high concentration of glucose that may affect oocyte maturation. In oocytes, glucose is essential for optimal completion of nuclear maturation to metaphase II stage, but

supraphysiological levels may be detrimental and compromise subsequent developmental capacity (Hashimoto *et al.* 2000). Exposure of mouse oocytes to diabetic conditions during folliculogenesis has a negative effect on meiotic maturation and subsequent development, due to decreased communication between the somatic and germ cell compartments (Colton *et al.* 2003). In the mouse, alterations in energy substrate supplementation profoundly influence meiotic regulation *in vitro*. For example, high glucose induces germinal vesicle breakdown in COCs in the presence of meiotic inhibitors (Downs & Mastropolo 1994). In the present study, a lower cleavage rate was observed in oocytes derived from the insulin-treated group. It seems that insulin supplied a surplus amount of glucose for oocytes that resulted in an alteration of oocyte cytoplasmic maturation which adversely affected cleavage rate. However, no differences were observed in the developmental competence of oocytes as assessed by frequency of development to blastocyst stage and total cell numbers. Indeed, although not significant, slightly higher numbers of

blastocysts were produced in oocytes derived from the insulin-treated follicle culture group. In addition, these blastocysts contained slightly increased total cell numbers than the oocytes from the insulin-negative group.

Any effects of the insulin family of growth factors on the oocyte *in vivo* are possibly regulated by IGFBPs (Funston *et al.* 1996). IGFBPs have a regulatory role in follicle development (Echternkamp *et al.* 1994) and are developmentally regulated in bovine granulosa and theca cells (Spicer & Echternkamp 1995). IGFBP-2 and possibly IGFBP-4 and -5 are high in small- and medium-sized follicles, but reduce or become undetectable in large or dominant follicles, resulting in increased IGF bioactivity and enhancing the amplification of gonadotrophin action on follicular cells (Spicer & Echternkamp 1995, Funston *et al.* 1996, de la Sota *et al.* 1996). In contrast, atresia is characterised by an increase in concentrations of the low molecular weight IGFBPs in the follicular fluid (Besnard *et al.* 1997).

The present study on the effects of insulin on IGFBPs in both culture media and follicular fluid indicate that in insulin-treated follicles of >4 mm in diameter, two protein bands with estimated molecular weights of 19–22 and 29–34 kDa corresponding to the low molecular weight IGFBP-4 and 2 were significantly reduced to undetectable levels. It is possible that this reduction of low molecular weight IGFBPs resulted in protection of larger follicles against atresia. However, in small follicles, insulin did not protect follicles against atresia, so the concentrations of IGFBPs increased. During follicular development in cattle, the IGFBP-3 content of follicular fluid increases (Armstrong *et al.* 1996). In addition, growth of follicles larger than 4 mm in diameter is FSH dependent (Gong *et al.* 1995). The presence of insulin may have a compensatory role in this group of follicles, resulting in healthier follicles. On the other hand, the disappearance of low molecular weight IGFBPs in larger follicles may be explained by intrinsic changes in the follicle and any effect of insulin on the follicle are ignored.

Analysis of the ultrastructure of the recovered oocytes showed that insulin-treated follicles yield oocytes which have features normally associated with matured oocytes. This may explain the reduced cleavage rate after fertilisation. As these oocytes were cultured in maturation medium for a further 24 h, to allow nuclear maturation, the early occurrence of cytoplasmic changes may suggest that the oocytes became hypermature or aged, thus resulting in a lower fertilisation rate than the control group. In porcine (May & Schomberg 1981), human (Hill & Osteen 1992) and bovine (Spicer *et al.* 1993) ovarian tissue, insulin stimulates granulosa cell progesterone secretion, granulosa cell luteinisation (Channing *et al.* 1967) and induces granulosa cell LH/hCG receptors (Osteen *et al.* 1985, May & Schomberg 1981). In the experiments presented here, the levels of progesterone were not compared between the two groups. If insulin supplementation increases progesterone level in the follicular fluid, then it may mimic

the phenomenon at the preovulatory surge of LH that leads to resumption of meiosis in the oocytes.

In conclusion, follicle culture in the presence of insulin has no detrimental effect on oocyte structure and does not improve bovine oocyte quality or preimplantation embryo development *in vitro*, but appears to promote cytoplasmic maturation. In addition, the presence of insulin during follicle culture may prevent the process of atresia in medium-sized antral follicles.

Acknowledgements

The authors would like to thank Dr David Armstrong and Mrs C Hogg for IGFBP analysis. The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

References

- Armstrong DG, Hogg CO, Campbell BK & Webb R 1996 Insulin-like growth factor (IGF)-binding protein production by primary cultures of ovine granulosa and theca cells. The effects of IGF-I, gonadotropin, and follicle size. *Biology of Reproduction* **55** 1163–1171.
- Armstrong DG, McEvoy TG, Baxter G, Robinson JJ, Hogg CO, Woad KJ, Webb R & Sinclair KD 2001 Effect of dietary energy and protein on bovine follicular dynamics and embryo production *in vitro*: associations with the ovarian insulin-like growth factor system. *Biology of Reproduction* **64** 1624–1632.
- Armstrong DG, Gong J & Webb R 2003 Interactions between nutrition and ovarian activity in cattle: physiological, cellular and molecular mechanisms. *Reproduction* **61** (Supplement) 403–414.
- Besnard N, Pisselet C, Monniaux D & Monget P 1997 Proteolytic activity degrading insulin-like growth factor-binding protein-2, -3, -4, and -5 in healthy growing and atretic follicles in the pig ovary. *Biology of Reproduction* **56** 1050–1058.
- Betteridge KJ, Smith C, Stubbings RB, Xu KP & King WA 1989 Potential genetic improvement of cattle by fertilization of fetal oocytes *in vitro*. *Journal of Reproduction and Fertility* **38** (Supplement) 87–98.
- Blum F, Beier H & Gross HJ 1987 Improved silver staining of plant proteins, RNA and DNA in polyacrylamide gels. *Electrophoresis* **8** 93–99.
- Breslow NE & Clayton DG 1993 Approximate inference in generalised linear mixed models. *Journal of the American Statistical Association* **88** 9–25.
- Campbell BK, Scaramuzzi RJ & Webb R 1995 Control of antral follicle development and selection in sheep and cattle. *Journal of Reproduction and Fertility* **49** (Supplement) 335–350.
- Chang AS, Dale AN & Moley KH 2005 Maternal diabetes adversely affects preovulatory oocyte maturation, development, and granulosa cell apoptosis. *Endocrinology* **146** 2445–2453.
- Channing CP, Tsai V & Sachs D 1976 Role of insulin, thyroxine and cortisol in luteinisation of porcine granulosa cells grown in chemically defined media. *Biology of Reproduction* **15** 235–247.
- Chi MM, Schlein AL & Moley KH 2000 High insulin-like growth factor 1 (IGF-1) and insulin concentrations trigger apoptosis in the mouse blastocyst via down-regulation of the IGF-1 receptor. *Endocrinology* **141** 4784–4792.
- Cicirelli MF, Pelech SL & Krebs EG 1988 Activation of multiple protein kinases during the burst in protein phosphorylation that precedes the first meiotic cell division in *Xenopus* oocytes. *Journal of Biological Chemistry* **263** 2009–2019.
- Colton SA, Humpherson PG, Leese HJ & Downs SM 2003 Physiological changes in oocyte–cumulus cell complexes from diabetic

- mice that potentially influence meiotic regulation. *Biology of Reproduction* **69** 761–770.
- Davoren JB & Hsueh AJW** 1984 Insulin enhances FSH-stimulated steroidogenesis by cultured rat granulosa cells. *Molecular and Cellular Endocrinology* **35** 97–105.
- Diamond MP, Webster BW, Carr RK, Wentz AC & Osteen KG** 1985 Human follicular fluid insulin concentrations. *Journal of Clinical Endocrinology and Metabolism* **61** 990–992.
- Downs SM & Mastropolo AM** 1994 The participation of energy substrates in the control of meiotic maturation in murine oocytes. *Developmental Biology* **162** 154–168.
- Duleba AJ, Spaczynski RZ, Olive DL & Behrman HR** 1997 Effects of insulin and insulin-like growth factors on proliferation of rat ovarian theca-interstitial cells. *Biology of Reproduction* **56** 891–897.
- Dumesic DA, Schramm RD & Abbott DH** 2005 Early origins of polycystic ovary syndrome. *Reproduction, Fertility and Development* **17** 349–360.
- Echternkamp SE, Howard HJ, Roberts AJ, Grizzle J & Wise T** 1994 Relationships among concentrations of steroids, insulin-like growth factor-I, and insulin-like growth factor binding proteins in ovarian follicular fluid of beef cattle. *Biology of Reproduction* **51** 971–981.
- Fouladi-Nashta AA, Waddington D & Campbell KHS** 1998 Maintenance of bovine oocytes in meiotic arrest and subsequent development *in vitro*. A comparative evaluation of antral follicle culture with other methods. *Biology of Reproduction* **59** 255–262.
- Fulghesu AM, Villa P, Pavona V, Guido M, Apa R, Carusa A, Lanzone A, Rossodivita A & Mancuso S** 1997 The impact of insulin secretion on the ovarian response to exogenous gonadotropins in polycystic ovary syndrome. *Journal of Clinical Endocrinology and Metabolism* **82** 644–648.
- Funston RN, Seidel GE Jr, Klindt J & Roberts AJ** 1996 Insulin-like growth factor I and insulin-like growth factor-binding proteins in bovine serum and follicular fluid before and after the preovulatory surge of luteinizing hormone. *Biology of Reproduction* **55** 1390–1396.
- Gong JG, Bramley TA & Webb R** 1993a The effect of recombinant bovine somatotrophin on ovarian follicular growth and development in heifers. *Journal of Reproduction and Fertility* **97** 247–254.
- Gong JG, Bramley TA, Wilmut I & Webb R** 1993b Effect of recombinant bovine somatotrophin on superovulatory response to pregnant mare serum gonadotrophin in heifers. *Biology of Reproduction* **48** 1141–1149.
- Gong JG, Bramley TA, Gutierrez CG, Peters AR & Webb R** 1995 Effects of chronic treatment with a gonadotrophin-releasing hormone agonist on peripheral concentrations of FSH and LH, and ovarian function in heifers. *Journal of Reproduction and Fertility* **105** 263–270.
- Harvey MB & Kaye PL** 1988 Insulin stimulates protein synthesis in compacted mouse embryos. *Endocrinology* **122** 1182–1184.
- Hashimoto S, Minami N, Yamada M & Imai H** 2000 Excessive concentration of glucose during *in vitro* maturation impairs the developmental competence of bovine oocytes after *in vitro* fertilization: relevance to intracellular reactive oxygen species and glutathione contents. *Molecular Reproduction and Development* **56** 520–526.
- Hill GA & Osteen KG** 1992 The effect of insulin on *in vitro* progesterone production of human granulosa cells. *International Journal of Fertility* **37** 93–98.
- Hsueh AJW, Adashi EY, Jones PBC & Welsh TH Jr** 1984 Hormonal regulators of the differentiation of cultured ovarian granulosa cells. *Endocrine Reviews* **5** 76–127.
- Hyttel P & Madsen I** 1987 Rapid method to prepare mammalian oocytes and embryos for transmission electron microscopy. *Acta Anatomica* **129** 12–14.
- Jones JI & Clemmons DR** 1995 Insulin-like growth factors and their binding proteins: biological actions. *Endocrine Reviews* **16** 3–34.
- Kaye PL, Bell KL, Beebe LFS, Dunglison GF, Gardner HG & Harvey MB** 1992 Insulin and the insulin-like growth factors (IGFs) in preimplantation development. *Reproduction, Fertility and Development* **4** 373–386.
- Kruip TA & Dieleman SJ** 1982 Macroscopic classification of bovine follicles and its validation by micromorphological and steroid biochemical procedures. *Reproduction, Nutrition, Development* **22** 465–473.
- Lessman CA & Schuetz AW** 1981 Role of follicle wall in meiosis reinitiation induced by insulin in *Rana pipiens* oocytes. *American Journal of Physiology* **241** E51–E56.
- Mason HD, Willis D, Holly JM, Cwyfan-Hughes SC, Seppala M & Franks S** 1992 Inhibitory effects of insulin-like growth factor-binding proteins on steroidogenesis by human granulosa cells in culture. *Molecular and Cellular Endocrinology* **89** R1–R4.
- Matsui M, Takahashi Y, Hishinuma A & Kanagawa H** 1997 Stimulation of the development of bovine embryos by insulin and insulin-like growth factor-I (IGF-I) is mediated through the IGF-I receptor. *Theriogenology* **48** 605–616.
- May JV & Schomberg DW** 1981 Granulosa cell differentiation *in vitro*: effect of insulin on growth and functional integrity. *Biology of Reproduction* **25** 421–431.
- May JV, McCarty K Jr, Reichert LE Jr & Schomberg DW** 1980 Follicle stimulating hormone mediated induction of functional luteinising hormone/human chorionic gonadotrophin receptors during monolayer culture of porcine granulosa cells. *Endocrinology* **107** 1041–1049.
- Monget P, Monniaux D, Pisselet C & Durand P** 1993 Changes in insulin-like growth factor-I (IGF-I), IGF-II, and their binding proteins during growth and atresia of ovine ovarian follicles. *Endocrinology* **132** 1438–1446.
- Moor RM, Hay ME, McIntosh JEA & Caldwell BV** 1973 Effect of gonadotrophins on the production of steroids by sheep ovarian follicles cultured *in vitro*. *Journal of Endocrinology* **58** 599–611.
- Mutter GL, Grills GS & Wolgemuth DJ** 1988 Evidence for the involvement of the proto-oncogene c-mos in mammalian meiotic maturation and possibly very early embryogenesis. *EMBO Journal* **7** 683–689.
- Naz RK, Leslie MH & Zhu X** 2001 Inhibition of murine sperm-oolemma binding by antibodies to an oocyte membrane (OM) antigen: implication in contraceptive vaccine development. *American Journal of Reproductive Immunology* **45** 52–61.
- Osteen KG, Anderson LD, Reichert LE & Channing CP** 1985 Follicular fluid modulation of functional LH receptor induction in pig granulosa cells. *Journal of Reproduction and Fertility* **74** 407–418.
- Poretsky L** 1991 On the paradox of insulin-induced hyperandrogenism in insulin-resistant states. *Endocrine Reviews* **12** 3–13.
- Poretsky L & Kalin MF** 1987 The gonadotrophic function of insulin. *Endocrine Reviews* **8** 132–141.
- Rao LV, Wikarczuk ML & Heyner S** 1990 Functional roles of insulin and insulin like growth factors in preimplantation mouse embryo development. *In Vitro Cellular and Developmental Biology* **26** 1043–1048.
- Reynolds ES** 1963 The use of lead citrate at high pH as an electron-opaque stain in electron microscopy. *Journal of Cell Biology* **17** 208–212.
- Schultz GA, Hogan A, Watson AJ, Smith RM & Heyner S** 1992 Insulin, insulin-like growth factors and glucose transporters: temporal patterns of gene expression in early murine and bovine embryos. *Reproduction, Fertility and Development* **4** 361–371.
- de la Sota RL, Simmen FA, Diaz T & Thatcher WW** 1996 Insulin-like growth factor system in bovine first-wave dominant and subordinate follicles. *Biology of Reproduction* **55** 803–812.
- Spicer LJ, Alpizar A & Echternkamp SE** 1993 Effects of insulin, insulin-like growth factor I, and gonadotrophins on bovine granulosa cell proliferation, progesterone production, oestradiol production, and (or) insulin-like growth factor-I production *in vitro*. *Journal of Animal Science* **71** 1232–1241.

- Spicer LJ & Echternkamp SE** 1995 The ovarian insulin and insulin-like growth factor system with an emphasis on domestic animals. *Domestic Animal Endocrinology* **12** 223–245.
- Straus DS** 1981 Effects of insulin on cellular growth and proliferation. *Life Science* **29** 2131–2139.
- Stubbings RB, Liptrap RM, Betteridge KJ, Walton JS, Armstrong DJ & Basrur PK** 1990 Requirements for bovine oocyte maturation *in vitro*. *Reproduction in Domestic Animals* **25** 158–166.
- Thompson JGD, Simpson AC, Paugh PA, Wright RW & Tervit HR** 1991 Glucose utilisation by sheep embryos derived *in vivo* and *in vitro*. *Reproduction, Fertility and Development* **3** 571–576.
- Tsafriri A & Channing CP** 1975 Influence of follicular maturation and culture conditions on the meiosis of pig oocytes *in vitro*. *Journal of Reproduction and Fertility* **43** 149–152.
- Tulppala M, Stenman UH, Cacciatori B & Ylikorkala O** 1993 Polycystic ovaries and levels of gonadotrophins and androgens in recurrent miscarriage: prospective study in 50 women. *British Journal of Obstetrics and Gynaecology* **1993** **100** 348–352.
- Watson AJ, Watson PH, Arcellana-Panlilio M, Warnes D, Ealker SK, Schultz GA, Armstrong DT & Seamark RF** 1994 A growth factor phenotype map for ovine preimplantation development. *Biology of Reproduction* **50** 725–733.
- Watson ML** 1958 Staining of tissue sections for electron microscopy with heavy metals. *Journal of Biophysics and Biochemical Cytology* **4** 475–478.
- Zephne M, vanderSpuy & Dyer SJ** 2004 The pathogenesis of infertility and early pregnancy loss in polycystic ovary syndrome. *Best Practice and Research in Clinical Obstetrics and Gynaecology* **18** 755–771.

Received 24 November 2004

First decision 3 February 2005

Revised manuscript received 19 August 2005

Accepted 22 November 2005