Peptidase activities in *Crotalus durissus terrificus* semen

Camila Eduardo Marinho^{1,4}, Selma Maria Almeida-Santos², Sylvia Mendes Carneiro³, Simone Cristina Yamasaki¹ and Paulo Flavio Silveira¹

¹Laboratory of Pharmacology, ²Laboratory of Ecology and Evolution and ³Laboratory of Cellular Biology, Instituto Butantan, Av. Vital Brasil, 1500, São Paulo SP 05503-900, Brazil and ⁴Department of Physiology, Instituto de Biociências, Universidade de São Paulo, São Paulo SP 05508-900, Brazil

Correspondence should be addressed to P F Silveira; Email: pefesil@butantan.gov.br

Abstract

To understand the role of peptidases in seminal physiology of *Crotalus durissus terrificus*, activity levels of representative enzymes in semen and their sensitivities to inhibitors, cofactors, and peptide hormones were evaluated. The existence of seminal fractions and the association of peptidases with these fractions were also characterized for the first time in snakes. The prominent inhibitors of aminopeptidases (APs) were amastatin for acid, basic, and neutral; bestatin for basic; and diprotin A for dipeptidyl-IV. Cystyl and prolyl-imino APs were similarly susceptible to the majority of these inhibitors. The basic and neutral were characterized as metallo-peptidases, acid AP was activated by MnCl₂, and cystyl, prolyl-imino, and type I pyroglutamyl were characterized as sulphhydryl-dependent APs. Angiotensin II, vasotocin, bradykinin, fertilization-promoting peptide, and TRH altered the majority of these peptidase activities; these peptides are possible substrates and/or modulators of these peptidases. Peptidase activities were found in all seminal fractions: seminal plasma (SP), prostatesome-like (PR) structures, and soluble (S-) and membrane-bound fractions (MFs) of spermatozoa. The levels of activity of each peptidase varied among different seminal fractions. In SP, the higher activities were puromycin-insensitive neutral and basic APs. In PR, the higher activity was puromycin-insensitive neutral AP. In spermatozoa, the higher activity in subcellular SF was puromycin-sensitive neutral, while in MF both puromycin-sensitive and -insensitive neutral APs were equally higher than the other examined peptidases. Data suggested that these peptidases, mainly basic and neutral, have a high relevance in regulating seminal functions of *C. d. terrificus*.

Reproduction (2008) 136 767–776

Introduction

Enzyme hydrolysis leading to inactivation or processing of peptides has been assumed to be a limitation step for their biological activities. Many peptides, such as angiotensins (Angs; Fraser *et al.* 2006), kinins (Blaukat 2003), substance P, enkephalins (Sastry *et al.* 1991), endorphin (El-Haggar *et al.* 2006), oxytocin (OXT), vasopressin (AVP; Assinder *et al.* 2000), LHRH (Amory & Bremner 2003), TRH, and fertilization-promoting peptide (FPP; Green *et al.* 1996), as well some peptidases, particularly aminopeptidases (APs) and oligopeptidases have been related to seminal physiology in mammals (Fernández *et al.* 2002, Irazusta *et al.* 2004, Valdivia *et al.* 2004, Subirán *et al.* 2008). In this sense, Ang I and II are hydrolyzed by acid AP (APA; EC 3.4.11.7; Kugler 1982) and, also, by prolyl oligopeptidase (POP; EC 3.4.21.26; Barret *et al.* 1998). The formation of Ang IV from Ang III (Kugler 1982) and bradykinin (BK) from kallidin (Mizutani *et al.* 1993) is catalyzed by neutral AP (APN). Enkephalin is hydrolyzed by puromycin-insensitive APN (APN-PI; E.C 3.4.11.2) and puromycin-sensitive APN (APN-PS, E.C 3.4.11.14; Fernández *et al.* 2002). Substance P,

enkephalin, BK, OXT, AVP, LHRH (Barret *et al.* 1998), TRH, and FPP (Siviter & Cockle 1995) are hydrolyzed by POP. OXT and AVP are hydrolyzed by cystyl AP (CAP, EC 3.4.11.3; Davison *et al.* 1993). LHRH and TRH are susceptible to type I pyroglutamyl AP (PAP-I, EC 3.4.19.3; Cummins & O'Connor 1998), and FPP is a possible substrate of this enzyme. BK, kallidin, met-enkephalin, and somatostatin are hydrolyzed by basic AP (APB, EC 3.4.11.6; Barret *et al.* 1998). Substance P and endorphin-2 are substrates of dipeptidyl peptidase-IV (DPPIV, EC 3.4.14.5; Barret *et al.* 1998).

The distribution of peptidases in seminal fractions is fundamental to the regulatory role of these enzymes. Until the present, earlier studies of human seminal fractions evaluated the distribution of only three of the above-mentioned peptidases: APN (Fernández *et al.* 2002, Subirán *et al.* 2008), PAP, and POP (Valdivia *et al.* 2004). APN was detected in soluble (S-) and solubilized membrane-bound fractions (MFs) of seminal plasma (SP) and prostatesomes (PRs; Fernández *et al.* 2002). In sperm, APN was detected in the equatorial segment of the upper post-acrosomal region of the head, in the neck, and

along the tail (Subirán *et al.* 2008). Alterations of APN levels in human semen were associated with different abnormalities in the sperm of subfertile patients (Irazusta *et al.* 2004). PAP-I and POP activities were detected in SP and in the PR fraction, as well as in soluble and membrane-bound sperm subcellular fractions, and have higher activities in necrozoospermia than in normozoospermic semen (Valdivia *et al.* 2004). AP and amylase activities are also known to be the main agents in the liquefaction process of the ejaculated semen coagulum, in humans (Chatterjee *et al.* 1997). Moreover, AP inhibitors, bestatin and puromycin, are capable of diminishing cellular proliferation and viability in mammals (Takahashi *et al.* 1989, Constam *et al.* 1995). However, bestatin is capable of restoring and promoting follicular growth and ovulation after its suppression by stress (Nakamura *et al.* 1998).

To our knowledge, the fractionation of semen and the presence of seminal APs and POP have not yet been studied in snakes. Furthermore, species possessing sperm with long lifespans, such as the rattlesnake *Crotalus durissus terrificus*, are interesting experimental models to study the distribution and function of peptidases in seminal fractions. In *C. d. terrificus*, spermatogenesis begins in spring and has its peak in summer. The spermatozoa are maintained in the male tract until mating (middle of autumn; Almeida-Santos *et al.* 2004) and, subsequently, in the female tract until ovulation (spring; Almeida-Santos & Salomão 1997). The goal of the present study was to evaluate the existence of mammalian-like seminal fractions (SP, PR, and sperm), and to characterize the activity levels of soluble and membrane-bound APA, APB, APN-PI, APN-PS, CAP, PAP-I, DPPIV, POP, and prolyl iminopeptidase (PIP) and their sensitivities to inhibitors, cofactors, and possible natural substrates in whole semen, as well as their associations with different seminal fractions of *C. d. terrificus*.

Results

Figure 1 shows that lactate dehydrogenase (LDH) activity in SF was threefold higher than in MF of whole semen. Similar proportion was obtained in spermatozoa subcellular SF and MF (data not shown).

Table 1 shows values of peptidase activity in SF and MF of whole semen. APB, PIP, and POP activities were detected only in SF, while the others were detected in SF and MF.

Table 2 shows that APB activity was markedly inhibited by amastatin and bestatin, APA and APN by amastatin, and diprotin A had a stronger effect on DPPIV activity in SF and MF. CAP and PIP activity decreased in the presence of majority of examined inhibitors. Bestatin and puromycin were more efficient to inhibit PAP. POP activity was not evaluated by classical AP inhibitors, since it is an endooligopeptidase (the only one under study here), and this activity was shown to be increased

Figure 1 LDH activity (means ± S.E.M.) in soluble (white bar) and solubilized membrane-bound (black bar) fractions of whole semen of *C. d. terrificus*. Number of animals is given in parentheses. * $P < 0.0003$ (unpaired two-side Student's *t*-test).

or indifferent with the adopted agents (Olivo *et al.* 2008). CAP activity in SF, PAP-I, and PIP had a common pattern of decrease with BK, TRH, and FPP (with or without Ca^{2+}). Ang II and BK diminished APA activity in SF and MF. Ang II and arginine vasotocin (AVT) decreased CAP activity in SF and PIP activity. AVT also decreased CAP activity in MF. DL-dithiothreitol (DTT) improved CAP activity in SF and MF and PIP activity, and decreased APB activity. EDTA diminished the activities of APB and APN SF and MF. MnCl_2 decreased the activities of CAP in MF, PAP, and DPPIV SF and MF, and increased APA activity.

As shown in Fig. 2, a peak coincided with the dead volume (V_0) of seminal fraction from gel filtration in Sephadex G-200. This peak was analyzed by transmission electron microscopy and had numerous rounded or oval membranous vesicles of the size 50–200 nm, shown in Fig. 3.

Figure 4 shows AP activities in seminal fractions: SP, PR, and spermatozoa subcellular SF and MF. In SP, the higher activities were APN-PI and APB relative to other examined peptidases. In PR, APN-PI activity also had the higher level, followed by APB and APN-PS. In spermatozoa subcellular SF, APN-PS activity was higher, while in spermatozoa subcellular MF, APN-PS and APN-PI

Table 1 Activity levels of aminopeptidases: acid (APA), basic (APB), puromycin-insensitive neutral (APN-PI), puromycin-sensitive neutral (APN-PS), dipeptidyl peptidase-IV (DPPIV), pyroglutamy-I (PAP-I), prolyl-imino (PIP); and prolyl oligopeptidase (POP) in soluble (SF) and solubilized membrane-bound (MF) fractions of whole semen of *Crotalus durissus terrificus*.

Aminopeptidase	Activities (U/mg protein)	
	SF	MF
APA	691 ± 180 ^a (16)	838 ± 229 ^a (16)
APB	18 549 ± 3227 ^{b,*} (15)	Absent
APN-PI	30 372 ± 3107 ^c (13)	23 866 ± 1938 ^b (13)
APN-PS	9019 ± 1133 ^{d,*} (10)	12 637 ± 1290 ^c (11)
CAP	31 ± 14 ^a (14)	30 ± 13 ^a (15)
DPPIV	234 ± 57 ^{a,*} (13)	560 ± 153 ^a (11)
PAP-I	261 ± 34 ^a (14)	231 ± 71 ^a (16)
PIP	257 ± 66 ^{a,*} (15)	Absent
POP	1777 ± 252 ^{a,d,*} (14)	Absent

Values are means ± S.E.M. Number of animals is given in parentheses. U = picomoles of hydrolyzed substrate per minute. Different letters indicate different peptidase activity levels in the same fraction (One-way ANOVA $P < 0.0001$, Bonferroni $P < 0.05$). * $P < 0.05$ in comparison to MF (unpaired, two-side Student's *t*-test).

Table 2 Remaining activities of aminopeptidases: acid (APA), basic (APB), puromycin-insensitive neutral (APN-PI), puromycin-sensitive neutral (APN-PS), cystyl (CAP), dipeptidyl peptidase-IV (DPP-IV), pyroglutamyl-I (PAP-I), prolyl-imino (PIP); and prolyl oligopeptidase (POP) in soluble (SF) and solubilized membrane-bound (MF) fractions of whole semen of *Crotalus durissus terrificus* with classical aminopeptidase inhibitors and angiotensin II (AngII), arginine vasotocin (AVT), bradykinin (BK), fertilization-promoting peptide (FPP), fertilization-promoting peptide (FPP) plus calcium (FPP+Ca[2+]), TRH, DL-dithiothreitol (DTT), ethylenediaminetetraacetic acid (EDTA) and MnCl₂.

Agents	Remaining peptidase activity (% control)											
	APA			APN			CAP			DPP-IV		
	SF	MF	APB	SF	MF	SF	SF	MF	MF	SF	MF	POP
Amastatin	11 ± 1*	21 ± 3*	5 ± 2*	9 ± 4*	14 ± 2*	83 ± 18	0 ± 7*	51 ± 7*	89 ± 14	78 ± 19	92 ± 27	78 ± 22
Bestatin	76 ± 7*	33 ± 7*	7 ± 2*	69 ± 9*	21 ± 5*	8 ± 2*	0 ± 5*	52 ± 9*	91 ± 12	56 ± 9*	45 ± 12*	16 ± 9*
Diprotin A	99 ± 8	84 ± 11	112 ± 23	100 ± 12	54 ± 8*	48 ± 5*	0 ± 4*	51 ± 9*	68 ± 11*	97 ± 21	92 ± 24	67 ± 5*
Puromycin	97 ± 13	75 ± 15	95 ± 11	93 ± 11	45 ± 9*	34 ± 7*	11 ± 4*	69 ± 11*	91 ± 11	56 ± 13*	61 ± 9*	44 ± 7*
AngII	40 ± 10*	35 ± 2*	105 ± 18	104 ± 8	84 ± 12	20 ± 9*	0 ± 3*	89 ± 11	85 ± 19	94 ± 11	82 ± 16	29 ± 7*
AVT	99 ± 2	78 ± 11	112 ± 17	105 ± 8	80 ± 11	28 ± 9*	0 ± 6*	97 ± 13	94 ± 21	199 ± 21	89 ± 32	118 ± 22
BK	77 ± 11*	65 ± 9*	95 ± 9	102 ± 7	84 ± 11	0 ± 6*	0 ± 5*	91 ± 12	92 ± 14	59 ± 16*	34 ± 18*	140 ± 20
FPP	81 ± 9	98 ± 16	99 ± 8	105 ± 18	88 ± 10	36 ± 11*	0 ± 6*	85 ± 15	91 ± 14	0 ± 12*	0 ± 9*	46 ± 6*
FPP+Ca[2+]	96 ± 11	84 ± 19	102 ± 9	104 ± 12	87 ± 21	10 ± 3*	0 ± 9*	72 ± 8*	82 ± 18	17 ± 8*	37 ± 11*	151 ± 21
TRH	92 ± 11	95 ± 7	101 ± 15	102 ± 11	84 ± 11	0 ± 3*	0 ± 6*	86 ± 9	89 ± 17	63 ± 13*	41 ± 19*	4 ± 1*
DTT	86 ± 9	71 ± 19	42 ± 10*	106 ± 11	88 ± 14	2510 ± 193*	7587 ± 654*	88 ± 8	83 ± 18	114 ± 18	129 ± 24	1302 ± 211*
EDTA	88 ± 9	92 ± 6	39 ± 9*	74 ± 9*	16 ± 3*	122 ± 24	236 ± 56	95 ± 10	86 ± 9	81 ± 23	90 ± 21	110 ± 14
MnCl ₂	356 ± 42*	452 ± 51*	86 ± 14	107 ± 8	70 ± 17	84 ± 18	0 ± 2*	67 ± 8*	73 ± 4*	0 ± 14*	0 ± 16*	74 ± 28

Values are means ± S.E.M. from triplicates of pool of 11 animals. Control (absence of any factor) = 100%. *P < 0.05 in comparison with the respective control (unpaired, two-side Student's t-test).

activities were higher than other examined peptidase activities. Considering activity levels of each peptidase among different seminal fractions, APB, PIP, and POP were higher in SP; APB, APN-PI, DPP-IV, and PAP-I in PR; APN-PS in spermatozoa subcellular SF; and PAP-I and CAP in spermatozoa subcellular MF. APA was equally distributed among all seminal fractions. It was noteworthy that APN-PS activity was absent in SP.

Discussion

The efficiency of the employed procedure to separate SF and MF was confirmed by the pronounced LDH activity (cytosolic marker) in SF of all samples, which allowed the characterization of the activity of each peptidase in different fractions, and consequently the association of intra- and extracellular peptides with the peptidase activities under study. Overall, peptides involved in intracellular signaling are primarily hydrolyzed outside the cell, mainly by membrane-bound peptidases which have, in general, the active sites in the extracellular side, which interact with released peptides (O'Cuinn 1998). Soluble peptidases act mainly in intracellular processes, but can also be exocytosed or act in recaptured peptides, which are internalized as a part of the peptide-receptor complex, as occurs with peptide hormones (Gibson *et al.* 1989). Under stimulation, cytosolic enzymes can be translocated to the cell surface and efficiently process extracellular substances (Albiston *et al.* 2004). In the present study, APB, PIP, and POP activities were detected only in SF and the other activities were detected in SF and MF, following the same pattern found in tissues and semen of mammals (Irazusta *et al.* 2001, 2004, Silveira *et al.* 2001, Varona *et al.* 2003).

The use of substances derived from naphthylamide has been the initial step to detect or confirm the involvement of APs and POP in physiological mechanisms. Variables that could not be controlled might have influenced peptidase activities in the present study, e.g. intra-season and geographical and circadian variations. This is possible considering the seasonality, since samples were obtained in different days of the same season (austral autumn – the mating season for *C. d. terrificus*). However, the fertility potential of semen can be presumed effective and homogeneous along this season, based on the existence of a high quality of seminal dynamic parameters of *Boa constrictor occidentalis* (Tourmente *et al.* 2007), the peak of testosterone (Zacariotti *et al.* 2005) and the decrease of spermatozoa defects (Zacariotti 2004) in *C. d. terrificus*. Geographical variation was not a factor as snakes were captured from locations with similar fauna and environmental conditions. Moreover, we can also exclude circadian variations, because the material was obtained in the same period of the day, with a maximum difference of 90 min.

Figure 2 Gel filtration chromatography on Sephadex G-200. The column was equilibrated and washed with Tris-saline buffer (30 mM Tris plus 130 mM NaCl), pH 7.4, flow rate 0.16 ml/min.

The anatomy and morphology of the male reproductive tract could also influence the features of the peptidase activities and their distribution in seminal fractions of *C. d. terrificus*. In reptiles, the ductus deferens are recognized as the site of spermatozoa storage in males and, differently from mammals, the epididymis does not participate in sperm maturation and storage (Sever *et al.* 2002). The ductus deferens of *C. d. terrificus* are convoluted and enter the cloaca independently of the ureters. As previously described by Almeida-Santos *et al.* (2004), the ductus deferens occupy about one-third of the body length and when freed from

Figure 3 Electron microphotography of prostasome-like structures from semen collected from the ductus deferens of *C. d. terrificus* in austral autumn.

the peritoneal membrane and manually distended, its length is about six times the total body length, although the right duct is always slightly longer than the left. Three regions can be distinguished: proximal (testicular), median, and distal (cloacal). In this snake, spermatozoa become increasingly more motile from the proximal to the distal region (Almeida-Santos *et al.* 2004). Regarding the presence of examined peptidase activities in the spermatozoa, the identification of their anatomical sources need to be investigated; for example, in the acrosomal complex, which is known in all vertebrate taxa as an enzyme-rich organelle that facilitates spermatozoal penetration into the ovum (Subirán *et al.* 2008). In squamates, the acrosomal complex is highly compartmentalized (Gribbins *et al.* 2007), and this compartmentalization has been suggested as an aid to the sequential release of acrosomal enzymes (Talbot 1991). Since the scope of the present study was not to compare the seminal fractions and peptidase composition from different areas of the ductus deferens, semen from the entire ductus deferens of *C. d. terrificus* was taken systematically to assure similar seminal fluid components for each analysis.

The APs under study are known to be sensitive to amastatin, bestatin, diprotin A, and puromycin (Cadel *et al.* 1995, Ronai *et al.* 1999, Sato 2003). POP is inhibited by Z-Pro-prolinal (García-Horsman *et al.* 2007) and belonging to a distinct class of enzymes, it was not included in the inhibition assay.

In the present study, APB susceptibility to bestatin distinguished this enzyme from APA and APN. By the susceptibility of DPPIV to diprotin A, and PAP-I, CAP, and PIP to puromycin, it was possible to differentiate them from APA and APB. The inhibition of DPPIV by diprotin A is a known feature of this enzyme (CD26) in mammals (Minelli *et al.* 1999). As distinguishing features, APB and APN are metallo-peptidases, because of their inhibition by EDTA (Kawata *et al.* 1980). APA was marked as activated by $MnCl_2$, which is a known characteristic of this enzyme in mammals (Ramírez *et al.* 1990). CAP and PAP-I were activated by DTT, a thiol-reducing agent, characterizing them as sulphydryl-dependent peptidases. DTT is an activator of the type I PAP and an inhibitor of the type II (O'Cuinn *et al.* 1990), and the degradation of FPP (possible substrate of PAP-I) is also increased by the presence of DTT (Cockle *et al.* 1994), which confirms that PAP-I is among the enzymes studied in the present work and FPP is one of its substrates. Overall, the present results show that seminal peptidase activities of *C. d. terrificus* have distinct biochemical properties, and therefore it is possible to attribute them to different proteins. It was notable that enzyme activities were affected by the evaluated peptides, which are their possible substrates (e.g. Ang II on APA, AVT on CAP, and PIP, and particularly TRH and FPP with or without Ca^{2+} on CAP, PIP, and PAP-I). These results indicate the competition among these

Figure 4 Distribution of activity levels of aminopeptidases: acid (APA), basic (APB), puromycin-insensitive neutral (APN-PI), puromycin-sensitive neutral (APN-PS), cystyl (CAP), dipeptidyl peptidase-IV (DPPIV), pyroglutamyl-I (PAP-I), prolyl-imino (PIP); and prolyl oligopeptidase (POP) in seminal plasma, prostate-like and spermatozoa subcellular soluble (SF) and solubilized membrane-bound (MF) fractions of *C. d. terrificus*. Values are means $\pm$ S.E.M. from triplicates of samples (see Materials and Methods). U = picomoles of hydrolyzed substrate per min. Different letters indicate different peptidase activity levels in the same fraction (one-way ANOVA, $P < 0.0001$, Bonferroni $P < 0.001$). Asterisks indicate the higher activity of each peptidase when comparisons were made among all seminal fractions (one-way ANOVA, $P < 0.0003$, Bonferroni $P < 0.05$).

peptides and the synthetic substrates, strongly suggesting that those peptides are natural substrates of seminal peptidases of *C. d. terrificus*.

The semen of the rattlesnake contains SP and spermatozoa, as well as vesicles resembling mammalian PRs. Although ultrastructural analyses are needed to determine peculiarities of these vesicles seen in *C. d. terrificus*, compared with PRs of mammals, they could be considered as PR structures since they were obtained by the same purification procedure employed to obtain human PRs (Fernández *et al.* 2002, Vivacqua *et al.* 2004) and had peptidase activities (Fernández *et al.* 2002), shape (Fernández *et al.* 1997, Vivacqua *et al.* 2004), and size (Olsson & Ronquist 1990, Vivacqua *et al.* 2004) similar to human PRs. Mammalian PRs are membranous vesicles (50–550 nm; 150–200 nm are the most frequent), which contain large amounts of cholesterol, sphingomyelin, calcium, and several enzymes (Kravets *et al.* 2000). They are surrounded by a bilaminar or multilaminar lipoprotein membrane with unusual lipid composition due to quantitative predominance of

cholesterol over phospholipids (Kravets *et al.* 2000). PRs have been related to sperm motility, liquefaction, and immunosuppression (Kravets *et al.* 2000). Since PRs can be identified only after purification procedures it is difficult to infer how frequent they are in *C. d. terrificus*. Another important question that remains to be investigated is the origin of these vesicles in *C. d. terrificus*. In humans, PRs are secreted by the prostate gland (Kravets *et al.* 2000). In bovines, PRs are originated in the seminal vesicles (Agrawal & Vanha-Perttulla 1987). Seminal vesicles secrete proteins, vitamin C, fructose, prostaglandin, and fibrinogen, while the prostate gland secretes a whitish liquid with low viscosity which contains citrate and Ca^{2+} (Minelli *et al.* 1998). It is noteworthy that *C. d. terrificus* possesses PR structures, since squamates do not possess a prostate gland, seminal vesicles, and bulbo-urethral and urethral glands (Sever 2004). The renal sexual segment is known to act as an accessory sex organ (Sever *et al.* 2002), from which those structures could have originated. This renal sexual segment of snakes secretes a complex of glycogen,

mucopolysaccharides, lipids, mucoproteins, and phosphatases (Kühnel & Krisch 1974) in a pattern that can be related to spermatogenic cycle and mating activity (Sever *et al.* 2002). However, since in the present study the semen was collected from the ductus deferens, it would not have mixed with fluids from the renal sexual segment but probably from other areas of major secretory activity in the snake urogenital system, e.g. the ampulla ductus deferentis and the ductuli epididymides of the epididymis (Volsoe 1944), from which the PR structures could have originated. The ductuli epididymides or ductuli efferents have been observed in the snake *Seminatrix pygaea* as efferent ducts from seminiferous tubules of the testis that lead into the epididymis, from which sperm pass into the ductus deferens (Sever 2004). In several mammals, the terminal portion of the ductus deferens is differentiated into an ampulla, which is considered a multifaceted organ, morphologically similar to the seminal vesicles (Riva *et al.* 1982). Among reptiles, the ampulla has been reported only in squamates, being more prominent in lizards than in snakes (Akbarsha *et al.* 2005). The ampulla of squamates and mammals shares the irregular narrow folded epithelium (Sever 2004).

The higher levels of the majority of the examined peptidase activities were found in SP and PR fractions. APN and APB activities were more expressive in all seminal fractions. The conversion of kallidin to BK (Mizutani *et al.* 1993) and inactivation of BK (Barret *et al.* 1998) are, respectively, actions of APN and APB, indicating the importance of the regulation of these peptides to seminal function in *C. d. terrificus*. However, considering our results and the large range of peptides that are susceptible to these peptidases, it is not possible to assure that the high levels of peptidases are restricted only to the kinin system. In other words, the relevance of other evaluated peptidases cannot be ignored. For instance, DPPIV and POP activities were also highly expressed in PR and SP of *C. d. terrificus* respectively. Considering their known hydrolytic effects over endorphin-2 and enkephalin, respectively (Barret *et al.* 1998), and the fact that endogenous opioid peptides seem to have a marked role in seminal physiology (O'Hara *et al.* 1994, Agirregoitia *et al.* 2006), DPPIV and POP activities may have a relevant role in seminal physiology of *C. d. terrificus*.

In conclusion, this is the first report of fractionation and peptidase composition of semen of reptiles, and demonstrated the presence of acid, basic, neutral (puromycin-sensitive and -insensitive), cystyl, dipeptidyl-IV, type I pyroglutamyl, and prolyl-imino APs, as well as POP activities in SP, spermatozoa, PR structures, and whole semen from ductus deferens of *C. d. terrificus*. Metallo-dependent APs inhibited by amastatin and bestatin (mainly those that act on neutral or basic amino acids) had marked levels in whole semen and in all seminal fractions, indicating a high physiological

importance. Ang II, AVT, BK, FPP, and TRH altered the majority of the peptidase activities under study, suggesting them as possible substrates and/or modulators of these peptidases.

Materials and Methods

Animals

Adult male snakes (*C. d. terrificus*, Serpentes, Viperidae, Crotalinae; $n=26$) were captured from their natural environment in the states of São Paulo and Minas Gerais (Brazil) during austral autumn, when testosterone is at its peak (Zacariotti *et al.* 2005) and mating occurs (Almeida-Santos & Salomão 1997). The animals were identified by the Laboratory of Herpetology of the Instituto Butantan and housed individually in wooden cages (inside length $\times$ width $\times$ height of 35 $\times$ 26 $\times$ 22 cm) and acclimated to controlled conditions of temperature (25 °C), humidity (65.3 $\pm$ 0.9%), and photoperiod (12 h light:12 h dark—lights on at 0600 h) in a restricted-access room for a period of 10 days.

After anesthesia with CO₂ exposure for 3 h, the ductus deferens of the reproductive tract were removed by laparotomy (Langlada *et al.* 1994), and semen was obtained from these ducts. The animals were then destined for other studies after euthanizing by decapitation.

The animal and research protocols used in this study are in agreement with the Brazilian Council Directive (COBEA-BRAZIL) and were approved by the Ethics Committee of the Instituto Butantan (193/04).

Collection of *C. d. terrificus* semen

As previously described by Almeida-Santos *et al.* (2004), the ductus deferens were stretched out on polystyrene plates for semen extraction. Pressure with a cell scraper (TPP – Techno Plastic Products AG, Trasadingen, Switzerland) along the whole extension of the two ducts was applied. The total amount of semen obtained was used for the following procedures.

Preparation of soluble and solubilized membrane-bound fractions from whole semen of *C. d. terrificus*

In order to obtain SF and MF, individual samples of whole semen were resuspended in 1 ml of 10 mM Tris–HCl buffer (pH 7.4), homogenized with a Teflon pestle in a glass potter (2 min at 800 r.p.m.) and ultracentrifuged (100 000 *g* for 35 min; Hitachi model HIMAC CP56GII). The resulting supernatants correspond to the SF of whole semen. To avoid contamination with the SF, the resulting pellet was washed three times with the same buffer and was then homogenized (2 min at 800 r.p.m.) in 1 ml of 10 mM Tris–HCl buffer plus 0.1% (v/v) Triton X-100 (Calbiochem, San Diego, CA, USA), and then ultracentrifuged (100 000 *g* for 35 min). The resulting supernatants correspond to the MF of whole semen. All steps were carried out at 4 °C. SF and MF were stored in polystyrene tubes at –80 °C until their use in LDH, protein, and peptidase activities assays.

Fractionation of semen of *C. d. terrificus*

In order to obtain SP, PR, and spermatozoa subcellular SF and MF of *C. d. terrificus* semen, the methodology described by Fernández *et al.* (2002) for human semen was adapted as follows. The pool of whole semen of five animals was homogenized in 1 ml Tris-saline buffer (30 mM Trizma (Sigma) plus 130 mM NaCl (Merck), pH 7.4) and centrifuged (600 *g* for 20 min; Hitachi HIMAC CP56GII). The supernatant contained SP and PR and was reserved for later use, while the pellet contained spermatozoa. To avoid contamination with SP and PR, the pellet was washed with 1 ml Tris-saline buffer and subsequently resuspended in 2.5 ml of the same buffer and centrifuged (600 *g* for 10 min). The resulting supernatant was discarded and the pellet was washed with 2.5 ml Tris-saline buffer and subsequently resuspended in 2.5 ml of the same buffer and centrifuged (1000 *g* for 15 min). The supernatant was discarded, and the resulting pellet contained only spermatozoa. This pellet was homogenized in 2.5 ml 10 mM Tris-HCl buffer (pH 7.4; 2 min at 800 r.p.m.) and stored at -80°C , until use for separation of subcellular SF and MF. Then, the homogenate was sonicated (6 pulses of 30 s, with intervals of 15 s) and ultracentrifuged (100 000 *g* for 35 min). The pellet was reserved and the supernatant was ultracentrifuged (100 000 *g* for 35 min), and the resulting pellet was discarded while the supernatant was considered the spermatozoa subcellular SF. The reserved pellet from the penultimate ultracentrifugation was, then, homogenized at 800 r.p.m. (2 min, in 1 ml 10 mM Tris-HCl buffer) and ultracentrifuged (100 000 *g* for 35 min). After discarding the resulting supernatant, the pellet was homogenized at 800 r.p.m. (2 min, in 1 ml 10 mM Tris-HCl buffer), and was considered as the spermatozoa subcellular MF.

The resulting supernatant from the first centrifugation (SP and PR) was centrifuged (1000 *g* for 20 min) in order to eliminate debris and residual spermatozoa. The resulting pellet was discarded and the supernatant was stored at -80°C . After defrost, this supernatant was ultracentrifuged (100 000 *g* for 120 min). The obtained pellet was reserved and the supernatant was ultracentrifuged at the same conditions. The resulting pellet was discarded and the resulting supernatant contained SP. The reserved pellet from the penultimate ultracentrifugation was washed with 0.6 ml Tris-saline buffer and subsequently resuspended in the same volume. To obtain 0.6 ml more of this resuspension, all needful procedures were repeated with another pool of whole semen of five animals. Both the resuspensions were mixed (1.2 ml), filtered in a membrane filter with a pore size 0.22 μm (Millipore, Bedford, MA, USA), and then submitted to gel filtration in Sephadex G-200 (Sigma; 1.5×30 cm), at a flow rate of 0.16 ml/min, in a column pre-equilibrated with Tris-saline buffer. The dead volume (V_0) was determined by injecting a solution of Blue dextran (Sigma). PR was not retained by the column (Fernández *et al.* 2002) and, then, was collected in the V_0 (fractions 60–120 min or 9.6–19.2 ml), in a volume of about 9.6 ml. The eluate with PR was ultracentrifuged (100 000 *g* for 120 min) and the resultant pellet (PR fraction) was homogenized in 1 ml Tris-saline buffer. All procedures were performed at 4°C . After obtaining SP, PR, SF, and MF of spermatozoa, samples were transferred to polystyrene tubes and maintained at -80°C until the measurements of protein and peptidase activities. Part of PR fraction was destined to the electron microscopy.

Electron microscopy of PR fraction

The PR pellet was fixed for 2 h in a solution of 1.5% glutaraldehyde (Electron Microscopy Sciences-EMS, Hatfield, PA, USA) and 1% paraformaldehyde (Sigma) in 0.1 M cacodylate buffer, pH 7.3, and post-fixed in 1% osmium tetroxide in the same buffer. After dehydration in ethanol graded series, the pellet was embedded in Embed-812 resin (Electron Microscopy Sciences-EMS). Silver ultrathin sections were stained with uranyl acetate and lead citrate, examined and microphotographed (LEO 906E Transmission Electron Microscope, Zeiss, Göttingen, Germany).

Protein

Protein content was measured at 630 nm in triplicates by the method of Bradford (1976; Bio-Rad protein assay). Protein contents were extrapolated by comparison with standard curves of BSA (Sigma) in the same diluent.

LDH

As a marker for the fractionation procedure, LDH activity was determined (Bergmeyer & Brent 1972) in samples of 3 μl of SF and MF in triplicates incubated with 297 μl of NADH (β -NADH, reduced form; Sigma), dissolved in 100 mM phosphate buffer, pH 7.4, containing 1.6 mM sodium pyruvate (Sigma) and 200 mM NaCl. Absorbance was read in 96-well flat bottom microplates, at 0 and 10 min, in Bio-Tek PowerWave X spectrophotometer absorbance reader at 340 nm. Values of LDH activity were obtained by subtracting the absorbance reading at 10 min from time zero of incubation at 37°C , and extrapolated by comparison with a standard curve of NADH dissolved in 100 mM phosphate buffer, pH 7.4, containing 200 mM NaCl. LDH activity was expressed as mmol NADH oxidized/min/mg protein.

Peptidase activity assays

Peptidase activities were quantified on the basis of the amount of 4-methoxy- β -naphthylamine (Sigma; for DPPIV and CAP) or β -naphthylamine (Sigma; for all other peptidases) released (Irazusta *et al.* 2001, Silveira *et al.* 2001, Gasparello-Clemente & Silveira 2002, Gasparello-Clemente *et al.* 2003, Varona *et al.* 2003), which is the result of the enzyme activity of 20–50 μl samples incubated with prewarmed substrate solution at concentrations of 0.125 mM (APA, APN-PS, APN-PI, CAP, PAP, PIP, and POP), 0.2 mM (DPPIV), and 0.5 mM (APB) in respective 0.05 M buffers containing BSA 0.1 mg/ml in 96-well flat bottom microplates for 30 min at 37°C in a total volume of 300 μl . The content of naphthylamine was estimated fluorometrically (microplate fluorescence reader Bio-Tek FL600FA) at 460/40 nm emission wavelength and 360/40 nm excitation wavelength. The fluorescence value obtained at time zero was subtracted from values at time 30 min and the relative fluorescence was then converted to picomoles of β -naphthylamine (Sigma) or 4-methoxy- β -naphthylamine (Sigma) by comparison with a correspondent standard curve, which was dissolved in the same diluent as the incubation.

Peptidase activities were expressed as picomoles of hydrolyzed substrate per min (U) per milligram of protein, in which

the existence of a linear relationship between time of hydrolysis and protein content in the fluorometric assay was a previous condition. Considering enzyme activity measures as a comparative tool, the possible unspecific degradation during homogenization was not considered.

The following substrates and conditions were used:

- APA, L-aspartic acid α -(β -naphthylamide; Sigma; solubilized in 0.012 M NaOH) in Tris-HCl buffer, pH 7.4, with 1 mM MnCl_2 ;
- APB, L-arginine β -naphthylamide (Sigma; solubilized in H_2O) in phosphate buffer, pH 6.5, with 150 mM NaCl, and 0.02 mM puromycin (Sigma);
- APN, L-alanine- β -naphthylamide (Sigma; solubilized in 0.012 M HCl) in phosphate buffer, pH 7.4, with 1 mM DTT (Sigma), with or without puromycin – APN-PI activity is the result of the incubation with puromycin, while APN-PS is the result of values of incubates without puromycin minus those with puromycin;
- CAP, H-Cys-4-methoxy- β -naphthylamide (Bachem Bioscience Inc., Torrance, CA, USA; solubilized in 0.012 M HCl) Tris-maleate, pH 5.9;
- DPPIV, Glycyl-Proline-4-methoxy- β -naphthylamide (Bachem Bioscience Inc.; solubilized in dimethyl sulfoxide, DMSO (Sigma)) in Tris-HCl buffer, pH 8.3;
- PAP-I, L-pyrogutamic acid- β -naphthylamide (Sigma; solubilized in DMSO) in phosphate buffer, pH 7.4, with 2 mM DTT (DTT inhibits PAP-II and activates PAP-I (O'Cuinn *et al.* 1990)) and 2 mM EDTA (Merck);
- PIP, L-proline- β -naphthylamide (Sigma; solubilized in DMSO) in phosphate buffer, pH 7.4;
- POP, Z-Gly-Pro- β -naphthylamide (Bachem Bioscience Inc.; solubilized in DMSO) in phosphate buffer, pH 7.4, with 2 mM DTT.

Characterization of peptidase activities in SF and MF of whole semen

The characteristics of peptidase activities with their classical inhibitors

The susceptibilities of APA, APB, APN, CAP, DPPIV, PAP-I, and PIP peptidase activities were comparatively evaluated in a SF and/or MF pool of whole semen from 11 animals with: amastatin ([[(2S,3R)-3-Amino-2- -5-methylhexanoyl]-Val-Val-Asp-OH.HCl; Bachem Bioscience Inc.): 0.026 mM in acetic acid 8%; bestatin ([[(2S,3R)-3-Amino-2-hydroxy-4- phenylbutanoyl]-L-leucine; Bachem Bioscience Inc.): 0.026 mM in acetic acid 8%; diprotin A (H-Lle-Pro-Lle-OH; Bachem Bioscience Inc.): 0.026 mM in deionized water; puromycin (3'-[α -Amino-*p*-methoxyhydrocin-namamido]-3-deoxy-*N,N*-dimethyladenosine): 0.026 mM in deionized water. The samples were incubated in the presence or absence of those substances, for 30 min, at 25 °C, and peptidase activities were quantified.

The characteristics of peptidase activities with peptide hormones, DTT, EDTA, and MnCl_2

The susceptibilities of APA, APB, APN, CAP, DPPIV, PAP-I, and PIP peptidase activities were comparatively evaluated in a SF

and/or MF pool of whole semen from 11 animals with: 0.315 μM Ang II (Sigma); 0.314 μM AVT (reptilian analog of OXT and AVP; Sigma); 0.310 μM BK (Sigma); 0.94 μM FPP (Sigma); 0.94 μM FPP + 1.8 mM CaCl_2 ; 0.911 μM TRH (Sigma); 0.97 mM DTT; 2 mM EDTA; 1 mM MnCl_2 . The samples were incubated in the presence or absence of these substances, for 30 min, at 25 °C, and peptidase activities were subsequently quantified.

Statistical analysis

Data were analyzed statistically using GraphPad Prism and Instat softwares (GraphPad Software Inc., San Diego, CA, USA). Regression analysis was performed to obtain standard curves. ANOVA, followed by Bonferroni test, compared values among all peptidase activities in the same seminal fraction. Student's *t*-test was performed to compare values of LDH between SF and MF, and peptidase activities in different incubation conditions relative to the respective controls. Differences were considered statistically significant at a minimum level of $P < 0.05$.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP, Brazil (research grant no. 04/14971-9). C E M (05/03745-0) and S C Y (07/06829-6) were recipients of a FAPESP fellowship.

Acknowledgements

The authors thank Valdir José Germano for his efficient technical assistance.

References

- Agirregoitia E, Valdivia A, Carracedo A, Casis L, Gil J, Subiran N, Ochoa C & Irazusta J 2006 Expression and localization of delta-, kappa-, and mu-opioid receptors in human spermatozoa and implications for sperm motility. *Journal of Clinical Endocrinology and Metabolism* **91** 4969–4975.
- Agrawal Y & Vanha-Perttula T 1987 Effect of secretory particles in bovine seminal vesicles secretion on sperm motility and acrosome reaction. *Journal of Reproduction and Fertility* **79** 409–419.
- Akbarsha MA, Tamilarasan V, Kadalmani B & Daisy P 2005 Ultrastructural evidence for secretion from the epithelium of ampulla ductus deferentis of the fan-throated lizard *Sitana ponticeriana* Cuvier. *Journal of Morphology* **266** 94–111.
- Albiston AL, Fernando R, Ye S, Peck GR & Chai SY 2004 Alzheimer's, angiotensin IV and an aminopeptidase. *Biological & Pharmaceutical Bulletin* **27** 765–767.
- Almeida-Santos SM & Salomão MG 1997 Long-term sperm storage in the neotropical rattlesnake *Crotalus durissus terrificus* (Viperidae: Crotalinae). *Japanese Journal of Herpetology* **17** 46–52.

- Almeida-Santos SM, Laporta-Ferreira IL, Antoniazzi MM & Jared C 2004 Sperm storage in males of the snake *Crotalus durissus terrificus* (Crotalinae: Viperidae) in southeastern Brazil. *Comparative Biochemistry and Physiology, Part A* **139** 169–174.
- Amory JK & Bremner WJ 2003 Regulation of testicular function in men: implications for male hormonal contraceptive development. *Journal of Steroid Biochemistry and Molecular Biology* **85** 357–361.
- Assinder SJ, Carey M, Parkinson T & Nicholson HD 2000 Oxytocin and vasopressin expression in the ovine testis and epididymis: changes with the onset of spermatogenesis. *Biology of Reproduction* **63** 448–456.
- Barret AJ, Rawlings ND & Woessner JF 1998 *Handbook of Proteolytic Enzymes*. London: Academic Press.
- Bergmeyer HU & Brent E 1972 Assay with pyruvate and NADH. In: *Methods in Enzymatic Analysis*, vol 2, pp 574–577. Ed HU Bergmeyer. London: Academic Press Inc.
- Blaukat A 2003 Structure and signalling pathways of kinin receptors. *Andrologia* **35** 17–23.
- Bradford MM 1976 A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72** 248–254.
- Cadel S, Pierotti AR, Foulon T, Crémillon C, Barré N, Segrétain D & Cohen P 1995 Aminopeptidase-B in the rat testes: isolation, functional properties and cellular localization in the seminiferous tubules. *Molecular and Cellular Endocrinology* **110** 149–160.
- Chatterjee S, Laloraya M & Kumar PG 1997 Free radical-induced liquefaction of ejaculated human semen: a new dimension in semen biochemistry. *Archives of Andrology* **38** 107–111.
- Cockle SM, Prater GV, Thetford CR, Hamilton C, Malone PR & Mundy AR 1994 Peptides related to thyrotrophin-releasing hormone (TRH) in human prostate and semen. *Biochimica et Biophysica Acta* **1227** 60–66.
- Constam DB, Tobler AR, Resing-Ehl A, Kemler I, Hersh LB & Fontana A 1995 Puromycin-sensitive aminopeptidase. Sequence analysis, expression, and functional characterization. *Journal of Biological Chemistry* **270** 26931–26939.
- Cummins PM & O'Connor B 1998 Pyroglutamyl peptidase: an overview of the three known enzymatic forms. *Biochimica et Biophysica Acta* **1429** 1–17.
- Davison JM, Sheills EA, Phillips P, Barron WM & Lindheimer MD 1993 Metabolic clearance of vasopressin and an analogue resistant to vasopressinase in human pregnancy. *American Journal of Physiology* **264** F348–F353.
- El-Haggag S, El-Ashmawy S, Attia A, Mostafa T, Roaiah MM, Fayed A, Ghazi S, Zohdy W & Roshdy N 2006 Beta-endorphin in serum and seminal plasma in infertile men. *Asian Journal of Andrology* **8** 709–712.
- Fernández JA, Heeb MJ, Radtke K-P & Griffin JH 1997 Potent blood coagulant activity of human semen due to prostatic-bound tissue factor. *Biology of Reproduction* **56** 757–763.
- Fernández D, Valdivia A, Irazusta J, Ochoa C & Casis L 2002 Peptidase activities in human semen. *Peptides* **23** 461–468.
- Fraser LR, Adeoya-Osiguwa SA, Baxendale RW & Gibbons R 2006 Regulation of mammalian sperm capacitation by endogenous molecules. *Frontiers in Bioscience* **11** 1636–1645.
- García-Horsman JA, Männistö PT & Venäläinen JI 2007 On the role of prolyl oligopeptidase in health and disease. *Neuropeptides* **41** 1–24.
- Gaspardo-Clemente E & Silveira PF 2002 Fluorometric assay using naphthylamide substrates for assessing novel venom peptidase activities. *Toxicon* **40** 1617–1626.
- Gaspardo-Clemente E, Casis L, Varona A, Gil J, Irazusta J & Silveira PF 2003 Aminopeptidases in visceral organs during alterations in body fluid volume and osmolality. *Peptides* **24** 1367–1372.
- Gibson AM, McDermott JR, Lauffart B & Mantle D 1989 Specificity of action of human brain alanyl aminopeptidase on Leu-enkephalin and dynorphin-related peptides. *Neuropeptides* **13** 259–262.
- Green CM, Cockle SM, Watson PF & Fraser LR 1996 A possible mechanism of action for fertilization promoting peptide, a TRH-related tripeptide that promotes capacitation and fertilizing ability in mammalian spermatozoa. *Molecular Reproduction and Development* **45** 244–252.
- Gribbins KM, Mills EM & Sever DM 2007 Ultrastructural examination of spermiogenesis within the testis of the ground skink, *Scincella laterale* (Squamata, Sauria, Scincidae). *Journal of Morphology* **268** 181–192.
- Irazusta J, Silveira PF, Gil J, Varona A & Casis L 2001 Effects of hydrosaline treatments on prolyl endopeptidase activity in rat tissues. *Regulatory Peptides* **101** 141–147.
- Irazusta J, Valdivia A, Fernández D, Agirreagoitia E, Ochoa C & Casis L 2004 Enkephalin-degrading enzymes in normal and subfertile human semen. *Journal of Andrology* **25** 733–739.
- Kawata S, Takayama S, Ninomiya K & Makisumi S 1980 Porcine liver aminopeptidase B. Substrate specificity and inhibition by amino acids. *Journal of Biochemistry* **88** 1601–1605.
- Kravets FG, Lee J, Singh B, Trocchia A, Pentyala SN & Khan SA 2000 Prostatomes: current concepts. *Prostate* **43** 169–174.
- Kugler P 1982 Aminopeptidase A is angiotensinase A. II. Biochemical studies on aminopeptidase A and M in rat kidney homogenate. *Histochemistry* **74** 247–261.
- Kühnel W & Krisch B 1974 On the sexual segment of the kidney in the snake (*Natrix natrix*). *Cell and Tissue Research* **148** 417–429.
- Langlada FG, Santos S & Ferreira IL 1994 Techniques of artificial insemination in *Crotalus durissus terrificus* (Viperidae-Crotalinae). *Brazilian Journal of Veterinary Research and Animal Science* **31** 141–144.
- Minelli A, Moroni M, Martínez E, Mezzasoma I & Ronquist G 1998 Occurrence of prostatic-like membrane vesicles in equine seminal plasma. *Journal of Reproduction and Fertility* **114** 237–243.
- Minelli A, Allegrucci C, Mezzasoma I, Ronquist G, Lluís C & Franco R 1999 CD26 and adenosine deaminase interaction: its role in the fusion between horse membrane vesicles and spermatozoa. *Biology of Reproduction* **61** 802–808.
- Mizutani S, Goto K, Nomura S, Ino K, Goto S, Kikkawa F, Kurauchi O, Goldstein G & Tomoda Y 1993 Possible action of human placental aminopeptidase N in feto-placental unit. *Research Communications in Chemical Pathology and Pharmacology* **82** 65–80.
- Nakamura K, Fujiwara H, Higuchi T, Honda T, Yamada S, Nakayama T, Fujita J, Maeda M, Tachibana T, Suganami H *et al.* 1998 Bestatin, an aminopeptidase inhibitor, promotes follicular growth and ovulation suppressed by stress in mice. *Journal of Endocrinology* **45** 547–553.
- O'Cuinn G 1998 Peptide metabolism in cytoplasm of brain cells. *Biochemical Society Transactions* **26** 279–292.
- O'Cuinn G, O'Connor B & Elmore M 1990 Degradation of thyrotropin-releasing hormone and luteinising hormone-releasing hormone by enzymes of brain tissue. *Journal of Neurochemistry* **54** 1–13.
- O'Hara BF, Donovan DM, Lindberg I, Brannock MT, Ricker DD, Moffatt CA, Klaunberg BA, Schindler C, Chang TS, Nelson RJ *et al.* 1994 Proenkephalin transgenic mice: a short promoter confers high testis expression and reduced fertility. *Molecular Reproduction and Development* **38** 275–284.
- Olivo RA, Nascimento NG, Teixeira CFP & Silveira PF 2008 Methotrexate and cyclosporine treatments modify the activities of dipeptidyl peptidase IV and prolyl oligopeptidase in murine macrophages. *Clinical & Developmental Immunology* **2008** 794050 (doi: 10.1155/2008/794050).
- Olsson I & Ronquist G 1990 Nucleic acid association to human prostatic prostates. *Archives of Andrology* **24** 1–10.
- Ramírez M, Arechaga G, García S, Sánchez B, Lardelli P & Gandarias JM 1990 Mn²⁺-activated aspartate aminopeptidase activity, subcellular localization in young and adult rat brain. *Brain Research* **522** 165–167.
- Riva A, Testa-Riva F, Usai E & Cossu M 1982 The ampulla ductus deferentis in man, as viewed by SEM and TEM. *Archives of Andrology* **8** 157–164.
- Rónai AZ, Timár J, Makó E, Erdő F, Gyarmati Z, Tóth G, Orosz G, Fürst S & Székely JI 1999 Diprotin A, an inhibitor of dipeptidyl aminopeptidase IV (EC 3.4.14.5) produces naloxone-reversible analgesia in rats. *Life Sciences* **64** 145–152.
- Sastry BV, Janson VE & Owens LK 1991 Significance of substance P- and enkephalin-peptide systems in the male genital tract. *Annals of the New York Academy of Sciences* **632** 339–353.
- Sato Y 2003 Aminopeptidases and angiogenesis. *Endothelium* **10** 287–290.
- Sever DM 2004 Ultrastructure of the reproductive system of the black swamp snake (*Seminatrix pygmaea*): IV. Occurrence of an ampulla ductus deferentis. *Journal of Morphology* **262** 714–730.
- Sever DM, Stevens RA, Ryan TJ & Hamlett WC 2002 Ultrastructure of the reproductive system of the black swamp snake (*Seminatrix pygmaea*): III. Sexual segment of the male kidney. *Journal of Morphology* **252** 238–254.

- Silveira PF, Irazusta J, Gil J, Agirregoitia N & Casis L** 2001 Interactions among challenges of hydromineral balance, angiotensin-converting enzyme, and cystine aminopeptidase. *Peptides* **22** 2137–2144.
- Siviter RJ & Cockle SM** 1995 Peptides related to thyrotrophin-releasing hormone are degraded in seminal plasma by an enzyme similar to prolil endopeptidase. *Journal of Endocrinology* **144** 61–66.
- Subirán N, Agirregoitia E, Valdivia A, Ochoa C, Casis L & Irazusta J** 2008 Expression of enkephalin-degrading enzymes in human semen and implications for sperm motility. *Fertility and Sterility* **89** 1571–1577.
- Takahashi T, Ikai A & Takahashi K** 1989 Purification and characterization of proline-beta-naphthylamidase, a novel enzyme from pig intestinal mucosa. *Journal of Biological Chemistry* **264** 11565–11571.
- Talbot P** 1991 Compartmentalization in the acrosome. In *Comparative Spermatology – 20 Years After*, pp 255–259. Ed. B Baccetti. New York: Raven Press.
- Tourmente M, Cardozo GA, Guidobaldi HA, Giojalas LC, Bertona M & Chiaraviglio M** 2007 Sperm motility parameters to evaluate the seminal quality of *Boa constrictor occidentalis*, a threatened snake species. *Research in Veterinary Science* **82** 93–98.
- Valdivia A, Irazusta J, Fernández D, Múgica J, Ochoa C & Casis L** 2004 Pyroglutamyl peptidase I and prolil endopeptidase in human semen: increased activity in necrozoospermia. *Regulatory Peptides* **122** 79–84.
- Varona A, Silveira PF, Irazusta A, Valdivia A & Gil J** 2003 Effects of changes in hydromineral balance on rat brain aspartyl, arginyl, and alanyl aminopeptidase activities. *Hormone and Metabolic Research* **35** 36–42.
- Vivacqua A, Siciliano L, Sabato M, Palma A & Carpino A** 2004 Prostatome as zinc ligands in human seminal plasma. *International Journal of Andrology* **27** 27–31.
- Volsoe H** 1944 Structure and seasonal variation of the male reproductive organs of *Vipera berus* (L.). *Spolia Zoologica Musei Hauniensis V. Skrifter* **5** 1–157.
- Zacariotti RL** 2004 Estudo longitudinal do espermograma e dos níveis de testosterona sérica de cascavel (*Crotalus durissus terrificus*, Laurenti, 1768) proveniente da natureza do Estado de São Paulo. *Dissertação (Mestrado em Reprodução Animal)*. Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, São Paulo.
- Zacariotti RL, Betkowsky SE, Grego KF, Santanna SS, Fernandes W, Oliveira CA & Guimaraes MABV** 2005 Spermogram and serum testosterone levels in wild caught Brazilian rattlesnakes (*Crotalus durissus terrificus*). *Proceedings of the Annual Conference of The American Association of Zoo Veterinarians*, 2005, Omaha, NE, USA pp 320.

Received 7 May 2008

First decision 30 May 2008

Revised manuscript received 1 September 2008

Accepted 10 September 2008