Effects of HSPA8, an evolutionarily conserved oviductal protein, on boar and bull spermatozoa

Roslyn M A Elliott¹, Rhiannon E Lloyd¹, Alireza Fazeli², Edita Sostaric², A Stephen Georgiou², Nana Satake^{1,3}, Paul F Watson³ and William V Holt¹

¹Institute of Zoology, Regent's Park, London NW1 4RY, UK, ²Academic Unit of Reproductive and Developmental Medicine, The University of Sheffield, Level 4, The Jessop Wing, Tree Root Walk, Sheffield S10 2SF, UK and ³The Royal Veterinary College, Royal College Street, London NW1 0TU, UK

Correspondence should be addressed to W V Holt; Email: bill.holt@ioz.ac.uk

Abstract

Previous studies have shown that a soluble protein fraction derived from preparations of apical plasma membrane (APM) of the oviductal epithelium enhances the *in vitro* survival of mammalian spermatozoa. Here, we show that the survival enhancing property of the soluble protein fraction seems to depend significantly upon heat shock 70 kDa protein 8 (HSPA8 previously known as HSPA10). The following findings in the present study enabled us to draw this conclusion: first, using proteomic analysis, we identified a subset of 70 kDa oviductal surface proteins that bound to spermatozoa, one of which was HSPA8. Second, pre-treatment of the soluble protein fraction with anti-HSPA8 antibody reduced the 24 h (at 39 °C) sperm survival enhancement effect normally induced by the presence of 200 µg/ml soluble APM proteins. Third, complementary experiments showed that substituting the soluble protein fraction with bovine recombinant HSPA8 (0.5–2 µg/ml) also elicited the sperm survival effect. Finally, we also tested the effect of bovine recombinant HSPA8 on bull spermatozoa and found similar, dose-responsive, sperm survival promoting effects. The conserved nature of HSPA8 between mammalian species suggests that this protein may represent a common biological mechanism for the maintenance of sperm survival in the oviduct.

Reproduction (2009) **137** 191–203

Introduction

Sperm storage in the female reproductive tract has been extensively documented in a range of vertebrates that includes sharks (Pratt & Tanaka 1994), salamanders (Sever & Brizzi 1998), snakes and turtles (Galbraith 1993, Gist & Congdon 1998) as well as many birds and mammals. Some of these species, especially the snakes and turtles, store spermatozoa for several months, or even years, prior to fertilization. In general, mammals are only able to store spermatozoa for a few days in the female reproductive tract (Rodriguez-Martinez *et al.* 2005, Holt *et al.* 2006, Rodriguez-Martinez 2007), although some bat species have evolved successful means of storing spermatozoa for several months over winter in the uterus or oviduct (Racey & Potts 1970, Racey 1979, Racey *et al.* 1987, Holt *et al.* 2006). The mechanisms involved in prolonging sperm survival in the female reproductive tract have remained largely elusive, although the widely observed (Hunter 1981, Pacey *et al.* 1995, Hunter & Rodriguez-Martinez 2004) formation of intimate associations between spermatozoa and the oviductal epithelial surface suggests that membrane–membrane contact is likely to be important.

Such inter-membrane contact is likely to stimulate mutual cell signalling dialogues between the spermatozoa and oviductal cells, with resultant effects on the activity of both cell types. Oviductal epithelial cells (OEC) are now known to respond by modulating gene transcription and protein synthesis (Georgiou *et al.* 2005) while (boar) spermatozoa respond to the presence of oviductal proteins with reduced motility and an accelerated increase in intracellular pH (Satake *et al.* 2006).

Co-culture of spermatozoa with OEC has repeatedly been shown to prolong sperm viability in some species, including pigs. In this laboratory, we have consistently demonstrated the enhancement of boar sperm viability by a soluble fraction of oviductal apical plasma membrane (sAPM) proteins (Elliott *et al.* 2001, Fazeli *et al.* 2003, Holt *et al.* 2005), derived from APM vesicles, as originally described by Smith & Nothnick (1997). In bioassays involving incubation of boar spermatozoa with sAPM for 24 h at 39 °C, sperm viability was enhanced in a dose-dependent manner (Fazeli *et al.* 2003). Moreover, the enhancement effect was abolished if sAPM proteins had been denatured (Fazeli *et al.* 2003).

These results confirmed that one or more proteins within the sAPM contribute to the enhancement of sperm viability.

However, sAPM represents a complex mixture of proteins, any of which might be active at low concentration. This might explain why a relatively high concentration of sAPM (200 µg/ml) is necessary in order to demonstrate the viability enhancing effect. In a previous publication (Holt *et al.* 2005), we described in detail the preparation method for sAPM and demonstrated that we could distinguish oviduct epithelium surface-exposed proteins by the use of an *in situ* biotinylation technique. In the current study, we have again used *in situ* biotinylated oviductal sAPM preparation and have tracked the binding of this protein preparation to the surface of co-incubated spermatozoa. We have also studied the sperm binding topology in some detail using confocal microscopy and used proteomic analyses to identify heat shock 70 kDa protein 8 (HSPA8 previously known as HSPA10, HSC71, HSC70 and HSP73) as a probable significant functional component of the sAPM. Ultimately, using bovine recombinant HSPA8 protein, we replicated the sperm-survival enhancement effect of sAPM, not only in boar spermatozoa but also in bull spermatozoa.

Results

Association of sAPM proteins with spermatozoa

Two different approaches were used to examine the association of surface oviductal sAPM proteins and spermatozoa. In the first experiment, *in situ* biotinylated sAPM proteins were bound to Immobilized NeutrAvidin coated beads, which were then co-incubated with spermatozoa to see whether any physical binding occurred. The control sample consisted of NeutrAvidin-coated beads with no surface-bound sAPM. The second experiment involved visualization of sAPM proteins by confocal microscopy. Aliquots of sAPM were directly labelled with FITC to provide visible evidence of proteins associating with spermatozoa.

Examination of live images of the sperm-bead interactions revealed that unexpectedly small numbers of spermatozoa became bound to the bead surfaces. Even those that became bound tended to be released again after a few seconds. This lack of firm binding was not caused by any shortage of spermatozoa in the surrounding media and may therefore indicate that some degree of sperm selection was involved. Visual estimates nevertheless showed that the number of spermatozoa associated with sAPM-treated beads (regardless of whether the sAPM had been biotinylated *ex-situ* or *in-situ*) was about fivefold greater than became bound to untreated beads (five sperm/bead for *ex-situ* labelled sAPM; 4.4 sperm/bead for *in situ*

labelled sAPM; control values were 0.75 and 0.82 sperm/bead respectively; $P=0.02$). As it was not possible to examine the whole of the bead surface by video microscopy, some sperm-bead preparations (made using live spermatozoa labelled with SYBR14) were fixed and examined by confocal microscopy to check the uniformity and low level of binding. Slices through an sAPM-treated bead and a control bead are shown in Fig. 1A and B respectively. The confocal microscopy confirmed that relatively few spermatozoa were bound to the bead surfaces.

Fluorescence microscopy of spermatozoa incubated with FITC-labelled sAPM showed that binding mostly occurred on the apical segment of the acrosome in plasma-membrane intact propidium iodide (PI–ve) spermatozoa (mean $\pm$ S.E.M.: $63.1 \pm 6.1\%$; $n=8$ boars), although some spermatozoa exhibited more extensive binding over the whole of the sperm head or over the

Figure 1 Confocal images through a single sAPM-coated bead (A) incubated with SYBR-14 labelled spermatozoa and a control, untreated, bead (B); relatively few spermatozoa were bound to the bead (see white arrows). Scale bars = 100 µm. (C) shows that many sperm heads bind FITC- labelled sAPM at the apical region of the acrosome (white arrows), while a few spermatozoa bind sAPM over their entire head, middle-piece and tail. Scale bar = 20 µm.

head and whole or part of the flagellum (mean $\pm$ S.E.M.: $18.3 \pm 3.24\%$; $n=8$ boars). Approximately 18% spermatozoa (mean $\pm$ S.E.M.: $18.6 \pm 5.1\%$; $n=8$ boars) did not appear to bind any of the labelled sAPM at all. Examples are presented in Fig. 1C.

Identification of *in situ* biotinylated oviductal epithelial proteins that bind to spermatozoa

Immunoblotting of the combined solubilized sperm membrane protein fraction with sperm surface-bound biotin-labelled sAPM demonstrated specific binding of a protein fraction of ~ 70 kDa. In order to identify this protein(s), the corresponding protein band from a series of replicate SDS-PAGE gels was excised for proteomic analysis.

Figure 2A presents a PAGE gel showing that while many proteins were contained within the *in situ* biotinylated sAPM fraction, after incubation of the

sAPM with spermatozoa only one band (MWt ~ 70 kDa) was strongly detectable on the blot (Fig. 2B, Lane 1). Lane 2 in this figure shows that when spermatozoa were incubated with APM (i.e. non-solubilized biotinylated APM), they bound not only this protein band, but also another with a molecular weight of about 97 kDa (Fig. 2B, lane 2).

To identify the proteins contained within the sperm-binding 70 kDa band, the relevant band was excised (Fig. 2C) from 16 lanes of two gels prepared in parallel and their protein composition analyzed in detail. The proteomic analysis identified 19 proteins (listed in Table 1) within this band. It was not feasible to investigate all the 19 proteins in detail, therefore, as a first step we chose to investigate one of these proteins, namely HSPA8, in more detail. Western blotting showed a strong immunoreactivity between sAPM and anti-HSPA8, confirming the presence of HSPA8 in sAPM.

Neutralization of sAPM with anti-HSPA8

In view of the ability of proteins within the sAPM to interact with spermatozoa, the next step was to see whether the ability of sAPM to enhance sperm viability would be compromised by adsorption with anti-HSPA8. A preliminary experiment showed that anti-HSPA8 diluted to $1 \mu\text{g/ml}$ largely prevented the binding of FITC-labelled sAPM to spermatozoa. This dilution factor was therefore used in tests aimed at blocking the sAPM effects on sperm viability. When the 24 h bioassays were repeated using sAPM and sAPM treated with anti-HSPA8, it was found that sperm survival was significantly higher in the former compared with the latter ($P=0.013$; Fig. 3A).

Figure 2 (A) One-dimensional protein profile of sAPM (both lanes); the protein band at 70 kDa (arrow) corresponds to the positively stained band at 70 kDa in the blot illustrated in (B) (Lane 1); this band was excised from a series of gels and subjected to proteomic analysis. The blot (B) (lane 1) indicates that the 70 kDa sAPM protein was transferred to spermatozoa during sperm-sAPM incubation. A second band of ~ 90 kDa (B, lane 1; arrow) was also transferred to spermatozoa, but was not investigated further. Figure B (Lane 2) shows that the same two subsets of APM proteins were also transferred to spermatozoa during incubation. Lane 3 shows the biotinylation profile of *in situ* biotinylated sAPM; Lanes 4 and 5 show the absence of the biotinylation signal in unlabelled sAPM proteins and spermatozoa respectively. (C). PAGE protein profile of sAPM (both lanes). (D) A western blot of sAPM probed with an antibody against HSPA8.

Effects of HSPA8 on boar spermatozoa incubated for 24 h at 39 °C

To test the effects of HSPA8 itself on boar sperm survival at 39 °C under capacitating conditions, washed spermatozoa were incubated in Tyrode's medium with lactate and pyruvate (TLP) containing 0, 2, 4, 8 and $16 \mu\text{g/ml}$ HSPA8 for 24 h and then evaluated for its viability. There was a significant overall enhancement of sperm survival, represented by a 15% increase in viability index, due to the presence of HSPA8 ($P=0.007$). However, this beneficial effect appeared to be optimal at $2 \mu\text{g/ml}$ ($P=0.0007$; $F_{1/24}=14.9$) and there was no further benefit to be gained by further increasing the concentration of protein.

This outcome was pursued further by repeating the bioassays, but using lower concentrations of HSPA8 (0, 0.25, 0.5, 2 and $8 \mu\text{g/ml}$). As an additional comparison, the spermatozoa were also incubated in the presence of $200 \mu\text{g/ml}$ sAPM to evaluate the effectiveness of HSPA8 against the original oviductal preparation. The beneficial

Table 1 Protein components of the sperm-binding 70 kDa sAPM band.

Protein number	Protein name	Previous name; aliases. Protein family	Accession number (Uniprot)	MOWSE score (MASCOT)	Theoretical protein MW (kDa)	Peptides identified	Sample peptide sequence(s)	Sample peptide e-value(s)
1	Heat shock protein 90 kDa beta (Grp94), member 1 (HSP90B1)	TRA1; GP96 and GRP94. HSP90 protein family	A53211	445	92456	11	K.LIINSLYK.N, K.IYFMAGSSR.K	1.4, 1.5
2	Heat shock protein 90 kDa α (cytosolic), class A member 1 (HSP90AA1)	HSPC1, HSPCA; HSP89, HSP90, FLJ31884, HSP90N. HSP90 protein family	Q2VPJ6_HUMAN	74	68329	3	R.APFDLFENR.K, R.GVVDSDELPLNISR.E	5.3, 0.13
3	Heat shock protein 90 kDa α (cytosolic), class B member 1 (HSP90AB1)	HSPC2, HSPCB. HSP90 protein family	Q1PSW2_RAT	114	83230	3	R.ELISNASDALDK.I, R.GVVDSDELPLNISR.E	0.00062, 0.13
4	Heat shock 70 kDa protein 5 (glucose-regulated protein, 78 kDa) (HSPA5)	GRP78; BIP. HSPA4 protein family	HHRTGB	357	72302	9	K.ITITNDQNR.L, R.NELESYAYSLK.N	2.7, 0.075
5	Heat shock 70 kDa protein 8 (HSPA8)	HSPA10; HSC71, HSC70, HSP73. HSP70 protein family	1NGB	54	71241	1	R.IINQPTAAAIAYGLDK.K	0.0063
6	Clusterin (CLU)	CLI, APOJ; SGP-2, SP-40, TRPM_2, KUB1. CLU protein family	A42108	155	51741	6	K.SLLSSLEEAK.K, R.ASNIMDELQDR.F	0.28, 31
7	Activated leukocyte cell adhesion molecule (ALCAM)	CD166, MEMD	Q1HGM8_HUMAN	114	65061	2	R.SSNTYTLTDVR.R, K.SVQYDDVPEYK.D	0.00035, 0.032
8	Sec23 homolog B (<i>S. cerevisiae</i>) (SEC23B)		SEC23B	50	86407	2	K.IDMNLTDLLGELQR.D	0.16
9	Gelsolin (GSN)	Gelsolin protein family	FAHUP	145	85644	3	K.TGAQELLR.V, K.AGALNSNDAFVLK.T	1.8, 0.035
10	Oviductal glycoprotein 1, 120 kDa (mucin 9, oviductin) (OVGP1)	MUC9; CHIT5. Glycosyl hydrolase 18 family	AAA85445	280	58482	7	K.LLMGFPTYGR.T, K.SSAYTMNYWR.K	2.6, 0.043
11	Ezrin (EZR)	VIL2. Ezrin-radixin-moesin protein family	Q8C6P2_MOUSE	93	34989	2	K.LIMGFPTYGR.N, K.TLLSIGGWNFGTSR.F	2.6, 0.00038
			EZRI_BOVIN	335	68586	9	K.IALLEEAR.R, R.ALKLEER.K	1.6, 0.01
12	Tropomyosin 1 (TPM1)	Tropomyosin protein family	EZRI_MOUSE	261	69233	8	K.IALLEEAR.R, K.ALQLEER.R	1.6, 0.01
			TMRTF1	78	32938	2	R.IQLVEEELDR.A, K.TIDDLLEETLASAK.E	0.016, 2.8
13	Tropomyosin 3 (TPM3)	Tropomyosin protein family	Q59GR8_HUMAN	70	34382	2	R.IQLVEEELDR.A	0.016
			Q5VU71_HUMAN	118	32799	3	R.IQLVEEELDR.A	0.016

Protein number	Protein name	Previous name; aliases. Protein family	Accession number (Uniprot)	MOWSE score (MASCOT)	Theoretical protein MW (kDa)	Peptides identified	Sample peptide sequence(s)	Sample peptide e-value(s)
14	Keratin 1 (KRT1)	EHK1; KRT1A. Intermediate filament family	K2C1_HUMAN	307	65847	6	K.IEISELNR.V; K.YEELQL-TAGR.H	0.022, 0.4
15	Keratin 2 (KRT2)	KRT2A; KRTE. Intermediate filament family	K2C1_MOUSE	154	65052	3	R.TNAENEFVTIK.K; R.FLEQQNKVLQTK.W	0.00024, 0.0012
			Q9D2K8_MOUSE	154	65586	3	R.TNAENEFVTIK.K; R.FLEQQNQVLQTK.W	0.00024, 0.0012
			Q4VAQ2_HUMAN	172	65393	4	R.NLDLSIAEVK.A; K.VDLLNQEIEFLK.V	1.3, 0.69
16	Keratin 4 (KRT4)	CYK4; CK4, K4. Intermediate filament family	K2C4_RAT	92	57285	2	K.NEISELNR.M	0.03
17	Keratin 6A (KRT6A)	KRT6C, KRT6D; CK6C, K6C, CK6D, K6D. Intermediate filament family	Q8BIS2_MOUSE	82	60045	2	R.SLNLDLSIAEVK.A	0.35
18	Keratin 6B (KRT6B)	KRTL1. Intermediate filament family	KRHUEA	76	60008	2	R.NLDLSIAEVK.A	1.3
			I59009	67	60045	1	R.TDAENEFVTIK.K	0.00031
			I61771	67	60030	2	K.YEELQTAGR.H; R.NLDLSIAEVK.A	0.4, 1.3
19	Keratin 77 (KRT77)	KRT1B. Intermediate filament family	CAD91892	92	61688	2	R.FLEQQNQVLQTK.W	0.0012

effects of HSPA8 were again evident at 2 µg/ml ($P=0.0037$; $F_{1/84}=8.89$), as in the earlier experiment but to a lesser extent than both 0.25 and 0.5 µg/ml ($P=0.001$) for the comparison between TLP and 0.25 µg/ml and $P<0.0001$ for the specific comparison between TLP and both concentrations together). 8 µg/ml had no beneficial effect on maintaining sperm survival and it was concluded that the optimal concentration must be lower than 2 µg/ml (Fig. 3B).

A further series of bioassays was therefore undertaken using an even lower range of HSPA8 concentrations (0.1, 0.5 and 4 µg/ml); and sAPM (200 µg/ml) treatment was also included (Fig. 3C). Beneficial effects on sperm survival over a 24 h period were evident with HSPA8 at 0.1 µg/ml ($P=0.005$) and 0.5 µg/ml ($P=0.0001$). In this experiment, there was no beneficial effect observed with HSPA8 at 4 µg/ml. As 0.5 µg/ml HSPA8 had the greatest beneficial effect on sperm viability (Fig. 3A and C), it was concluded that this must be the optimal concentration of HSPA8.

In vitro fertilization assays

Spermatozoa were briefly (10 min) exposed to 0.5 µg/ml recombinant bovine HSPA8 protein before being diluted (1: 200) into the final fertilization droplet. A control treatment with no protein exposure was also set up. The summarised IVF results are shown in Table 2. Exposure to HSPA8 did not enhance the fertilization rate compared with the control, but it significantly enhanced the extent of monospermy (control versus HSPA8; 58.6 vs 88.9%; $P<0.05$). Exposing spermatozoa to HSPA8 also appeared to increase the rate of embryonic development so that by 60 h of incubation a markedly higher proportion of embryos had reached or passed the three or four cell stage (control versus HSPA8; 37.9 vs 74.1% and 3.5 and 22.2% respectively; $P<0.05$).

Effects of HSPA8 on bull spermatozoa incubated at 37 °C for 48 h

As HSPA8 had a beneficial effect on boar sperm viability, despite being a bovine recombinant protein, we hypothesized that its role in sperm viability maintenance would also apply to the bull. To test this hypothesis, a sperm viability experiment was set up using bull, instead of boar, spermatozoa.

When bull spermatozoa were incubated in the presence of HSPA8 for 48 h at 37 °C, ~ 50% remained viable. However, the inclusion of HSPA8 in the TLP medium increased the rate of sperm survival in a dose dependent manner up to an optimal value at 1 µg/ml (TLP control versus 1 µg/ml HSPA8; $P=0.0013$). Although, increasing the dose of HSPA8 further (to 4 µg/ml) still resulted in a beneficial effect on sperm survival ($P=0.006$) compared with TLP alone, the higher

Figure 3 (A) Viability index (mean $\pm$ S.E.M.) of boar spermatozoa incubated for 24 h at 39 °C in the presence of TALP only (Control), 200 µg/mlsAPM (sAPM), sAPM treated with anti-HSPA8 (anti-HSPA8+sAPM) and anti-HSPA8 only. (B) and (C). Viability index (mean $\pm$ S.E.M.) of boar spermatozoa incubated for 24 h at 39 °C presence of TLP, sAPM (200 µg/ml) or two concentration ranges ((B) and (C)) of bovine recombinant HSPA8. Significant beneficial effects of the protein treatments on sperm viability are indicated as follows: ** $P < 0.01$ *** $P < 0.0001$.

dose was less effective at maintaining sperm survival. When spermatozoa were incubated in media containing tubulin as a control protein treatment, there was no significant difference in outcome with respect to the TLP control. These results are summarised in Fig. 4.

Discussion

HSPA8 demonstrated consistent and significant enhancement of viability at low concentrations (0.5–4 µg/ml) for both porcine and bovine spermatozoa. As oviductal sperm storage is so widely observed in nature, it is likely that the fundamental physiological and biochemical mechanisms may also have been conserved across species. The present study is consistent with this view as it has revealed a role for the highly conserved protein HSPA8 in the sperm storage process; in fact, this molecule is sometimes regarded as the most highly conserved protein in nature (Daugaard *et al.* 2007).

In this study, we have demonstrated that the HSPA8 is present within a soluble protein subset prepared from the porcine oviductal epithelium, sAPM, and that recombinant bovine HSPA8 enhances the *in vitro* survival of both boar and bull spermatozoa. Thus, we conclude that the HSPA8 is likely to be a biologically active sAPM component responsible for improved *in vitro* sperm survival that we have observed and reported previously (Elliott *et al.* 2001). Our investigative strategy involved two complementary approaches; one was the direct proteomic identification of biotinylated sAPM proteins that became bound to the sperm surface. This approach was informative because the sAPM had been prepared from *in situ* biotinylated OEC; therefore the biotin-labelled proteins must have originated on or near the oviductal cell surface. The second approach aimed to use the proteomic information by incubating the sAPM with anti-HSPA8 in an effort to inactivate HSPA8, the selected protein of interest. The investigative strategy was then taken a step further by testing the biological activity of commercially available bovine recombinant HSPA8 in sperm survival bioassays. This protein was shown to be capable of reproducing the sperm-enhancing properties of sAPM, but at considerably lower concentrations.

As a prelude to these studies we also showed, by confocal microscopy, that when spermatozoa and fluorescent-labelled sAPM proteins are co-incubated, the sAPM proteins bind predominantly to the apical region of the sperm head. This together with evidence of boar sperm binding to sAPM coated beads supports the findings of our earlier work whereby a small number of surface labelled sAPM proteins were found to associate with spermatozoa during incubation (Satake *et al.* 2006). The majority of boar spermatozoa (63.1%) bound FITC-labelled sAPM proteins at the apical region of the sperm head; this is the region that associates with OEC *in vivo* (Rodriguez-Martinez 2007) and suggests that the sAPM-sperm interaction mimics the natural situation. This also correlates with binding patterns and interactions seen in other studies (Pollard *et al.* 1991, Pacey *et al.* 1995, Satake *et al.* 2006). Binding over the anterior acrosome also coincides with the distribution of sulphated glycolipids reported by Gadella *et al.* (1995); interestingly, the sulphated glycolipids are known to bind specifically to members of the HSP70 family, especially

Table 2 Effect of exposing spermatozoa to HSPA8 protein for 10 min prior to transfer into the IVF droplet.

Treatment	Number of oocytes	Degenerate oocytes	Failed to fertilize	Fertilized (%)	Monospermic (%)	2PN stage or later (% of fertilized)	3 or 4 cell embryos or later (% of fertilized)	7/8 cell embryos or later (% of fertilized)
Control	72	26	17	29 (40.3)	17 (58.6)	16 (55.1)	11 (37.9)	1 (3.5)
HSPA8	72	35	10	27 (37.5)	24 (88.9)*	27 (100)	20 (74.1)*	6 (22.2)*
Total	144	61	27	56	41	43	31	7

*Significantly higher than control values in the same column. $P < 0.05$.

HSPA8, through specific binding sites on the protein molecule (Boulanger *et al.* 1995, Lingwood *et al.* 2001, Mamelak *et al.* 2001). If this binding interaction is indeed functionally significant, it would explain the dose-dependent sperm–protein interaction observed here because the sperm surface-receptors would have to interact stoichiometrically with the exogenous protein. Since the anterior acrosomal sulphated glycolipid pattern reported by Gadella *et al.* (1995) undergoes a significant shift towards the equatorial segment during capacitation, we interpret our acrosomal patterns as indicating that most spermatozoa in our samples are uncapacitated; this would be expected as they had only been exposed to capacitation conditions for a very short period. This result would also agree with previous studies indicating that uncapacitated boar spermatozoa preferentially associate with OEC (Fazeli *et al.* 1999). That other spermatozoa either displayed extensive sAPM binding or no binding at all is therefore likely to be a reflection of sperm heterogeneity in the ejaculate; this is a well known property of boar ejaculates even though the specific reasons for the heterogeneity remain unclear. It is likely, however, that the differences correspond to inter-sperm differences in ability to colonise the oviduct.

Our attempt to mimic the sperm-oviduct interaction more precisely by incubating sAPM-coated beads with spermatozoa were of interest in that a surprisingly low number of spermatozoa became attached to each bead; although, this number was very low, it was nevertheless higher than the control beads that lacked surface-bound sAPM. There are a number of possible reasons for this, one plausible possibility being that the spermatozoa were only able to bind transiently to the bead surfaces, thus preventing the accumulation of spermatozoa. We did not, however, assess the effective density of sAPM bound to the bead surfaces, and a second possibility would be that the sAPM had been insufficiently coupled to the bead surface.

Further sperm/sAPM binding experiments enabled a subset of sperm-associating proteins to be identified using *in situ* biotinylated sAPM, followed by proteomic analysis of a significant 70 kDa sperm-binding protein band. Of the 19 proteins identified from the significant 70 kDa sperm-binding protein band, five were heat shock proteins and we chose to analyse one of these, namely

HSPA8, in more detail. A western blot of sAPM proteins probed with anti-HSPA8 confirmed its presence in sAPM.

sAPM was unable to enhance sperm viability if it had been pre-treated with anti-HSPA8. We interpreted this result as showing that HSPA8 could be a member of the sAPM protein milieu participating in the protection of sperm viability *in vitro*. Incubations (bioassays) of spermatozoa with serial dilutions of the commercial bovine recombinant HSPA8 confirmed this view by clearly demonstrating its effectiveness in enhancing sperm survival *in vitro* at concentrations that were ~ two orders of magnitude lower than the correspondingly bioactive sAPM concentration. It is of considerable interest that when the effectiveness of HSPA8 was tested with bull spermatozoa, it was again seen to enhance sperm survival at low concentrations and in a dose-dependent manner.

In this study, the HSPA8 protein was detected in sAPM via the direct proteomic analysis of the sperm-binding protein band and by western blotting with an antibody (anti-HSPA8). To our knowledge, this is the first report suggesting membrane bound or extracellularly located HSPA8 may have a role in the protection of spermatozoa in the oviduct, but the associated literature is consistent with the evidence we present here. HSPA8 is a constitutive member of the heat shock protein 70 family

Figure 4 Viability index (mean ± S.E.M.) of bull spermatozoa incubated for 48 h with various concentrations of bovine recombinant HSPA8. Significant beneficial effects of the protein treatments on sperm viability are indicated as follows: * $P < 0.05$ ** $P < 0.001$ *** $P < 0.0001$.

that includes a number of HSP70 variants. Some are stress-inducible and others are constitutive with involvement in housekeeping functions (for reviews, see Tavaría *et al.* 1996, Daugaard *et al.* 2007). HSPA8 has been reported to be involved in a multitude of the housekeeping chaperone functions including folding of nascent polypeptides, protein translocation across membranes, chaperone-mediated autophagy, prevention of protein aggregation under stress conditions and disassembly of clathrin coated vesicles (Daugaard *et al.* 2007). Thus, HSPA8 is considered as an essential housekeeping gene and it has been reported that HSPA8 knockout-mice cannot be created due to the essential role of HSPA8 in cell survival.

When we undertook a parallel series of experiments using FITC-labelled HSPA8, we were unable to demonstrate the occurrence of any clear binding patterns to spermatozoa (data not shown). Logically, we conducted these experiments using HSP8A concentrations that matched the biologically significant range ($\sim 1\text{--}4\text{ }\mu\text{g/ml}$). It could be argued that the concentration of FITC-labelled HSP8A protein was too low to be detectable on spermatozoa. However, it is also possible that HSP8A binds transiently to the functional sperm surface, exerting its effects instead by plasma membrane-mediated signalling mechanisms that probably operate via exceedingly low protein concentrations. This explanation would also be consistent with the large discrepancy between the concentration of sAPM and HSPA8 needed to elicit improved sperm survival.

The reduction of polyspermic fertilization and improved rate of embryonic development *in vitro* by briefly exposing spermatozoa to HSPA8 before interaction with oocytes is of considerable interest but difficult to explain. The results agree with a previous and larger IVF study, using the same conditions, in which spermatozoa were exposed to sAPM for 10 min before fertilization (Satake *et al.* 2007). In that study, the overall fertilization rate was significantly enhanced from 78 to 86% ($n=1235$ oocytes) when spermatozoa were pre-incubated with sAPM, and in the group of fertilized oocytes, polyspermic fertilization was also significantly reduced (47 vs 21%). Given that HSPA8 is a component of sAPM, it may also have been one of the agents responsible for producing these earlier results. However, the possibility that the HSPA8 protein acts in concert with proteins that are either already present in the sperm plasma membrane or within the oviductal epithelium *in vivo* cannot be ruled out.

Like the study reported here, experiments by Boilard and colleagues (Boilard *et al.* 2001) identified three chaperone proteins from cow oviducts that bound to bull spermatozoa; two of these were identified as Heat shock 60 kDa protein 1 (chaperonin; HSPD1 previously known as HSP60) and HSPA5 (Boilard *et al.* 2004). Interestingly, HSPA5 is also present in our sperm-binding 70 kDa sAPM band. Subsequent studies of these proteins have

also confirmed their presence in the human female reproductive tract (Lachance *et al.* 2005). These proteins were found to elevate intracellular calcium concentrations in spermatozoa; (Lachance *et al.* 2007), thus confirming that the heat shock proteins are able to act upon spermatozoa via plasma membrane mediated cell-signalling mechanisms. This provides supportive evidence that HSPA8 may act via a similar route. In their study, Lachance *et al.* (2007) were unable to detect any viability enhancing effects of HSPD1 and HSPA5 on human spermatozoa, although given that their duration of incubation was only 4 h, any effects may have been difficult to detect.

Heat shock proteins of similar size and function to those identified here as components of the oviductal epithelium have also been identified as components of spermatozoa themselves. They are known to be synthesized during spermatogenesis (Allen *et al.* 1988, Maekawa *et al.* 1989) and they adopt specific localizations within sperm e.g. within their plasma membrane (Boilard *et al.* 2004, Mitchell *et al.* 2007). While endogenous heat shock proteins clearly have roles in sperm function, data in the present study clearly show that exogenous heat shock proteins, such as HSPA8 can also have an important effect on sperm function.

Our data showing an important role for HSPA8 in the maintenance of boar and bull sperm survival *in vitro* implies that the mechanism of action could be highly conserved across species. Consequently, the protein could be a useful additive to sperm extenders used for artificial insemination for more than one species. Indeed, inter-species variability in effectiveness is a major difficulty for the development of sperm extenders in mammals, but we speculate that such a highly conserved protein as HSPA8 may be useful in overcoming this problem.

Materials and Methods

Materials and reagents

Reproductive organs from sows were obtained from local abattoirs within 20 min of slaughter, immediately chilled in ice-cold PBS (PBS) and transported to the laboratory. boar semen in beltsville thawing solution (BTS; Pursel & Johnson 1975) was supplied by JSR Genetics (Thorpe Willoughby, Yorkshire, UK). All reagents, except where stated, were obtained from Sigma-Aldrich; PBS pH 7.2 without calcium and magnesium (Invitrogen); electrophoresis equipment and reagents (Bio-Rad); western blotting reagents including Sulfo-NHS-LC-biotin (Pierce, Rockford, IL, USA); live-dead stains SYBR-14, PI and ethidium homodimer (ETHD-1; Molecular Probes, Eugene, OR, USA); methanol and glacial acetic acid (BDH, Poole Dorset); antibodies (Abcam plc, Cambridge, UK and Stressgen, Ann Arbor, MI, USA). Bovine recombinant HSPA8 was obtained from Stressgen (Cat no. SPP-751). Two modifications of Tyrode's medium (TLP and TALP) were used in this study; TLP consisted of 2 mM CaCl_2 , 3.1 mM KCl, 0.4 mM

MgCl₂·6H₂O, 100 mM NaCl, 25 mM NaHCO₃, 0.3 mM NaH₂PO₄·H₂O, 10 mM HEPES, 21.6 mM sodium lactate and 1 mM sodium pyruvate. Sometimes the TLP was supplemented with 12 mg/ml BSA and we therefore refer to it as TALP (TLP+Albumin).

Apical plasma membrane preparations

Unlabelled fractions

The OEC from the reproductive tissue of freshly slaughtered sows were harvested manually and the sAPM fractions were isolated and solubilized as described previously (Holt *et al.* 2005). Briefly, epithelial cells were scraped from the lumina of oviducts and sequentially and alternately treated with 10 mM MgCl₂ and differential centrifugation to release the APMs. Soluble proteins were extracted from the membranes with 1 M NaCl and high-speed centrifugation. The supernatant was separated from the pellet and concentrated using Microcon 3 000 Dalton cut-off centrifugal filter devices (Millipore UK Ltd, Watford, UK). The concentrated sAPM protein fraction was washed twice with TLP and stored at -80 °C. Protein concentration was measured using the Bradford (1976) method (kit supplied by Bio-Rad).

Biotin labelled sAPM fractions

Prior to epithelial cell collection and sAPM preparation the oviductal lumina were labelled with biotin ester Sulpho-NHS-LC-biotin (Pierce). This involved filling the oviducts with the prepared biotin ester (0.5 mg/ml) and incubating them for 30 min. The excess biotin was quenched by flushing with 50 mM NH₄Cl. sAPM proteins were prepared using the same method as that for the unlabelled fractions.

Sperm preparation

Viable boar sperm samples were separated from the storage diluent, usually BTS, by washing through a Percoll–saline concentration gradient (Holt & Harrison 2002). Semen aliquots (7 ml) were layered over 2 ml 35% Percoll on 2 ml 70% Percoll and centrifuged for 15 min at 200 *g* followed by 15 min at 900 *g*. The supernatant was aspirated to leave ~ 0.4 ml containing the sperm pellet.

Bull semen was supplied by Genus Freezing Unit, (Ruthin, Wales), in 1:5 dilution of Eqcellsire (IMV L'Aigle, France) and cooled to room temperature prior to overnight dispatch to our laboratory. A total of eight ejaculates were used in the current study. The samples were centrifuged for 10 min at 600 *g* and the supernatant removed. The sperm pellet was washed once and then diluted in TALP to give a sperm concentration of 100 × 10⁶/ml (final concentration 50 × 10⁶ sperm/ml).

Evaluation of sperm viability

Sperm viability (as defined by plasma membrane integrity) was assessed using SYBR-14 (200 nM) and ETHD-1 (2 µM) in PBS supplemented with 0.01% (v/v) formalin. Spermatozoa were incubated in this solution for 15 min at 39 °C in 5% CO₂ and at

least 200 cells were counted and classified as live (plasma membrane-intact; green) or dead (plasma membrane damaged; red). The data were expressed as mean viability index ± S.E.M. Viability index was defined as the ratio of % viable spermatozoa at time = 24 h/% viable spermatozoa at time = 0 h.

Evaluation of interactions between in situ biotinylated sAPM and spermatozoa

Samples of *in situ* biotinylated sAPM were adjusted to 3 mg/ml in 200 µl Immobilized Neutravidin coated agarose beads in PBS (Pierce). The samples were incubated for 1 h on a small shaker at 600 rpm (room temperature). The bead/sAPM samples were centrifuged to remove unbound protein and washed in 5 × 1 ml PBS for 2 min each at 2500 *g*. A sample of beads without added sAPM was also washed. The final pellet of beads was resuspended in 100 µl TLP. Boar semen was washed in a Percoll–saline concentration gradient as described above; 2 µl aliquants of washed spermatozoa were added to each of three tubes containing 25 µl each of 1) TLP only, 2) beads without sAPM and 3) beads with sAPM; these were incubated for 20 min at 39 °C in 5% CO₂. Clean electropositive slides (VWR, Lutterworth, UK) were prepared by warming to 37 °C. After 20 min an aliquot (10 µl) of the incubated sample was mounted on a slide beneath a coverslip. The coverslips were raised > 65 µm above the slide surface using small amounts of silicon grease mixed with 65 µm diameter glass beads. The sperm-bead interactions were recorded on video tape for observation; two observers made blind and independent estimates of the number of spermatozoa bound to beads.

Microscopic localization of sAPM proteins on the surface of spermatozoa

sAPM proteins were labelled with FITC using a ProtOn Fluorescein labelling kit (Vector Laboratories Inc., Burlingame, CA, USA) following the manufacturer's instructions. Briefly, 100 µl sAPM (7.0 mg/ml) was supplemented with 2 µl ProtOn FITC reagent and incubated for 30 min at room temperature. The reaction was stopped with 2 µl ethanolamine (1 M) and incubated for 5 min at room temperature. The sample was passed through the salt extraction column provided (by centrifugation at 500 *g* for 1 min) to remove excess fluorescein. The protein concentration of the sample was then assessed (Bradford 1976).

The labelled protein (2.4 mg/ml) was diluted in TLP (1:10 v/v). 2 µl Percoll–saline washed spermatozoa was added to 100 µl aliquots of this labelled protein solution. The samples were incubated at 39 °C for 10 min prior to the addition of 1 µl PI for a further 3 min. These samples were centrifuged for 3 min at 3000 *g*; the supernatant was removed and the samples resuspended in 80 µl TLP. Aliquots were examined by confocal microscopy (Zeiss LSM 510, Oberkochen Germany). At least ten images of spermatozoa from each boar were recorded. Sperm/sAPM localization was assessed subjectively as to whether there was 'no binding', 'binding to the acrosomal region' or 'binding to the head and mid-piece region'. Approximately 60 plasma membrane intact spermatozoa were evaluated from each of eight boar ejaculates.

In vitro fertilization technique

Sow ovaries were obtained from a local abattoir (A & G Barber, Purleigh, Essex, UK); these were placed in 0.9% saline solution supplemented with 100 IU/ml penicillin-G and 100 µg/ml streptomycin sulphate solution (pen/strep; Invitrogen Life Technologies) and maintained in a thermos flask at 25–30 °C for transportation to the laboratory. On arrival the ovaries were washed in lukewarm water then rinsed in warm saline solution with pen/strep supplements.

Follicular fluid from 3–6 mm diameter ovarian follicles were aspirated using an 18.5 gauge needle attached to a 10 ml disposable syringe containing 2 ml collection medium 1 (TCM 199 supplemented with 5 mM NaHCO₃, 15 mM Hepes (Na⁺ salt or free acid), 0.05 g/l kanamycin sulphate, 0.4% (w/v) BSA fraction V and 0.04 g/l heparin). The aspirated fluid was then filtered through a 70 µm cell strainer to harvest the oocytes. The oocytes were washed and collected into pre-warmed collection medium 2 (TCM 199 (with Hank's salts and sodium bicarbonate) supplemented with 0.1% (w/v) polyvinyl alcohol, 3.05 mM D-glucose, 0.91 mM sodium pyruvate and pen-strep solution). Cumulus–oocyte complexes (COCs) were collected under a stereomicroscope; oocytes had at least two complete layers of cumulus cells and evenly granulated ooplasm. The selected COCs were washed in collection medium 2 and transferred in groups of 50 COCs into pre-equilibrated 500 ml wells of IVM medium (TCM 199 (with Earle's salts, L-glutamine and sodium bicarbonate) and supplemented with 0.1% polyvinyl alcohol, 3.05 mM D-glucose, 0.91 mM sodium pyruvate, 0.57 µM cysteine and 10 ng/ml epidermal growth factor and pen-strep solution (Abeydeera *et al.* 2000) and pre-equilibrated). Before oocyte transfer 0.5 µg/ml LH and 0.5 µg/ml FSH were added to the 500 µl maturation droplets. The preparations were incubated for 48 h.

After IVM incubation 0.1% (w/v) hyaluronidase was added for 30 min to the maturation droplets to loosen the cumulus cells. COCs were then washed in pre-equilibrated mTBM (modified Tris-buffered medium (mTBM (Abeydeera & Day 1997): 113.1 mM NaCl, 3 mM KCl, 20 mM Tris-base, 11 mM D-glucose and 5 mM sodium pyruvate supplemented with 5 mM NaHCO₃ and 0.4% (w/v) BSA). After equilibration of the medium, 10 mM CaCl₂ was added to avoid precipitation). Groups of 30–40 COCs were placed in pre-equilibrated 98 µl droplets of mTBM covered in mineral oil. Approximately 3 × 10⁶ ml of Percoll-washed spermatozoa were incubated for 10 min in 1 ml pre-equilibrated mTBM supplemented with, or left without, 0.5 µg/ml bovine recombinant HSPA8. 2 µl sperm suspension were added to the IVF droplets, to give a final concentration of 6 × 10⁴ spermatozoa/ml. The oocytes and spermatozoa were co-incubated for 6 h.

Following sperm–oocyte co-incubation, the presumptive zygotes were collected into 1 ml pre-equilibrated NCSU23 medium (North Carolina State University-23 medium: 108.73 mM NaCl, 4.78 mM KCl, 1.19 mM K₂PO₄, 1.19 mM MgSO₄·7H₂O, 1 mM L-glutamine, 5.55 mM D-glucose, 7 mM taurine, 5 mM hypotaurine, 25.07 mM NaHCO₃ and 1.71 mM CaCl₂ supplemented with 0.4% (w/v) BSA and pen-strep solution) in a 5 ml glass tube and vortexed for

2 min to remove loosely bound spermatozoa and cumulus cells. The presumptive zygotes were then washed twice in fresh pre-equilibrated 40 µl droplets of NCSU23 and transferred to 20 µl of the same medium in groups of 20–25. They were cultured for 48 h and then fixed in 2.5% paraformaldehyde fixative with 0.5% triton X-100 for 30 min at 39 °C. The embryos were washed (× 3) in warm PBS and stained with 10 µg/ml PI for 10 min before being mounted in mounting medium (Vectashield, Vector Laboratories). They were examined by confocal microscopy (Zeiss LSM 510, Germany) to identify fertilization status. The oocytes/embryos were classified as 1) unfertilized oocytes (the presence of metaphase plates or female pronuclei), 2) fertilized, with male and female pronuclei, 3) one-cell embryos, 4) cleaved embryos with two or more cells 5) embryos with more than four cells.

Identification of the oviductal epithelial proteins that bound to spermatozoa

Biotinylated APM and sAPM (prepared *ex situ* and *in situ*) was incubated with spermatozoa (150 µg/ml and 1 × 10⁸ spermatozoa/ml) at 39 °C and 5% CO₂ in air for 30 min. After incubation the sperm suspensions were concentrated by microcentrifugation and then washed three or four times (> 1:100 dilution per wash) by microcentrifugation at 5000 g for 4 min for removal of APM and sAPM. To prepare a suspension of biotinylated proteins bound to sperm membranes, the washed pellets were resuspended in SDS sample buffer (50 mmol Tris/l pH 6.8 and 0.5% SDS w/v), incubated at room temperature for 90 min and then centrifuged at 14 000 g for 10 min. The supernatant (soluble fraction, comprising sperm membrane proteins and surface-bound biotinylated proteins) was aspirated for evaluation by one-dimensional PAGE and immunoblotting to detect biotinylated proteins.

A biotin-positive band in the immunoblot of sperm proteins corresponded to a 70 kDa band in the PAGE gel preparation. This band was therefore excised and analyzed further. To prepare sufficient protein for proteomic analysis sAPM proteins were separated using 16 single-dimensional PAGE lanes and the appropriate band in each was extracted for analysis. Specifically, this was achieved as follows. Protein (15 µg) was loaded into each of eight wells in two 10-well gels. The two outer wells of each gel were loaded with Kaleidoscope standards for identification of molecular weight. Electrophoresis was performed, using the discontinuous buffer system (Laemmli 1970) for 45 min at 190 volts after which the gels were fixed (40% methanol v/v and 7% glacial acetic acid) for 1 h and stained overnight in Brilliant Blue G-Colloidal Concentrate. The gels were destained in 10% acetic acid in 25% methanol (v/v) for 30 s and rinsed in 25% methanol before being placed carefully onto clean glass plates. Using a new sterile blade, the prominent band (Fig. 2A arrow) corresponding to the 70 kDa *in situ* sAPM biotinylated protein that bound to spermatozoa (Fig. 2B arrows) was excised from each of the 16 separate sAPM lanes. Each piece was cut into 1 mm² pieces and placed in 1 ml dH₂O in 1.8 ml centrifuge tubes for further proteomic processing.

Proteomic analysis

Gel pieces were washed in 25% (v/v) methanol, and then incubated for 1 h at 37 °C in Coomassie destain solution consisting of 40% (v/v) acetonitrile (ACN: VWR International Ltd., Leicester, UK) in 200 mM ammonium bicarbonate. The destain solution was removed, and gel pieces were incubated with ACN for 15 min at room temperature. ACN was subsequently removed, and the gel pieces were dried in a vacuum centrifuge. Dried gel pieces were stored at 4 °C until required. Proteins were digested with 20 ng/μl sequencing grade modified trypsin (Promega) in 50 mM ammonium bicarbonate at 37 °C for 12 h. The supernatant from the trypsin digest was transferred to a siliconized microcentrifuge tube. Peptides were sequentially extracted three times by incubation with peptide extraction solution consisting of 25 mM ammonium bicarbonate (10 min at room temperature), 5% formic acid (15 min at 37 °C), and ACN (15 min at 37 °C). Each extraction was followed by centrifugation and removal of supernatants. The original supernatant and the supernatants from the three sequential extractions were combined and dried in a vacuum centrifuge for 4–6 h. The dried peptides were dissolved in 7 μl 0.1% (v/v) formic acid in 3% (v/v) ACN in water. Samples were centrifuged for 5 min at 12 000 *g*, and the supernatants were subjected to liquid chromatography electrospray ionization tandem mass spectrometry (LC-ESI-MS/MS).

Liquid chromatographic separations of the tryptic digests were performed using a reverse phase CapLC system (Waters, Manchester, UK). Peptides were desalted by a PepMap C₁₈ microguard column (300 μm internal diameter × 1 mm; LC-Dionex, Leeds, UK) and were then transferred to the analytical column (PepMap C₁₈; 75 μm internal diameter × 15-cm column (LC-Dionex). The peptides were eluted in a 60 min gradient. The compositions of the hydrophilic and hydrophobic solvents were 5% ACN, 0.1% formic acid and 95% ACN, 0.1% formic acid. The column eluent was sprayed directly into the nano-ESI source of a Q-TOF microcolumn (Waters). An initial MS scan was performed, and selection of ions for CID was automated by Mass Lynx software (Waters). CID selection criteria were set for 2+ and 3+ ions within the range of 400–2000 *m/z* above 10 ion counts.

Spectra were searched against the mass spectrometry data base in a sequence query search using MASCOT 2.2.03 software (www.matrixscience.com). The taxonomy was limited to filter for only mammalian matches, and trypsin was used as the cleavage enzyme with one missed cleavage site allowed. The peptide tolerance was set to 1.2 kDa, and the MS/MS tolerance was set to 0.6 kDa. Carbamidomethyl modifications of cysteine and oxidized methionine were set as variable modifications. Matches were considered valid if MS/MS data for multiple unique peptides per protein were identified, each with ion scores above the threshold of statistical significance (values generated by MASCOT). Manual examination of MS/MS data was performed for single peptide matches. In these cases, a continuous stretch of peptide sequence covered by either the y- or b-ion series was required for the protein match to be considered valid.

Western blot analysis of sAPM

The proteomic analysis indicated that spermatozoa avidly bound a series of heat shock proteins. Of these, we chose to examine HSPA8 in more detail. First, for independent confirmation that HSPA8 was indeed present in sAPM a western blot were performed. sAPM proteins were separated by SDS-PAGE (10% gel), blotted on to nitrocellulose and incubated in blocking buffer for 1 h. Anti-HSPA8 (SPA-815), at the manufacturer's recommended dilution was added to the blot and incubated for 1 h at room temperature. The blot was washed 6 × 10 min in tris-buffered saline (Pierce) supplemented with tween-20 (TTBS) followed by the addition of the goat anti-rat secondary antibody conjugated with alkaline phosphatase (AP; ab7098, Abcam), at the manufacturer's recommended dilution for a further hour. The blot was washed 6 × 10 min in TTBS and, finally, incubated for 3 min in Lumi-phos for AP (Pierce) detection. The blot was exposed to X-ray film for periods between 5 and 60 s.

Sperm viability assays

Boar sperm viability in the presence of antibody-treated sAPM

Antibody-treated sAPM (200 μg/ml) was prepared by incubating sAPM with 1 μg/ml anti-HSPA8 (Ab1427) in TLP for 2 h on a rotary shaker at room temperature. Then, semen from eight boars was individually washed with Percoll-saline as previously described and each sample (in TALP) was adjusted to give a final concentration of 25×10^6 spermatozoa/ml. 15 μl droplets of sperm suspension and antibody-treated sAPM (with untreated sAPM and TLP as control treatments) were mixed 1:1 in 20 mm diameter wells in four well Multi-dish plates (Sigma Aldrich) and then covered with mineral oil. They were incubated at 39 °C in 5% CO₂ for 24 h. The viability of spermatozoa was assessed at zero hours and after 24 h.

Dose response effect of HSPA8 protein on boar sperm viability

Boar spermatozoa were prepared for the bioassay as previously described. Semen samples from seven boars were used for series 1 and eight boars for series 2. Recombinant bovine HSPA8 (Stressgen; SPP-751) was diluted in TLP to give final concentrations of 16, 8, 4 and 2 μg/ml (series 1) and 8, 2, 0.5 and 0.25 μg/ml (series 2) when incubated with equal volumes of spermatozoa. Control samples were incubated in TLP only or TLP + sAPM (200 μg/ml). The samples were incubated at 39 °C in 5% CO₂ for 24 h. Sperm viability was examined as described earlier.

Effect of HSPA8 protein on bull sperm viability

Serial dilutions of bovine recombinant HSPA8 and a control protein α tubulin were prepared in TALP to give final concentrations of 4, 1, 0.5 and 0.1 μg/ml. Spermatozoa were added to droplets of TALP, TALP + α tubulin (Abcam plc) or TALP + recombinant HSPA8 in equal volumes of 25 μl each. The 50 μl droplets were covered with mineral oil and incubated at 37 °C in a 5% CO₂ incubator for 48 h. The viability of samples was assessed at zero hours and then at 48 h following the same procedure as that for boar spermatozoa.

Statistical analyses

Percentage and Viability Index data were log transformed and treatment effects were examined using analyses of variance (Statistica for Windows V7; Statsoft UK, Letchworth, UK). Individual contrasts between treatments were examined using orthogonal polynomial coefficients. Treatment effects in the IVF experiment were analyzed using Fisher's exact χ^2 -tests. Sperm – bead binding effects were assessed using the non-parametric Mann–Whitney *U* test.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This study was funded by grants from the Biotechnology and Biological Sciences Research Council and the London Development Agency's Heptagon Fund.

Acknowledgements

We are grateful to Dr Elena Badia for her assistance with elucidating the proteomics data and to Dr Abdul Al-Haider for assistance with the IVF experiments. We also thank JSR Genetics (Thorpe Willoughby, UK) for supplying the boar semen used in these experiments.

References

- Abeydeera LR & Day BN 1997 Fertilization and subsequent development *in vitro* of pig oocytes inseminated in a modified tris-buffered medium with frozen-thawed ejaculated spermatozoa. *Biology of Reproduction* **57** 729–734.
- Abeydeera LR, Wang WH, Cantley TC, Rieke A, Murphy CN, Prather RS & Day BN 2000 Development and viability of pig oocytes matured in a protein-free medium containing epidermal growth factor. *Theriogenology* **54** 787–797.
- Allen RL, O'Brien DA, Jones CC, Rockett DL & Eddy EM 1988 Expression of heat shock proteins by isolated mouse spermatogenic cells. *Molecular and Cellular Biology* **8** 3260–3266.
- Boilard M, Reyes-Moreno C & Sirard MA 2001 Binding of chaperonins to bovine spermatozoa by direct contact to apical plasma membrane of oviduct epithelial cells. *Biology of Reproduction* **64** 112.
- Boilard M, Reyes-Moreno C, Lachance C, Massicotte L, Bailey JL, Sirard MA & Leclerc P 2004 Localization of the chaperone proteins GRP78 and HSP60 on the luminal surface of bovine oviduct epithelial cells and their association with spermatozoa. *Biology of Reproduction* **71** 1879–1889.
- Boulanger J, Faulds D, Eddy EM & Lingwood CA 1995 Members of the 70 kDa heat shock protein family specifically recognize sulfoglycolipids: role in gamete recognition and mycoplasma-related infertility. *Journal of Cellular Physiology* **165** 7–17.
- Bradford MM 1976 A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72** 248–254.
- Daugaard M, Rohde M & Jaattela M 2007 The heat shock protein 70 family: highly homologous proteins with overlapping and distinct functions. *FEBS Letters* **581** 3702–3710.
- Elliott RMA, Duncan A, Watson PF, Holt WV & Fazeli AR 2001 Peripheral bound membrane proteins are involved in the maintenance of boar sperm viability by oviductal apical plasma membrane preparations *in vitro*. *Molecular Biology of the Cell* **12** 117a.
- Fazeli A, Duncan AE, Watson PF & Holt WV 1999 Sperm–oviduct interaction: induction of capacitation and preferential binding of uncapacitated spermatozoa to oviductal epithelial cells in porcine species. *Biology of Reproduction* **60** 879–886.
- Fazeli A, Elliott RM, Duncan AE, Moore A, Watson PF & Holt WV 2003 *In vitro* maintenance of boar sperm viability by a soluble fraction obtained from oviductal apical plasma membrane preparations. *Reproduction* **125** 509–517.
- Gadella BM, Lopes-Cardozo M, Vangolde LMG, Colenbrander B & Gadella TWJ 1995 Glycolipid migration from the apical to the equatorial subdomains of the sperm head plasma-membrane precedes the acrosome reaction – evidence for a primary capacitation event in boar spermatozoa. *Journal of Cell Science* **108** 935–946.
- Galbraith DA 1993 Multiple paternity and sperm storage in turtles. *Herpetological Journal* **3** 117–123.
- Georgiou AS, Sostaric E, Wong CH, Snijders AP, Wright PC, Moore HD & Fazeli A 2005 Gametes alter the oviductal secretory proteome. *Molecular and Cellular Proteomics* **4** 1785–1796.
- Gist DH & Congdon JD 1998 Oviductal sperm storage as a reproductive tactic of turtles. *Journal of Experimental Zoology* **282** 526–534.
- Holt WV & Harrison RA 2002 Bicarbonate stimulation of boar sperm motility via a protein kinase A-dependent pathway: between-cell and between-ejaculate differences are not due to deficiencies in protein kinase A activation. *Journal of Andrology* **23** 557–565.
- Holt WV, Elliott RMA, Fazeli A, Satake N & Watson PF 2005 Validation of an experimental strategy for studying surface-exposed proteins involved in porcine sperm–oviduct contact interactions. *Reproduction, Fertility, and Development* **17** 683–692.
- Holt WV, Elliott RM, Fazeli A, Sostaric E, Georgiou AS, Satake N, Prathalingam N & Watson PF 2006 Harnessing the biology of the oviduct for the benefit of artificial insemination. *Society of Reproduction and Fertility* **62** 247–259.
- Hunter RHF 1981 Sperm transport and reservoirs in the pig oviduct in relation to the time of ovulation. *Journal of Reproduction and Fertility* **63** 109–117.
- Hunter RH & Rodriguez-Martinez H 2004 Capacitation of mammalian spermatozoa *in vivo*, with a specific focus on events in the Fallopian tubes. *Molecular Reproduction and Development* **67** 243–250.
- Lachance C, Boilard M & Leclerc P 2005 Effect of HSP60 and GRP78 on human sperm functions. *Journal of Andrology* (Supplement) S39.
- Lachance C, Bailey JL & Leclerc P 2007 Expression of Hsp60 and Grp78 in the human endometrium and oviduct, and their effect on sperm functions. *Human Reproduction* **22** 2606–2614.
- Laemmli UK 1970 Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227** 680–685.
- Lingwood CA, Mylvaganam M, Whetstone H, Mamelak D & Kiso M 2001 Specific binding and modulation of hsp70 by sulfogalactolipids. *FASEB Journal* **15** A866.
- Maekawa M, O'Brien DA, Allen RL & Eddy EM 1989 Heat-shock cognate protein (hsc71) and related proteins in mouse spermatogenic cells. *Biology of Reproduction* **40** 843–852.
- Mamelak D, Mylvaganam M, Tanahashi E, Ito H, Ishida H, Kiso M & Lingwood C 2001 The aglycone of sulfogalactolipids can alter the sulfate ester substitution position required for hsc70 recognition. *Carbohydrate Research* **335** 91–100.
- Mitchell LA, Nixon B & Aitken RJ 2007 Analysis of chaperone proteins associated with human spermatozoa during capacitation. *Molecular Human Reproduction* **13** 605–613.
- Pacey AA, Hill CJ, Scudamore IW, Warren MA, Barratt CLR & Cooke ID 1995 The interaction *in vitro* of human spermatozoa with epithelial cells from the human uterine (Fallopian) tube. *Human Reproduction* **10** 360–366.
- Pollard JW, Plante C, King WA, Hansen PJ, Betteridge KJ & Suarez SS 1991 Fertilizing-capacity of bovine sperm may be maintained by binding to oviductal epithelial-cells. *Biology of Reproduction* **44** 102–107.
- Pratt HL & Tanaka S 1994 Sperm storage in male elasmobranchs; a description and a survey. *Journal of Morphology* **219** 297–308.
- Pursel VG & Johnson LA 1975 Freezing of boar spermatozoa: fertilizing capacity with concentrated semen and a new thawing procedure. *Journal of Animal Science* **40** 99–102.

- Racey PA** 1979 The prolonged storage and survival of spermatozoa in Chiroptera. *Journal of Reproduction and Fertility* **56** 391–402.
- Racey PA & Potts DM** 1970 Relationship between stored spermatozoa and the uterine epithelium in the pipistrelle bat (*Pipistrellus pipistrellus*). *Journal of Reproduction and Fertility* **22** 57–63.
- Racey PA, Uchida TA, Mori T, Avery MI & Fenton MB** 1987 Sperm–epithelium relationships in relation to the time of insemination in little brown bats (*Myotis lucifugus*). *Journal of Reproduction and Fertility* **80** 445–454.
- Rodriguez-Martinez H** 2007 Role of the oviduct in sperm capacitation. *Theriogenology* **68** S138–S146.
- Rodriguez-Martinez H, Saravia F, Wallgren M, Tienthai P, Johannisson A, Vazquez JM, Martinez E, Roca J, Sanz L & Calvete JJ** 2005 Boar spermatozoa in the oviduct. *Theriogenology* **63** 514–635.
- Satake N, Elliott RM, Watson PF & Holt WV** 2006 Sperm selection and competition in pigs may be mediated by the differential motility activation and suppression of sperm subpopulations within the oviduct. *Journal of Experimental Biology* **209** 1560–1572.
- Satake N, Alhaider AK, Holt WV & Watson PF** 2007 Exposure of spermatozoa to solubilized extracts of the oviductal epithelium apical plasma membrane enhances fertilization in porcine *in vitro* fertilization. *Reproduction, Fertility, and Development* **19** 272.
- Sever DM & Brizzi R** 1998 Comparative biology of sperm storage in female salamanders. *Journal of Experimental Zoology* **282** 460–476.
- Smith TT & Nothnick WB** 1997 Role of direct contact between spermatozoa and oviductal epithelial cells in maintaining rabbit sperm viability. *Biology of Reproduction* **56** 83–89.
- Tavaria M, Gabriele T, Kola I & Anderson RL** 1996 A hitchhiker's guide to the human Hsp70 family. *Cell Stress and Chaperones* **1** 23–28.

Received 9 July 2008

First decision 20 August 2008

Revised manuscript received 27 September 2008

Accepted 7 November 2008