

Theca: the forgotten cell of the ovarian follicle

J M Young and A S McNeilly

Human Reproductive Sciences Unit, Medical Research Council, Edinburgh EH16 4TJ, UK

Correspondence should be addressed to J M Young; Email: j.young@hrsu.mrc.ac.uk

Abstract

Theca cells function in a diverse range of necessary roles during folliculogenesis; to synthesize androgens, provide crosstalk with granulosa cells and oocytes during development, and provide structural support of the growing follicle as it progresses through the developmental stages to produce a mature and fertilizable oocyte. Thecal cells are thought to be recruited from surrounding stromal tissue by factors secreted from an activated primary follicle. The precise origin and identity of these recruiting factors are currently not clear, but it appears that thecal recruitment and/or differentiation involves not just one signal, but a complex and tightly controlled combination of multiple factors. It is clear that thecal cells are fundamental for follicular growth, providing all the androgens required by the developing follicle(s) for conversion into estrogens by the granulosa cells. Their function is enabled through the establishment of a vascular system providing communication with the pituitary axis throughout the reproductive cycle, and delivering essential nutrients to these highly active cells. During development, the majority of follicles undergo atresia, and the theca cells are often the final follicular cell type to die. For those follicles that do ovulate, the theca cells then undergo hormone-dependent differentiation into luteinized thecal cells of the corpus luteum. While the theca is an essential component of follicle development and ovulation, we do not yet fully understand the control of recruitment and function of theca cells, an important consideration since their function appears to be altered in certain causes of infertility.

Reproduction (2010) **140** 489–504

What are theca cells?

Reproduction is the result of a coordinated signaling network between the gonads, pituitary, and hypothalamus. The ovary is responsible for nurturing growing oocytes until an estradiol signal from primed preovulatory follicles induces GnRH, and a consequent luteinizing hormone (LH) surge, to release a mature oocyte, which is then capable of being fertilized to produce an embryo. Within the ovary, there are a number of cell types that support the growth and development of oocytes until ovulation. The oocyte is surrounded by a layer of granulosa cells, which change morphology and proliferate when an oocyte begins the process of folliculogenesis. Activated follicles are thought to recruit precursor thecal cells from the stromal layer surrounding the granulosa cells and oocyte. Together, these form the follicle structure that synthesizes steroid hormones (Hillier *et al.* 1994). Thecal cells are not capable of producing estrogen but do produce androgens in response to LH, which are then converted into estrogen by follicle stimulating hormone (FSH)-induced aromatase in the neighboring granulosa cells of selected growing follicles. Over the past decade, research has focused on granulosa cells and their interaction with the oocyte, and the theca has been somewhat forgotten as a necessary and vital part of the developmental process.

The classification system for folliculogenesis has been well defined in numerous publications (McNatty *et al.* 1999, 2007, Montgomery *et al.* 2001, Barnett *et al.* 2006, Edson *et al.* 2009). Briefly (see Fig. 1), primordial follicles (type 1) are in the resting stage before being activated to start development and the oocyte is surrounded by one layer of flattened granulosa cells; type 1a are the follicles transitioning through to the primary (type 2) stage when the granulosa cells become cuboidal. Primary follicles have one layer of cuboidal granulosa cells, secondary follicles (type 3) have two to four layers of granulosa cells, large preantral (type 4) follicles have four to six layers of granulosa cells, and antral follicles (type 5) have more than five layers of granulosa cells. It is after secondary follicle formation that the thecal cells begin to emerge and form a layer around the granulosa–oocyte structure. Throughout folliculogenesis, the rates of atresia increase, and the early stages of folliculogenesis proceed very slowly (Gougeon 1986, Hirshfield 1994); therefore, most follicles are observed at early stages of development. At the antral stage, follicles become gonadotropin dependent and form large antral follicles (type 5+), most of which undergo atresia, and few are selected for ovulation (reviewed by Scaramuzzi *et al.* (1993) and Edson *et al.* (2009)).




Figure 1 Thecal cell development and function during folliculogenesis. Thecal cells are vital for successful folliculogenesis. A primordial follicle consists of an oocyte and surrounding granulosa cells (GCs), and thecal layers are not formed until the follicle is activated and reaches the secondary stage of development. Thecal cells are required for the production of androgens to provide a structural scaffold, and they form the network of cells that support the vascular system, and after ovulation, thecal cells luteinize and form cells of the corpus luteum.

Origin of theca cells

Theca cells are first observed once a follicle has two or more layers of granulosa cells, which is around the time when thecal cells become LH responsive and steroidogenic enzymes are activated (Magoffin & Weitsman 1994). These specialized cells have long been thought to originate from fibroblast-like precursor cells within the ovarian stroma (Erickson *et al.* 1985, Orisaka *et al.* 2006, Honda *et al.* 2007). The putative undifferentiated progenitor theca cells do not express LH receptors (LHRs) or steroidogenic enzymes and are therefore not LH responsive, showing that initiation of theca cell differentiation is gonadotropin independent (Magoffin & Weitsman 1994). As theca cells are only associated with growing follicles, one would assume that the follicle itself produces factors that signal to the stroma to recruit cells that form the theca. Very few studies have investigated the factors that recruit thecal cells to the activated primary and secondary follicles. It is not currently clear whether cells surrounding the activated

follicle differentiate into the theca layer, or whether they are in fact recruited from the stroma to form the theca layers. However, results from mature thecal cells cultured *in vitro* do provide clues about the origin of some recruiting factors and, more importantly, the complexity of this system and provide evidence of steroidogenic regulators with potential differentiative roles. Selected factors are discussed later in the review.

Structure of theca cells

Electron microscopic analysis of normal thecal development in ovine follicles throughout folliculogenesis showed that thecal layers from small follicles (<3 mm diameter) were composed of flattened theca cells together with capillaries and bundles of collagen lying next to the basal lamina (O'Shea *et al.* 1978). The thecal cells were either fibroblast-like cells or presumed steroidogenic cells with large amounts of smooth endoplasmic reticulum. As the follicles grew, the thecal

cells hypertrophied, became less flattened and richer in endoplasmic reticulum, eventually producing pseudopodia, and contained many droplets of lipid but showed no signs of degradation.

Theca cells are highly differentiated with structural features characteristic of steroid-secreting cells including abundant mitochondria with vesicular cristae, agranular endoplasmic reticulum, and lipid vesicles (reviewed by Magoffin (2005)). The mitochondria contain the first enzyme in the steroidogenic pathway, cholesterol side-chain cleavage cytochrome P450 (CYP11A), and the endoplasmic reticulum contains the remaining enzymes necessary to produce androgens. The lipid vesicles store the precursors for steroid hormone synthesis as cholesterol esters which are transported into the mitochondria by steroidogenic acute regulatory protein (STAR; reviewed by Manna *et al.* (2009)). Thecal cells are vital components of the follicle, providing both structural support and being the exclusive producer of ovarian androgens, which are necessary as substrates for estradiol production in the neighboring granulosa cells.

Steroidogenesis

Androgens synthesized in thecal cells are transported to the granulosa cells where P450 aromatase converts these androgens to estrone and 17 β -estradiol. The steroidogenic enzymes are produced in a cell-specific manner (Wood & Strauss 2002), and in addition to the large amount of androgen receptor expressed in thecal and granulosa cells (but not in oocytes; Li *et al.* 2009a), studies indicate that a complete oocyte-independent system for androgen production exists within the growing follicle. Indeed, androgen and subsequent estradiol production can occur in mouse ovaries devoid of oocytes (McNeilly *et al.* 2000). Silencing 17 α -hydroxylase (CYP17) in the rat ovary caused a decline in androstenedione, 17 α -hydroxyprogesterone, and testosterone production, and also reduced progesterone levels (Li *et al.* 2009b); showing ovarian androgen biosynthesis can be inhibited by silencing CYP17 expression alone, and indicating a potential target for therapeutic development.

Androgen production from thecal cells in the gonadotropin-dependent stage is largely under the control of LH from the pituitary (Baird *et al.* 1981, Palermo 2007). LH is released in a pulsatile manner, and the frequency and amplitude of these pulses vary across the reproductive cycle in response to ovarian steroidogenic feedback. The pulse frequency of LH dictates the amounts of steroid hormones produced, where each pulse of LH is followed by an increase in androstenedione and estradiol secreted from the ovary in many species (Baird *et al.* 1976, 1981, Peluso *et al.* 1984, Schallenberger *et al.* 1984, Walters & Schallenberger 1984). When theca cells were cultured *in vitro*, low levels of LH also stimulated androgen production (Campbell *et al.* 1998, Ryan *et al.* 2008),

whereas, at high doses, LH inhibited androstenedione production and stimulated progesterone secretion as well as changing cell morphology indicating that the high LH levels induced luteinization in these cells (Campbell *et al.* 1998). LH has been shown to increase levels of STAR and steroidogenic enzymes (CYP11A1, CYP17, and 3- β -hydroxysteroid dehydrogenase (HSD3B)) and *LHR* gene expression (Magoffin & Weitsman 1993a, 1993b, 1993c, 1994, Lavoie & King 2009; see Fig. 2).

Insulin also plays an important role in thecal cell function. *In vitro* studies using thecal cells from porcine, bovine, and ovine ovaries have shown that insulin induced dose-dependent cell proliferation, increased steroid production, and increased the expression of genes encoding STAR, CYP11A1, and CYP17, thus promoting steroidogenesis (Morley *et al.* 1989, Campbell *et al.* 1995, 1998, Wrathall & Knight 1995, Mamluk *et al.* 1999, Smith *et al.* 2005; see Fig. 2).

The onset of thecal steroidogenic enzyme gene expression is similar in those mammalian species studied in depth (Pollack *et al.* 1997, Kerban *et al.* 1999, Lundy *et al.* 1999, Watson *et al.* 2000, Logan *et al.* 2002). Theca cells are first able to produce steroids just prior to antrum formation, as shown by the onset of expression of STAR, CYP11A1, CYP17, HSD3B, and LHR in thecal cells of preantral (large type 4) follicles, and the mRNA and protein localization at specific stages mirrored one another (Logan *et al.* 2002) and does not require gonadotropins (Scaramuzzi *et al.* 1993). mRNA encoding steroidogenic enzymes were also observed in the theca of bovine preantral follicles, although, unlike sheep, STAR expression was limited to thecal cells (Bao & Garverick 1998). Steroidogenic factor 1, a well-studied transcription factor regulating P450 enzymes and STAR, was expressed by granulosa cells, and protein was observed in both thecal and granulosa cells (Logan *et al.* 2002). Overall, the expression patterns found in the sheep ovary are similar to those observed in other mammalian species.

Angiogenesis

Small primordial follicles are located in the avascular region of the ovarian cortex, especially in larger species, and do not have their own vascular system. Once follicle growth is activated and a thecal layer has been recruited, a follicle develops its own vascular network within the surrounding thecal layer (see Fig. 1). Thecal cell proliferation begins early in the secondary stage of follicle development, although endothelial cell staining is still absent at this point showing that the theca forms before vascularization begins (Fraser & Duncan 2009). There are many potential factors involved in controlling angiogenesis in the developing follicle but vascular endothelial growth factor (VEGF) has a central role and has been studied extensively. VEGF, a potent mitogen for endothelial cells (Ferrara & Davis-Smyth 1997), stimulates vascular permeability (Connolly 1991,




Figure 2 Modulation of steroidogenic enzymes. Expression levels of genes encoding steroidogenic enzymes are regulated by many different factors within the ovary. The process of androgen production requires the following enzymes: cholesterol side-chain cleavage cytochrome P450 (CYP11A1), 17 α -hydroxylase (CYP17) and 3- β -hydroxysteroid dehydrogenase (HSD3B), as well as steroidogenic acute regulatory protein (STAR). Theca cells from sheep, humans, and primates principally produce androstenedione, whereas rodents produce testosterone as precursors for estradiol production in the neighboring granulosa cells. The modulating factors are depicted here, where in green circles the stimulatory factors are shown, and the inhibitory factors are listed in red boxes. BMP, bone morphogenetic protein; GDF9, growth differentiation factor 9; HGF, hepatocyte growth factor; hCG, human chorionic gonadotropin; IGF, insulin-like growth factor; SCF, stem cell factor/kit ligand; TGFB, transforming growth factor β .

Senger *et al.* 1993), and is highly expressed in granulosa cells and at lower levels in the thecal layer of follicles from the secondary stage onwards in primate ovaries (Taylor *et al.* 2004). In rodents, cows, and pigs, VEGF expression is weak in early follicle development but increases as the follicle progresses to ovulation (Maisonpierre *et al.* 1997, Barboni *et al.* 2000, Greenaway *et al.* 2005), although levels of VEGF decrease in granulosa cells in sheep and marmoset follicles just prior to ovulation (Ravindranath *et al.* 1992, Redmer *et al.* 2001, Taylor *et al.* 2004). In primates, mRNA encoding VEGF has been observed at the secondary stage in both the theca and granulosa cells (Taylor *et al.* 2004). VEGF was upregulated in rat ovaries during the primordial to primary transition thus preceding vascularization (Kezele *et al.* 2005). Certainly at early stages of follicle development, inhibition of VEGF prevents endothelial cell proliferation, and decreases thecal cell proliferation therefore hindering follicle development (Wulff *et al.* 2002, Fraser *et al.* 2005, Fraser & Duncan 2009). Suppression of gonadotropins using a GnRH antagonist results in reduced thecal and endothelial cell proliferation, and lower vascular density in antral follicles (Taylor *et al.* 2004), an effect probably due to reduced production of VEGF (Fraser & Duncan 2009).

While VEGF is critically involved in regulating follicle development in many species (Barboni *et al.* 2000, Mattioli *et al.* 2001, Wulff *et al.* 2001a, 2001b, 2002, Hunter *et al.* 2004, Martelli *et al.* 2008), many other

factors also contribute and modulate angiogenesis and vasculogenesis in mammals, such as transforming growth factor β (TGFB) superfamily members and their antagonists, angiopoietins, fibroblast growth factor (FGF), and gonadotropins, but whether they directly affect theca cell function remains to be explored.

Life span of theca cells

Folliculogenesis is a process that spans many weeks where the majority of follicles undergo atresia and only a few become dominant and go on to ovulate successfully (reviewed by Scaramuzzi *et al.* (1993)). During atresia, cell death is not confined to a specific cell type, but the entire follicle is degraded during this process. There are various ways that ovarian cells have been reported to die including apoptosis, autophagy, necrosis, and cornification (Jolly *et al.* 1994, Van Wezel *et al.* 1999, D'Haeseleer *et al.* 2006). In bovine follicles, oocytes of preantral follicles are the first component to die, whereas in later stages of development, the granulosa cells die first, although in particular types of atresia, thecal cells also die very early (Rodgers & Irving-Rodgers 2009; see Fig. 3). In the cow, atretic antral follicles have been classified into two types; antral atresia and basal atresia. Basal atresia occurred only in small antral follicles (<5 mm in bovine ovaries) where the theca cell layer becomes disrupted, having high levels of collagen, early death of endothelial and thecal cells, reduced




Figure 3 Theca cell fate. Atresia during follicle development is a more common fate of thecal cells than progression through to luteinization. In preantral follicles, the oocytes die first, and as follicles progress through development, follicles undergo basal or antral atresia. Thecal cells appear to be more susceptible to cell death early in follicular development, whereas in basally atretic follicles, the thecal layers are disrupted and are less vascular. Throughout all stages of development, antral granulosa cells are commonly the first cells to become atretic; hence, thecal cells are often the last to die during follicle atresia and the entire follicle is degraded during this process. FF, follicular fluid.

insulin-like factor 3 expression, and reduced androgen production associated with higher levels of progesterone in the follicular fluid in these follicles (Irving-Rodgers *et al.* 2003, Clark *et al.* 2004). These observations suggest that the theca is more susceptible to cell death early in follicular development compared to at later stages (Irving-Rodgers *et al.* 2001). In contrast, in atretic antral follicles where pyknotic nuclei were first observed in antral granulosa cells, thecal cells were the last to be disrupted (see Fig. 3). Thus, thecal cells from follicles undergoing atresia appear to respond differently depending on the stage of follicle development.

Since the blood supply is vital for follicle survival and transport of endocrine factors, vascularization is important in determining the fate of the follicle, and maintaining the blood supply is necessary for follicular health. The effect of atresia in the vasculature depends on the stage of follicle development when atresia is occurring. Small atretic follicles have reduced the numbers of capillaries in the thecal layers; the endothelium begins to degrade, and thecal capillaries become blocked by degrading material (O'Shea *et al.* 1977). Extensive hypertrophy of theca cells is observed during the early stages of atresia in human, rat, and rabbit, but not in sheep or bovine follicles (Himelstein-Braw *et al.* 1976, O'Shea *et al.* 1978, Erickson *et al.* 1985, Clark *et al.* 2004), which, together with the loss of granulosa cell function as they undergo

apoptosis, results in androgens being the predominant steroids secreted by atretic antral follicles (Moor *et al.* 1978). In larger bovine antral follicles, the vasculature is well established in non-atretic follicles (Shimizu *et al.* 2003), while the thecal layers of atretic follicles show signs of apoptosis from the outer layers progressing internally, and throughout the vascular network (Jiang *et al.* 2003). Whether the changes in the vasculature observed during follicle atresia are the cause or the effect of the atretic process itself is yet to be established.

Follicle fate is regulated by apoptotic factors such as nodal, which is produced by the thecal cells and acts to promote apoptosis in the neighboring granulosa cells (Wang *et al.* 2006, Craig *et al.* 2007). The oncogene *Skil* (also known as *SnoN*), involved in regulating TGF β superfamily signal transduction, has been mapped in the mouse ovary recently and is expressed in the theca throughout development and during atresia (Xu *et al.* 2009). This factor can modulate differentiation, proliferation, and apoptosis of several cell types (reviewed by Deheuninck & Luo (2009)), and was shown to have a specific manner of expression relating to follicle atresia and luteinization, suggesting that SKIL may play roles in these processes (Xu *et al.* 2009). VEGF (Redmer *et al.* 2001, Fraser *et al.* 2005), FGF (Shimizu *et al.* 2002, 2003), and TGF β superfamily members (Tomic *et al.* 2002) have also been linked to follicle atresia.

Major factors influencing theca cell differentiation

The current evidence suggests that an activated follicle produces factors that induce thecal cell differentiation from stroma, but the exact identity and combination of the proteins remain unknown. Not unexpectedly, it appears that no single factor appears to be responsible, but complex networks of signals function synergistically to result in a fully functional steroidogenic thecal layer surrounding a developing follicle.

A recent *in vitro* study using bovine ovarian tissue showed that ovarian stromal cells cultured in the presence of granulosa cells from small antral follicles transformed into putative thecal cells with increased lipid droplets and androstenedione production (Orisaka *et al.* 2006). Studies suggest that granulosa cells, but not activated oocytes, are involved in the functional differentiation and acquisition of LH responsiveness in stromal cells, and the cellular origin of the stroma determines whether or not granulosa cells influence thecal cell differentiation and functionality such as expression of necessary steroidogenic enzymes and LHRs (Orisaka *et al.* 2006). It appears that stromal cells from the cortical region may be preprogrammed differently to medullary stromal cells, and able to respond to granulosa cell communication in a manner that medullary stromal cells are not.

Using neonatal mouse ovaries, putative thecal stem cells were purified and induced to differentiate *in vitro* (Honda *et al.* 2007). When these cells were treated with LH, insulin-like growth factor 1 (IGF1), stem cell factor (SCF, also known as kit ligand), or granulosa cell-conditioned media, the cells differentiated into thecal cells, and showed signs of lipid droplet accumulation, formation of smooth endoplasmic reticulum,

mitochondria with tubular cristae, and produced androstenedione at later stages of culture. When these putative thecal cells were injected into ovaries of live mice, they were found to surround growing follicles akin to natural thecal cells *in vivo*.

Various factors have been studied *in vitro* for their effects on promoting thecal steroidogenesis. Selected molecules are discussed in the following sections.

Insulin-like growth factor

The ovary has a complete repertoire of IGF system components. IGF receptors are found on human thecal cells (Poretsky *et al.* 1985), and IGF1, the synthesis of which is regulated by FSH in the granulosa cells (Adashi *et al.* 1985, Hammond *et al.* 1985, Hernandez *et al.* 1989, Oliver *et al.* 1989), increases thecal cell proliferation *in vitro* (Hillier *et al.* 1991a, Stewart *et al.* 1995, Spicer & Chamberlain 1998, Huang *et al.* 2001, Mazerbourg & Hsueh 2003, Campbell *et al.* 2006, Kwintkiewicz & Giudice 2009). IGF1 alone stimulated the expression of LHRs (Magoffin & Weitsman 1994) and steroidogenic enzymes, CYP11A1 and HSD3B but not CYP17, and acts synergistically with LH to increase expression of these enzymes (Magoffin & Weitsman 1993a, 1993b, 1993c), and hence androgen synthesis in thecal cells *in vitro* (Hillier *et al.* 1991b; see Fig. 2). Interestingly, human stromal tissue cultured *in vitro* synthesized androgens when stimulated by insulin and IGF (Barbieri *et al.* 1983, 1984) giving additional evidence of IGF involvement in promoting thecal differentiation (see Fig. 4). IGF2 can stimulate bovine thecal steroidogenesis and acts through the type 1 IGF receptors (Spicer *et al.* 2004), so it may also play important roles in theca functionality.




Figure 4 Thecal cell recruitment and differentiation. Evidence suggests that thecal stem cells reside in the ovarian stroma and are recruited by factors released from follicles after they are activated. Individual factors may be responsible for the recruitment of these cells, whereas others may promote differentiation and proliferation. The origin of these factors is not currently clear, but it appears that a complex and tightly controlled set of signals from multiple factors is required for thecal recruitment and differentiation. AMH, anti-Müllerian hormone; BMP, bone morphogenetic protein; GDF9, growth differentiation factor 9; IGF, insulin-like growth factor; KGF, keratinocyte growth factor; LIF, leukemia inhibitory factor; SCF, stem cell factor (kit ligand).

Stem cell factor (SCF/kit ligand)

SCF, also known as kit ligand, is a growth factor that acts through the c-kit tyrosine kinase receptor (Zsebo *et al.* 1990, Besmer 1991), is synthesized by granulosa cells, and acts on differentiated thecal cells as well as undifferentiated stromal cells and oocytes where the receptor c-kit is located (Manova *et al.* 1990, 1993, Motro *et al.* 1991, Horie & Broxmeyer 1993, Motro & Bernstein 1993, Parrott & Skinner 1997). The addition of neutralizing antibodies to SCF and IGF1 together to follicle-conditioned media reduced the stimulatory effects on theca cell differentiation by more than 90% *in vitro* (Huang *et al.* 2001). In rat theca, neither SCF nor IGF1 alone stimulated androstenedione production, whereas, in combination, these factors dose dependently induced androgen production, but to a lesser extent than that induced by LH. The effects of adding these factors together with LH showed that IGF1 acted synergistically with LH to increase androgen production, whereas SCF had no added effect when in combination with LH. SCF alone decreased CYP17 and had no effect on the expression of CYP11A, HSD3B, or LHR expression (see Fig. 2). IGF1 alone had no effect on the expression of STAR and CYP17, but increased mRNA levels of LHR, CYP11A, and HSD3B. However, the combination of IGF1 and SCF increased the expression of STAR, CYP11A, CYP17, HSD3B, and LHR, giving strong evidence that these factors may act synergistically to regulate thecal cell differentiation into steroid-producing cells, at least in the rat.

SCF stimulated the growth of bovine primary cell cultures of theca and stromal cells under sub-confluent conditions *in vitro*, but when the cells were grown to confluence, SCF stimulated androstenedione production (Parrott & Skinner 1997). In contrast, SCF did not affect human chorionic gonadotropin-induced androgen production by bovine stromal cells (Parrott & Skinner 2000) suggesting that, in contrast to the rat, SCF alone affects bovine thecal cells once they have been differentiated.

The expression of SCF can also be modulated by leukemia inhibitory factor (LIF), keratinocyte growth factor (KGF), and hepatocyte growth factor (HGF; Parrott *et al.* 1994, Parrott & Skinner 1998, Nilsson *et al.* 2002). Thecal cells produce KGF and HGF (Parrott *et al.* 1994), which act on granulosa cells to produce SCF, which then signals back to the theca to produce KGF and HGF in a positive feedback mechanism (Parrott & Skinner 1998; see Fig. 5). Therefore, SCF may potentially act as a final common factor involved in thecal cell differentiation and activation.

Basic bFGF

Basic FGF (bFGF) has been shown to affect somatic cell mitosis, steroid synthesis, differentiation, and apoptosis (Tilly *et al.* 1992, Lavranos *et al.* 1994, Vernon & Spicer 1994). bFGF is expressed by primordial and primary oocytes, and granulosa cells of larger preantral follicles, and in the theca of rodent, bovine, and human follicles (van Wezel *et al.* 1995, Yamamoto *et al.* 1997, Berisha *et al.* 2000, Nilsson *et al.* 2001). The receptors for bFGF




Figure 5 Androgen production from theca cells. Factors released by cells comprising the follicle (granulosa cells, thecal cells, and oocytes) can modulate androgen production in thecal cells in addition to external influences such as gonadotropins and insulin. These molecules can stimulate (green) or inhibit (red) thecal androgen production both directly and/or indirectly. Specific molecules have been observed to act in opposing manners, indicating species-specific differences. bFGF, basic fibroblast growth factor; BMP, bone morphogenetic protein; GDF9, growth differentiation factor 9; HGF, hepatocyte growth factor; IGF, insulin-like growth factor; IGFBP, IGF-binding protein; IL1, interleukin-1; KGF, keratinocyte growth factor; LIF, leukemia inhibitory factor; SCF, stem cell factor/kit ligand; TGFB, transforming growth factor β ; TNF, tumor necrosis factor α .

have been found in granulosa and theca cells (Shikone *et al.* 1992, Wandji *et al.* 1992, Shimizu *et al.* 2002, Schams *et al.* 2009). In rat follicle cultures, bFGF activated primordial follicle development to the same extent as SCF (Nilsson *et al.* 2001) and also promoted cell growth in bovine thecal and stromal cells, and is therefore thought to act in a similar fashion to SCF by regulating somatic cell growth and development (Nilsson *et al.* 2001). While bFGF may also influence thecal development indirectly through stimulating SCF expression (Nilsson & Skinner 2004), bFGF, at least in the mouse, is not essential for folliculogenesis since the bFGF null mouse is fertile (Ortega *et al.* 1998).

TGFB superfamily members

TGFB superfamily members are now well established as having vital roles in controlling follicular growth and development (Shimasaki *et al.* 2004, Knight & Glister 2006, Xia & Schneyer 2009). The superfamily consists of a large group of proteins that include the bone morphogenetic proteins (BMPs), growth differentiation factors (GDFs), anti-Müllerian hormone (AMH; also known as Müllerian inhibiting substance (MIS)), activins, and inhibin, all of which are expressed in the ovary. These molecules bind to the BMP/TGFB/activin receptors to initiate phosphorylation cascades to influence gene expression in the cell nucleus. BMPs act through the SMAD1/5/8 pathway, whereas TGFB/activin/GDFs act through SMAD2/3. Evidence shows that specific members are involved in thecal cell recruitment, proliferation, and differentiation, which may or may not interact with the gonadotropins. Specific proteins will be discussed in this section, outlining the relevant research in this area and how each factor may contribute to theca growth and function.

Activin and inhibin

Activin and inhibin influence follicle activation, hormone synthesis, and luteolysis within the ovary (reviewed by Knight & Glister (2001)). Activin promoted the development of preantral follicles in sheep (Thomas *et al.* 2003) and human (Telfer *et al.* 2008) ovarian strips in culture, and this was also observed in rat studies (Li *et al.* 1995, Zhao *et al.* 2001), while in the mouse, activin from secondary follicles inhibited activation of small follicles (Mizunuma *et al.* 1999).

Studies on isolated human thecal cells cultured *in vitro* showed that activin suppressed androgen and progesterone production (Hillier *et al.* 1991a, Shukovski *et al.* 1993), while the activin antagonist inhibin enhanced LH-mediated androgen production (Hillier *et al.* 1991b). It has been suggested that granulosa cells secrete inhibin to control the amount of androgen synthesized by theca cells as substrates for estrogen synthesis in granulosa cells (Hillier *et al.* 1991b).

Thus, activins have direct effects on thecal cell function, and are regulated by the extracellular antagonists inhibin and follistatin (Findlay 1993, Welt *et al.* 2002).

In some species, an indirect effect of activins may occur through stimulation of granulosa cell proliferation or preantral follicle development (Li *et al.* 1995, Choi *et al.* 2008), which may be enhanced by additional IGF1 (Li *et al.* 1998). Activin can act by upregulating FSH receptors and aromatase gene expression in granulosa cells, and is involved in promoting estradiol production (Nakamura *et al.* 1993, El-Hefnawy & Zeleznik 2001, Ogawa *et al.* 2003, Park *et al.* 2005). Furthermore, since estradiol may suppress activin expression, this forms a possible interaction between activin and estrogen signaling during folliculogenesis (Kipp *et al.* 2007). Increased estradiol and inhibin production by the putative preovulatory follicle(s) would act to suppress activin production through estradiol, and block activin action at the theca through inhibin, thus enhancing thecal androgen production.

Inhibin is a critical factor in the control of steroid production and for control of gonadotropin secretion (McNeilly 2001, Padmanabhan & McNeilly 2001, McNeilly *et al.* 2003), but the effects of inhibin on thecal cell recruitment and differentiation are not known. Inhibin alone increased androgen production from human thecal cells in culture and also blocked the inhibitory effect of added activins (Hillier *et al.* 1991b). Inhibin is thought to modulate hormone production through antagonizing activin and BMPs, rather than through a signal cascade of its own (Wiater & Vale 2003, Farnworth *et al.* 2006).

Bone morphogenetic proteins

During folliculogenesis, BMPs are released at specific time points and act in either an autocrine or paracrine manner to modulate growth, differentiation, and function of follicular cells. BMP expression patterns have been investigated in rodents, ruminants, and primates, and observations suggest that species-specific differences occur for some molecules (reviewed by Shimasaki *et al.* (2004)).

BMP4 is expressed in the stromal cells surrounding primordial follicles, and BMP4/7 are expressed in the theca layer of antral follicles (Shimasaki *et al.* 1999, Nilsson & Skinner 2003, Lee *et al.* 2004). In neonatal rat ovaries, BMP4/7 increased the formation of primary follicles (Nilsson & Skinner 2003) and the percentage of growing follicles in the adult rat ovary (Nilsson & Skinner 2003, Lee *et al.* 2004).

BMP6 is produced by oocytes, but its function appears to differ between rodents and ruminants (Otsuka *et al.* 2001, Glister *et al.* 2004, Shi *et al.* 2009b). In sheep (Souza *et al.* 2002) and bovine (Kayani *et al.* 2009) ovaries, full complements of BMP/activin receptors were observed in granulosa, theca, and luteal tissues.

Human theca-derived tumor cells show reduced androstenedione production *in vitro* when treated with BMP4, and this effect is enhanced in the presence of cAMP agonists (Dooley *et al.* 2000). BMPs 2/4/6/7 all significantly decreased androstenedione secretion from ovine and bovine thecal cells *in vitro* while moderately increasing progesterone production and cell numbers (Glister *et al.* 2005, Campbell *et al.* 2006), and reducing *CYP17* gene expression, and to a lesser effect for *STAR*, *CYP11A1*, and *HSD3B* (Glister *et al.* 2005; see Fig. 2). The BMP antagonist, chordin, reversed the inhibitory effects of BMP7 on androgen production in bovine theca cells (Glister *et al.* 2005), while gremlin selectively reversed the effects of BMP4 but not BMP6 or 7. The BMP antagonist follistatin did not affect BMP-inhibited androgen production (Glister *et al.* 2005). Taken together, observations indicate that BMPs may act directly on theca to inhibit androgen production and also function to regulate the expression of factors from granulosa cells that act in a paracrine manner on thecal steroidogenesis. There are many BMP mutations that lead to aberrant fertility (McNatty *et al.* 2005), and these proteins may play roles in theca recruitment.

Growth differentiation factor 9

GDF9 is primarily produced specifically by the oocyte (McNatty *et al.* 2004), although it may also be produced in human (Shi *et al.* 2009a) and porcine (Paradis *et al.* 2009) granulosa and porcine theca cells (Paradis *et al.* 2009). Mutations of the *GDF9* gene lead to arrested folliculogenesis at the primary stage in mice, sheep, and humans (Shimasaki *et al.* 2004, Laissue *et al.* 2006, Kovanci *et al.* 2007, Nicol *et al.* 2009), suggesting that GDF9 is not vital for follicle activation but it is vital for onward primary growth. Follicles from GDF9 null mice and sheep (Thoka) lack supporting thecal cells indicating a role for GDF9 in the regulation of thecal recruitment, differentiation, and proliferation, but this appears to be dependent on the stage of follicle development (Dong *et al.* 1996, Elvin *et al.* 1999a, 1999b, Nicol *et al.* 2009). GDF9 alone, and in combination with IGF1, stimulated bovine thecal cell proliferation, but was found to inhibit IGF1- and LH-induced progesterone and androgen production, as well as decreasing LHR, LH-induced cAMP, and *CYP11A1* expression levels without altering IGF1 receptor, *STAR*, or *CYP17* levels (Spicer *et al.* 2008; see Fig. 2). The level of proliferation appeared higher in theca cells from small follicles compared to those from large follicles, and this may be related to higher levels of the putative GDF9 receptor, *ALK5*, in theca cells from small follicles. Thus, in small follicles, GDF9 could enhance proliferation, yet have no effect on promoting differentiation of thecal cells. GDF9 increased androgen production in rat thecal cells (Solovyeva *et al.* 2000), but reduced androgen production from human thecal cells indicating some

important species-specific differences (Yamamoto *et al.* 2002). Alternatively, the difference may be due to luteinization of the cells in culture.

GDF9 may also act in an indirect manner to modulate theca cell function, perhaps through regulating *SCF* expression (Dong *et al.* 1996, Joyce *et al.* 2000, Nilsson & Skinner 2002, Wang & Roy 2004). GDF9 also stimulated inhibin production (Hayashi *et al.* 1999, Kaivo-Oja *et al.* 2003, Roh *et al.* 2003), and the GDF9–inhibin- α double knockout mouse model is observed to have morphological thecal cells surrounding preantral follicles (Wu *et al.* 2004). However, these cells do not appear to have been differentiated into thecal cells since the expression of theca cell markers such as *CYP17A1*, *LHR*, or *KIT* are absent. These findings indicate that recruitment of putative theca-like cells can occur in the absence of GDF9, but differentiation of these cells does not appear to occur. GDF9 appears to have an indirect function on thecal cells, perhaps by promoting granulosa cell proliferation.

Transforming growth factor β

Both *TGFB1* and *TGFB2* are present in theca cells from follicles at the small preantral stage of development onwards and in stromal tissue and vascular systems in sheep ovaries, but not in granulosa cells or oocytes (Juengel *et al.* 2004). The receptors, *TGFB1* and *TGFB2*, had variable expression, and R1 was found in stromal and vascular cells, whereas R2 mRNA was found in thecal cells from the preantral stage onwards through development, as well as in the surface epithelium and some stromal cells. Furthermore, latent *TGFB*-binding proteins, which affect the bioavailability of *TGFBs* in tissues, are localized to the ovarian cortical stroma and theca externa of bovine antral follicles (Prodoehl *et al.* 2009). *TGFB1* has been reported to suppress androgen synthesis from human and rat thecal cells (Fournet *et al.* 1996, Attia *et al.* 2000; see Fig. 5), while in the mouse, *TGFB1* increased *Cyp11a*, *Cyp17*, and *Hsd3b* gene expression at various times during the culture period (Fournet *et al.* 1996), but inhibited *STAR* expression in a human thecal-like tumor cell line (Attia *et al.* 2000; see Fig. 2). A study using whole rat ovarian dispersed cell cultures suggested that *TGFB1* blocks steroidogenesis at the level of *CYP17* (Hernandez *et al.* 1990). These results are not clear-cut, but it appears that in the human at least, *TGFB* has a similar role to activin and BMPs in suppressing androgen production.

Hedgehog proteins

The hedgehog pathway has recently been shown to intersect with pathways involved with *FGF* receptor 2, BMPs, and other regulatory networks (Katoh 2009). Theca cells appear to be modulated by hedgehog signaling with the expression of hedgehog target genes

Ptch1, *Ptch2*, *Hip1*, and *Gli1* all present within theca cells (Wijgerde *et al.* 2005). Hedgehog proteins are expressed in the granulosa cells, but oocytes are unable to respond since they do not contain the necessary receptors. Hedgehog target genes are expressed in the pre-thecal cell compartment, and are therefore possible markers of pre-thecal cells and potentially involved in inducing theca cell differentiation. In cultured bovine thecal cells, sonic hedgehog-induced cell proliferation and androstenedione production (see Fig. 4), and hedgehog genes were shown to activate *Gli1* transcription factor in thecal cells (Spicer *et al.* 2009). In a transgenic mouse model, the hedgehog pathway was dominantly activated, and these mice displayed defective thecal development with reduced or absent smooth muscle actin normally seen in the thecal layer of growing follicles (Ren *et al.* 2009). The dominant activation of the hedgehog pathway therefore appears to block the differentiation of precursor cells into muscle cells that are normally located in the outer thecal layers, and are perhaps required for ovulation.

Synergism

A common trend appears to be emerging where the TGF β superfamily members are involved in fine tuning the modulation of androgen biosynthesis and steroidogenic enzyme gene expression. TGF β s, activin, and GDF9 all signal through SMAD2/3, whereas BMPs signal through the SMAD1/5/8 signaling pathway (Shimasaki *et al.* 2004). It makes sense that these two independent pathways influence the expression of completely separate groups of genes; otherwise, they would simply utilize the same pathway for the same effects. If the two pathways do in fact activate separate sets of genes, then these genes appear to be having similar effects on steroidogenesis. BMP4/6/7 suppress STAR, CYP11A1, CYP17, and HSD3B (Glister *et al.* 2005), and TGF β acts similarly by suppressing CYP17. GDF9 conversely appears to function in a more complicated manner, where it increases CYP17 expression while suppressing CYP11A1 levels. These contrasting results in different species coincide with the observed effects on androgen production, where in rat theca cells, GDF9 enhanced androgen synthesis (Solovyeva *et al.* 2000), but in bovine theca cells, GDF9 acted in an inhibitory manner (Spicer *et al.* 2008). There may be important species-specific differences in the function of GDF9 in particular, and this may also be the case with other TGF β superfamily members and therefore requires closer investigation.

There are also important species differences between rodents and humans with regard to stem cells. Mouse ES cells require LIF (Smith *et al.* 1992) and BMPs (Ying *et al.* 2003) to maintain pluripotency, whereas human counterparts rely on activin/nodal (Vallier *et al.* 2004, 2005, James *et al.* 2005) and FGFs (Xu *et al.* 2005). However, studies have shown that the same genes;

POU5F1 (*OCT4*), *SOX2*, and *NANOG*, are required for pluripotency in both species. Activin/nodal signaling controls expression of the key pluripotency factor *NANOG*. *NANOG* prevents differentiation induced by FGF signaling and limits transcriptional activity of SMAD2/3 (Vallier *et al.* 2009). By studying each factor in isolation, one is able to infer a specific function and role in ovarian folliculogenesis. However, we know that *in vivo* this is clearly not the case, and many different factors at varying, tightly controlled, concentrations all work synergistically to control the very delicate balance between the life and death of an ovarian follicle. The level and pattern of gene expression are vital factors; moreover, crosstalk between pathways and the presence of antagonists are additional levels of control for folliculogenesis and are yet to be fully elucidated.

Concluding remarks

Classically, it was thought that the oocyte was passively carried along the developmental process, and its maturation was controlled entirely by the production of endocrine hormones and surrounding somatic cell factors influencing the follicle as a whole. The latest concept in reproductive biology is that the oocyte itself is actively involved in regulating the surrounding somatic cells in order to provide an environment suitable for its own maturation. With this new and exciting theory in mind, it is possible that the oocyte itself is responsible for sending the signal for primordial follicle activation and thecal cell recruitment. However, since oocytes produce only limited factors, it is most likely that the interaction and communication between the oocyte and its somatic cells control the follicle development as a whole, and when one component fails, the entire process is halted. Nevertheless, thecal cells are vital for folliculogenesis in the ovary. They are specialized cells that are recruited to surround an activated follicle and provide structural support at first, and then by proliferating and differentiating, and acquiring a capillary network, they have become essential components of developing follicles. Their primary function is to synthesize androgens which act as substrates for estrogen production in granulosa cells, which is crucial for the pituitary–gonadal axis and endocrine control of reproduction. Androgen production is largely under the control of LH produced by the pituitary and transported to thecal cells via the blood stream. However, it is now clear that many other factors play an essential and important role in the modulation of theca function, including IGFs, insulin, FGF, SCF, TGF β superfamily members, and their related pathways and regulators. Theca cells have been somewhat forgotten more recently, where topical research has focused on granulosa cells and oocytes, but these specialized cells have a highly significant role in follicular function and are crucial for normal follicular development.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was funded by MRC (G7.002.0007.01) to A S McNeilly, which supports J M Young.

Acknowledgements

We would like to thank Ted Pinner for preparation of the figures, and Bruce Campbell for useful discussions.

References

- Adashi EY, Resnick CE, D'Ercole AJ, Svoboda ME & Van Wyk JJ 1985 Insulin-like growth factors as intraovarian regulators of granulosa cell growth and function. *Endocrine Reviews* **6** 400–420. (doi:10.1210/edrv-6-3-400)
- Attia GR, Dooley CA, Rainey WE & Carr BR 2000 Transforming growth factor beta inhibits steroidogenic acute regulatory (StAR) protein expression in human ovarian thecal cells. *Molecular and Cellular Endocrinology* **170** 123–129. (doi:10.1016/S0303-7207(00)00335-X)
- Baird DT, Swanston I & Scaramuzzi RJ 1976 Pulsatile release of LH and secretion of ovarian steroids in sheep during the luteal phase of the estrous cycle. *Endocrinology* **98** 1490–1496. (doi:10.1210/endo-98-6-1490)
- Baird DT, Swanston IA & McNeilly AS 1981 Relationship between LH, FSH, and prolactin concentration and the secretion of androgens and estrogens by the preovulatory follicle in the ewe. *Biology of Reproduction* **24** 1013–1025. (doi:10.1095/biolreprod24.5.1013)
- Bao B & Garverick HA 1998 Expression of steroidogenic enzyme and gonadotropin receptor genes in bovine follicles during ovarian follicular waves: a review. *Journal of Animal Science* **76** 1903–1921.
- Barbieri RL, Makris A & Ryan KJ 1983 Effects of insulin on steroidogenesis in cultured porcine ovarian theca. *Fertility and Sterility* **40** 237–241.
- Barbieri RL, Makris A & Ryan KJ 1984 Insulin stimulates androgen accumulation in incubations of human ovarian stroma and theca. *Obstetrics and Gynecology* **64** 735–80S.
- Barboni B, Turriani M, Galeati G, Spinaci M, Bacci ML, Forni M & Mattioli M 2000 Vascular endothelial growth factor production in growing pig antral follicles. *Biology of Reproduction* **63** 858–864. (doi:10.1095/biolreprod63.3.858)
- Barnett KR, Schilling C, Greenfield CR, Tomic D & Flaws JA 2006 Ovarian follicle development and transgenic mouse models. *Human Reproduction Update* **12** 537–555. (doi:10.1093/humupd/dml022)
- Berisha B, Schams D, Kosmann M, Amselgruber W & Einspanier R 2000 Expression and tissue concentration of vascular endothelial growth factor, its receptors, and localization in the bovine corpus luteum during estrous cycle and pregnancy. *Biology of Reproduction* **63** 1106–1114. (doi:10.1095/biolreprod63.4.1106)
- Besmer P 1991 The kit ligand encoded at the murine Steel locus: a pleiotropic growth and differentiation factor. *Current Opinion in Cell Biology* **3** 939–946. (doi:10.1016/0955-0674(91)90111-B)
- Campbell BK, Scaramuzzi RJ & Webb R 1995 Control of antral follicle development and selection in sheep and cattle. *Journal of Reproduction and Fertility. Supplement* **49** 335–350.
- Campbell BK, Baird DT & Webb R 1998 Effects of dose of LH on androgen production and luteinization of ovine theca cells cultured in a serum-free system. *Journal of Reproduction and Fertility* **112** 69–77. (doi:10.1530/jrf.0.1120069)
- Campbell BK, Souza CJ, Skinner AJ, Webb R & Baird DT 2006 Enhanced response of granulosa and theca cells from sheep carriers of the FecB mutation *in vitro* to gonadotropins and bone morphogenetic protein-2, -4, and -6. *Endocrinology* **147** 1608–1620. (doi:10.1210/en.2005-0604)
- Choi J, Lee B, Lee E, Yoon BK & Choi D 2008 Effect of activin A and insulin-like growth factor-I on *in vitro* development of preantral follicles isolated from cryopreserved ovarian tissues in the mouse. *Cryobiology* **57** 209–215. (doi:10.1016/j.cryobiol.2008.08.004)
- Clark LJ, Irving-Rodgers HF, Dharmarajan AM & Rodgers RJ 2004 Theca interna: the other side of bovine follicular atresia. *Biology of Reproduction* **71** 1071–1078. (doi:10.1095/biolreprod.104.029652)
- Connolly DT 1991 Vascular permeability factor: a unique regulator of blood vessel function. *Journal of Cellular Biochemistry* **47** 219–223. (doi:10.1002/jcb.240470306)
- Craig J, Orisaka M, Wang H, Orisaka S, Thompson W, Zhu C, Kotsuji F & Tsang BK 2007 Gonadotropin and intra-ovarian signals regulating follicle development and atresia: the delicate balance between life and death. *Frontiers in Bioscience* **12** 3628–3639. (doi:10.2741/2339)
- Deheuninck J & Luo K 2009 Ski and SnoN, potent negative regulators of TGF-beta signaling. *Cell Research* **19** 47–57. (doi:10.1038/cr.2008.324)
- D'Haeseleer M, Cocquyt G, Van Cruchten S, Simoens P & Van den Broeck W 2006 Cell-specific localisation of apoptosis in the bovine ovary at different stages of the oestrous cycle. *Theriogenology* **65** 757–772. (doi:10.1016/j.theriogenology.2005.07.008)
- Dong J, Albertini DF, Nishimori K, Kumar TR, Lu N & Matzuk MM 1996 Growth differentiation factor-9 is required during early ovarian folliculogenesis. *Nature* **383** 531–535. (doi:10.1038/383531a0)
- Dooley CA, Attia GR, Rainey WE, Moore DR & Carr BR 2000 Bone morphogenetic protein inhibits ovarian androgen production. *Journal of Clinical Endocrinology and Metabolism* **85** 3331–3337. (doi:10.1210/jc.85.9.3331)
- Edson MA, Nagaraja AK & Matzuk MM 2009 The mammalian ovary from genesis to revelation. *Endocrine Reviews* **30** 624–712. (doi:10.1210/er.2009-0012)
- El-Hefnawy T & Zeleznik AJ 2001 Synergism between FSH and activin in the regulation of proliferating cell nuclear antigen (PCNA) and cyclin D2 expression in rat granulosa cells. *Endocrinology* **142** 4357–4362. (doi:10.1210/en.142.10.4357)
- Elvin JA, Clark AT, Wang P, Wolfman NM & Matzuk MM 1999a Paracrine actions of growth differentiation factor-9 in the mammalian ovary. *Molecular Endocrinology* **13** 1035–1048. (doi:10.1210/me.13.6.1035)
- Elvin JA, Yan C, Wang P, Nishimori K & Matzuk MM 1999b Molecular characterization of the follicle defects in the growth differentiation factor 9-deficient ovary. *Molecular Endocrinology* **13** 1018–1034. (doi:10.1210/me.13.6.1018)
- Erickson GF, Magoffin DA, Dyer CA & Hofeditz C 1985 The ovarian androgen producing cells: a review of structure/function relationships. *Endocrine Reviews* **6** 371–399. (doi:10.1210/edrv-6-3-371)
- Farnworth PG, Stanton PG, Wang Y, Escalona R, Findlay JK & Ooi GT 2006 Inhibins differentially antagonize activin and bone morphogenetic protein action in a mouse adrenocortical cell line. *Endocrinology* **147** 3462–3471. (doi:10.1210/en.2006-0023)
- Ferrara N & Davis-Smyth T 1997 The biology of vascular endothelial growth factor. *Endocrine Reviews* **18** 4–25. (doi:10.1210/er.18.1.4)
- Findlay JK 1993 An update on the roles of inhibin, activin, and follistatin as local regulators of folliculogenesis. *Biology of Reproduction* **48** 15–23. (doi:10.1095/biolreprod48.1.15)
- Fournet N, Weitsman SR, Zachow RJ & Magoffin DA 1996 Transforming growth factor-beta inhibits ovarian 17 alpha-hydroxylase activity by a direct noncompetitive mechanism. *Endocrinology* **137** 166–174. (doi:10.1210/en.137.1.166)
- Fraser HM & Duncan WC 2009 SRB Reproduction, Fertility and Development Award Lecture 2008. Regulation and manipulation of angiogenesis in the ovary and endometrium. *Reproduction, Fertility, and Development* **21** 377–392. (doi:10.1071/RD08272)
- Fraser HM, Wilson H, Morris KD, Swanston I & Wiegand SJ 2005 Vascular endothelial growth factor Trap suppresses ovarian function at all stages of the luteal phase in the macaque. *Journal of Clinical Endocrinology and Metabolism* **90** 5811–5818. (doi:10.1210/jc.2005-1199)
- Glister C, Kemp CF & Knight PG 2004 Bone morphogenetic protein (BMP) ligands and receptors in bovine ovarian follicle cells: actions of BMP-4, -6 and -7 on granulosa cells and differential modulation of Smad-1 phosphorylation by follistatin. *Reproduction* **127** 239–254. (doi:10.1530/rep.1.00090)
- Glister C, Richards SL & Knight PG 2005 Bone morphogenetic proteins (BMP) -4, -6, and -7 potently suppress basal and luteinizing

- hormone-induced androgen production by bovine theca interna cells in primary culture: could ovarian hyperandrogenic dysfunction be caused by a defect in thecal BMP signaling? *Endocrinology* **146** 1883–1892. (doi:10.1210/en.2004-1303)
- Gougeon A** 1986 Dynamics of follicular growth in the human: a model from preliminary results. *Human Reproduction* **1** 81–87.
- Greenaway J, Gentry PA, Feige JJ, LaMarre J & Petrik JJ** 2005 Thrombospondin and vascular endothelial growth factor are cyclically expressed in an inverse pattern during bovine ovarian follicle development. *Biology of Reproduction* **72** 1071–1078. (doi:10.1095/biolreprod.104.031120)
- Hammond JM, Baranao JL, Skaleris D, Knight AB, Romanus JA & Rechler MM** 1985 Production of insulin-like growth factors by ovarian granulosa cells. *Endocrinology* **117** 2553–2555. (doi:10.1210/endo-117-6-2553)
- Hayashi M, McGee EA, Min G, Klein C, Rose UM, van Duin M & Hsueh AJ** 1999 Recombinant growth differentiation factor-9 (GDF-9) enhances growth and differentiation of cultured early ovarian follicles. *Endocrinology* **140** 1236–1244. (doi:10.1210/en.140.3.1236)
- Hernandez ER, Roberts CT Jr, LeRoith D & Adashi EY** 1989 Rat ovarian insulin-like growth factor I (IGF-I) gene expression is granulosa cell-selective: 5'-untranslated mRNA variant representation and hormonal regulation. *Endocrinology* **125** 572–574. (doi:10.1210/endo-125-1-572)
- Hernandez ER, Hurwitz A, Payne DW, Dharmarajan AM, Purchio AF & Adashi EY** 1990 Transforming growth factor-beta 1 inhibits ovarian androgen production: gene expression, cellular localization, mechanisms(s), and site(s) of action. *Endocrinology* **127** 2804–2811. (doi:10.1210/endo-127-6-2804)
- Hillier SG, Yong EL, Illingworth PJ, Baird DT, Schwall RH & Mason AJ** 1991a Effect of recombinant activin on androgen synthesis in cultured human thecal cells. *Journal of Clinical Endocrinology and Metabolism* **72** 1206–1211. (doi:10.1210/jcem-72-6-1206)
- Hillier SG, Yong EL, Illingworth PJ, Baird DT, Schwall RH & Mason AJ** 1991b Effect of recombinant inhibin on androgen synthesis in cultured human thecal cells. *Molecular and Cellular Endocrinology* **75** R1–R6. (doi:10.1016/0303-7207(91)90234-J)
- Hillier SG, Whitelaw PF & Smyth CD** 1994 Follicular oestrogen synthesis: the 'two-cell, two-gonadotrophin' model revisited. *Molecular and Cellular Endocrinology* **100** 51–54. (doi:10.1016/0303-7207(94)90278-X)
- Himelstein-Braw R, Byskov AG, Peters H & Faber M** 1976 Follicular atresia in the infant human ovary. *Journal of Reproduction and Fertility* **46** 55–59. (doi:10.1530/jrf.0.0460055)
- Hirshfield AN** 1994 Relationship between the supply of primordial follicles and the onset of follicular growth in rats. *Biology of Reproduction* **50** 421–428. (doi:10.1095/biolreprod50.2.421)
- Honda A, Hirose M, Hara K, Matoba S, Inoue K, Miki H, Hiura H, Kanatsu-Shinohara M, Kanai Y, Kono T et al.** 2007 Isolation, characterization, and *in vitro* and *in vivo* differentiation of putative thecal stem cells. *PNAS* **104** 12389–12394. (doi:10.1073/pnas.0703787104)
- Horie M & Broxmeyer HE** 1993 Involvement of immediate-early gene expression in the synergistic effects of steel factor in combination with granulocyte-macrophage colony-stimulating factor or interleukin-3 on proliferation of a human factor-dependent cell line. *Journal of Biological Chemistry* **268** 968–973.
- Huang CT, Weitsman SR, Dykes BN & Magoffin DA** 2001 Stem cell factor and insulin-like growth factor-I stimulate luteinizing hormone-independent differentiation of rat ovarian theca cells. *Biology of Reproduction* **64** 451–456. (doi:10.1095/biolreprod64.2.451)
- Hunter MG, Robinson RS, Mann GE & Webb R** 2004 Endocrine and paracrine control of follicular development and ovulation rate in farm species. *Animal Reproduction Science* **82–83** 461–477. (doi:10.1016/j.anireprosci.2004.05.013)
- Irving-Rodgers HF, van Wezel IL, Mussard ML, Kinder JE & Rodgers RJ** 2001 Atresia revisited: two basic patterns of atresia of bovine antral follicles. *Reproduction* **122** 761–775. (doi:10.1530/rep.0.1220761)
- Irving-Rodgers HF, Krupa M & Rodgers RJ** 2003 Cholesterol side-chain cleavage cytochrome P450 and 3beta-hydroxysteroid dehydrogenase expression and the concentrations of steroid hormones in the follicular fluids of different phenotypes of healthy and atretic bovine ovarian follicles. *Biology of Reproduction* **69** 2022–2028. (doi:10.1095/biolreprod.103.017442)
- James D, Levine AJ, Besser D & Hemmati-Brivanlou A** 2005 TGFbeta/activin/nodal signaling is necessary for the maintenance of pluripotency in human embryonic stem cells. *Development* **132** 1273–1282. (doi:10.1242/dev.01706)
- Jiang JY, Macchiarelli G, Tsang BK & Sato E** 2003 Capillary angiogenesis and degeneration in bovine ovarian antral follicles. *Reproduction* **125** 211–223. (doi:10.1530/rep.0.1250211)
- Jolly PD, Tisdall DJ, Heath DA, Lun S & McNatty KP** 1994 Apoptosis in bovine granulosa cells in relation to steroid synthesis, cyclic adenosine 3',5'-monophosphate response to follicle-stimulating hormone and luteinizing hormone, and follicular atresia. *Biology of Reproduction* **51** 934–944. (doi:10.1095/biolreprod51.5.934)
- Joyce IM, Clark AT, Pendola FL & Eppig JJ** 2000 Comparison of recombinant growth differentiation factor-9 and oocyte regulation of KIT ligand messenger ribonucleic acid expression in mouse ovarian follicles. *Biology of Reproduction* **63** 1669–1675. (doi:10.1095/biolreprod63.6.1669)
- Juengel JL, Bibby AH, Reader KL, Lun S, Quirke LD, Haydon LJ & McNatty KP** 2004 The role of transforming growth factor-beta (TGF-beta) during ovarian follicular development in sheep. *Reproductive Biology and Endocrinology* **2** 78. (doi:10.1186/1477-7827-2-78)
- Kaivo-Oja N, Bondestam J, Kamarainen M, Koskimies J, Vitt U, Cranfield M, Vuojolainen K, Kallio JP, Olkkonen VM, Hayashi M et al.** 2003 Growth differentiation factor-9 induces Smd2 activation and inhibin B production in cultured human granulosa-luteal cells. *Journal of Clinical Endocrinology and Metabolism* **88** 755–762. (doi:10.1210/jc.2002-021317)
- Katoh M** 2009 FGFR2 abnormalities underlie a spectrum of bone, skin, and cancer pathologies. *Journal of Investigative Dermatology* **129** 1861–1867. (doi:10.1038/jid.2009.97)
- Kayani AR, Glister C & Knight PG** 2009 Evidence for an inhibitory role of bone morphogenetic protein(s) in the follicular-luteal transition in cattle. *Reproduction* **137** 67–78. (doi:10.1530/REP-08-0198)
- Kerban A, Boerboom D & Sirois J** 1999 Human chorionic gonadotropin induces an inverse regulation of steroidogenic acute regulatory protein messenger ribonucleic acid in theca interna and granulosa cells of equine preovulatory follicles. *Endocrinology* **140** 667–674. (doi:10.1210/en.140.2.667)
- Kezele PR, Ague JM, Nilsson E & Skinner MK** 2005 Alterations in the ovarian transcriptome during primordial follicle assembly and development. *Biology of Reproduction* **72** 241–255. (doi:10.1095/biolreprod.104.032060)
- Kipp JL, Kilen SM, Woodruff TK & Mayo KE** 2007 Activin regulates estrogen receptor gene expression in the mouse ovary. *Journal of Biological Chemistry* **282** 36755–36765. (doi:10.1074/jbc.M705143200)
- Knight PG & Glister C** 2001 Potential local regulatory functions of inhibins, activins and follistatin in the ovary. *Reproduction* **121** 503–512. (doi:10.1530/rep.0.1210503)
- Knight PG & Glister C** 2006 TGF-beta superfamily members and ovarian follicle development. *Reproduction* **132** 191–206. (doi:10.1530/rep.1.01074)
- Kovanci E, Rohozinski J, Simpson JL, Heard MJ, Bishop CE & Carson SA** 2007 Growth differentiating factor-9 mutations may be associated with premature ovarian failure. *Fertility and Sterility* **87** 143–146. (doi:10.1016/j.fertnstert.2006.05.079)
- Kwintkiewicz J & Giudice LC** 2009 The interplay of insulin-like growth factors, gonadotropins, and endocrine disruptors in ovarian follicular development and function. *Seminars in Reproductive Medicine* **27** 43–51. (doi:10.1055/s-0028-1108009)
- Laissue P, Christin-Maitre S, Touraine P, Kuttann F, Ritvos O, Aittomaki K, Bourcigaux N, Jacquesson L, Bouchard P, Frydman R et al.** 2006 Mutations and sequence variants in GDF9 and BMP15 in patients with premature ovarian failure. *European Journal of Endocrinology* **154** 739–744. (doi:10.1530/eje.1.02135)
- Lavoie HA & King SR** 2009 Transcriptional regulation of steroidogenic genes: STARD1, CYP11A1 and HSD3B. *Experimental Biology and Medicine* **234** 880–907. (doi:10.3181/0903-MR-97)
- Lavranos TC, Rodgers HF, Bertonecello I & Rodgers RJ** 1994 Anchorage-independent culture of bovine granulosa cells: the effects of basic fibroblast growth factor and dibutyryl cAMP on cell division and differentiation. *Experimental Cell Research* **211** 245–251. (doi:10.1006/excr.1994.1084)

- Lee WS, Yoon SJ, Yoon TK, Cha KY, Lee SH, Shimasaki S, Lee S & Lee KA 2004 Effects of bone morphogenetic protein-7 (BMP-7) on primordial follicular growth in the mouse ovary. *Molecular Reproduction and Development* **69** 159–163. (doi:10.1002/mrd.20163)
- Li R, Phillips DM & Mather JP 1995 Activin promotes ovarian follicle development *in vitro*. *Endocrinology* **136** 849–856. (doi:10.1210/en.136.3.849)
- Li D, Kubo T, Kim H, Shimasaki S & Erickson GF 1998 Endogenous insulin-like growth factor-I is obligatory for stimulation of rat inhibin alpha-subunit expression by follicle-stimulating hormone. *Biology of Reproduction* **58** 219–225. (doi:10.1095/biolreprod58.1.219)
- Li M, Schatten H & Sun QY 2009a Androgen receptor's destiny in mammalian oocytes: a new hypothesis. *Molecular Human Reproduction* **15** 149–154. (doi:10.1093/molehr/gap006)
- Li Y, Liang XY, Wei LN, Xiong YL, Yang X, Shi HG & Yang ZH 2009b Study of RNA interference inhibiting rat ovarian androgen biosynthesis by depressing 17alpha-hydroxylase/17, 20-lyase activity *in vivo*. *Reproductive Biology and Endocrinology* **7** 73. (doi:10.1186/1477-7827-7-73)
- Logan KA, Juengel JL & McNatty KP 2002 Onset of steroidogenic enzyme gene expression during ovarian follicular development in sheep. *Biology of Reproduction* **66** 906–916. (doi:10.1095/biolreprod66.4.906)
- Lundy T, Smith P, O'Connell A, Hudson NL & McNatty KP 1999 Populations of granulosa cells in small follicles of the sheep ovary. *Journal of Reproduction and Fertility* **115** 251–262. (doi:10.1530/jrf.0.1150251)
- Magoffin DA 2005 Ovarian theca cell. *International Journal of Biochemistry & Cell Biology* **37** 1344–1349. (doi:10.1016/j.biocel.2005.01.016)
- Magoffin DA & Weitsman SR 1993a Differentiation of ovarian theca-interstitial cells *in vitro*: regulation of 17 alpha-hydroxylase messenger ribonucleic acid expression by luteinizing hormone and insulin-like growth factor-I. *Endocrinology* **132** 1945–1951. (doi:10.1210/en.132.5.1945)
- Magoffin DA & Weitsman SR 1993b Effect of insulin-like growth factor-I on cholesterol side-chain cleavage cytochrome P450 messenger ribonucleic acid expression in ovarian theca-interstitial cells stimulated to differentiate *in vitro*. *Molecular and Cellular Endocrinology* **96** 45–51. (doi:10.1016/0303-7207(93)90093-Y)
- Magoffin DA & Weitsman SR 1993c Insulin-like growth factor-I stimulates the expression of 3 beta-hydroxysteroid dehydrogenase messenger ribonucleic acid in ovarian theca-interstitial cells. *Biology of Reproduction* **48** 1166–1173. (doi:10.1095/biolreprod48.5.1166)
- Magoffin DA & Weitsman SR 1994 Insulin-like growth factor-I regulation of luteinizing hormone (LH) receptor messenger ribonucleic acid expression and LH-stimulated signal transduction in rat ovarian theca-interstitial cells. *Biology of Reproduction* **51** 766–775. (doi:10.1095/biolreprod51.4.766)
- Maisonpierre PC, Suri C, Jones PF, Bartunkova S, Wiegand SJ, Radziejewski C, Compton D, McClain J, Aldrich TH, Papadopoulos N *et al.* 1997 Angiopoietin-2, a natural antagonist for Tie2 that disrupts *in vivo* angiogenesis. *Science* **277** 55–60. (doi:10.1126/science.277.5322.55)
- Mamluk R, Greber Y & Meidan R 1999 Hormonal regulation of messenger ribonucleic acid expression for steroidogenic factor-1, steroidogenic acute regulatory protein, and cytochrome P450 side-chain cleavage in bovine luteal cells. *Biology of Reproduction* **60** 628–634. (doi:10.1095/biolreprod60.3.628)
- Manna PR, Dyson MT & Stocco DM 2009 Regulation of the steroidogenic acute regulatory protein gene expression: present and future perspectives. *Molecular Human Reproduction* **15** 321–333. (doi:10.1093/molehr/gap025)
- Manova K, Nocka K, Besmer P & Bachvarova RF 1990 Gonadal expression of c-kit encoded at the W locus of the mouse. *Development* **110** 1057–1069.
- Manova K, Huang EJ, Angeles M, De Leon V, Sanchez S, Pronovost SM, Besmer P & Bachvarova RF 1993 The expression pattern of the c-kit ligand in gonads of mice supports a role for the c-kit receptor in oocyte growth and in proliferation of spermatogonia. *Developmental Biology* **157** 85–99. (doi:10.1006/dbio.1993.1114)
- Martelli A, Russo V, Mauro A, Nardinocchi D, Rinaldi C, Bernabo N, Berardinelli P & Mattioli M 2008 Role of VEGF in pig preantral follicles. *Veterinary Research Communications* **32** (Supplement 1) S155–S157. (doi:10.1007/s11259-008-9114-2)
- Mattioli M, Barboni B, Turriani M, Galeati G, Zannoni A, Castellani G, Berardinelli P & Scapolo PA 2001 Follicle activation involves vascular endothelial growth factor production and increased blood vessel extension. *Biology of Reproduction* **65** 1014–1019. (doi:10.1095/biolreprod65.4.1014)
- Mazerbourg S & Hsueh AJ 2003 Growth differentiation factor-9 signaling in the ovary. *Molecular and Cellular Endocrinology* **202** 31–36. (doi:10.1016/S0303-7207(03)00058-3)
- McNatty KP, Heath DA, Lundy T, Fidler AE, Quirke L, O'Connell A, Smith P, Groome N & Tisdall DJ 1999 Control of early ovarian follicular development. *Journal of Reproduction and Fertility. Supplement* **54** 3–16.
- McNatty KP, Moore LG, Hudson NL, Quirke LD, Lawrence SB, Reader K, Hanrahan JP, Smith P, Groome NP, Laitinen M *et al.* 2004 The oocyte and its role in regulating ovulation rate: a new paradigm in reproductive biology. *Reproduction* **128** 379–386. (doi:10.1530/rep.1.00280)
- McNatty KP, Galloway SM, Wilson T, Smith P, Hudson NL, O'Connell A, Bibby AH, Heath DA, Davis GH, Hanrahan JP *et al.* 2005 Physiological effects of major genes affecting ovulation rate in sheep. *Genetics, Selection, Evolution* **37** (Supplement 1) S25–S38. (doi:10.1186/1297-9686-37-S1-S25)
- McNatty KP, Reader K, Smith P, Heath DA & Juengel JL 2007 Control of ovarian follicular development to the gonadotrophin-dependent phase: a 2006 perspective. *Society of Reproduction and Fertility Supplement* **64** 55–68.
- McNeilly AS 2001 Lactational control of reproduction. *Reproduction, Fertility, and Development* **13** 583–590. (doi:10.1071/RD01056)
- McNeilly JR, Saunders PT, Taggart M, Cranfield M, Cooke HJ & McNeilly AS 2000 Loss of oocytes in Dazl knockout mice results in maintained ovarian steroidogenic function but altered gonadotropin secretion in adult animals. *Endocrinology* **141** 4284–4294. (doi:10.1210/en.141.11.4284)
- McNeilly AS, Crawford JL, Taragnat C, Nicol L & McNeilly JR 2003 The differential secretion of FSH and LH: regulation through genes, feedback and packaging. *Reproduction Supplement* **61** 463–476.
- Mizunuma H, Liu X, Andoh K, Abe Y, Kobayashi J, Yamada K, Yokota H, Ibuki Y & Hasegawa Y 1999 Activin from secondary follicles causes small preantral follicles to remain dormant at the resting stage. *Endocrinology* **140** 37–42. (doi:10.1210/en.140.1.37)
- Montgomery GW, Galloway SM, Davis GH & McNatty KP 2001 Genes controlling ovulation rate in sheep. *Reproduction* **121** 843–852. (doi:10.1530/rep.0.1210843)
- Moor RM, Hay MF, Dott HM & Cran DG 1978 Macroscopic identification and steroidogenic function of atretic follicles in sheep. *Journal of Endocrinology* **77** 309–318. (doi:10.1677/joe.0.0770309)
- Morley P, Calaresu FR, Barbe GJ & Armstrong DT 1989 Insulin enhances luteinizing hormone-stimulated steroidogenesis by porcine theca cells. *Biology of Reproduction* **40** 735–743. (doi:10.1095/biolreprod40.4.735)
- Motro B & Bernstein A 1993 Dynamic changes in ovarian c-kit and Steel expression during the estrous reproductive cycle. *Developmental Dynamics* **197** 69–79. (doi:10.1002/aja.1001970107)
- Motro B, van der Kooy D, Rossant J, Reith A & Bernstein A 1991 Contiguous patterns of c-kit and steel expression: analysis of mutations at the W and Sl loci. *Development* **113** 1207–1221.
- Nakamura M, Minegishi T, Hasegawa Y, Nakamura K, Igarashi S, Ito I, Shinozaki H, Miyamoto K, Eto Y & Ibuki Y 1993 Effect of an activin A on follicle-stimulating hormone (FSH) receptor messenger ribonucleic acid levels and FSH receptor expressions in cultured rat granulosa cells. *Endocrinology* **133** 538–544. (doi:10.1210/en.133.2.538)
- Nicol L, Bishop SC, Pong-Wong R, Bendixen C, Holm LE, Rhind SM & McNeilly AS 2009 Homozygosity for a single base-pair mutation in the oocyte-specific GDF9 gene results in sterility in Thoka sheep. *Reproduction* **138** 921–933. (doi:10.1530/REP-09-0193)
- Nilsson EE & Skinner MK 2002 Growth and differentiation factor-9 stimulates progression of early primary but not primordial rat ovarian follicle development. *Biology of Reproduction* **67** 1018–1024. (doi:10.1095/biolreprod.101.002527)
- Nilsson EE & Skinner MK 2003 Bone morphogenetic protein-4 acts as an ovarian follicle survival factor and promotes primordial follicle development. *Biology of Reproduction* **69** 1265–1272. (doi:10.1095/biolreprod.103.018671)

- Nilsson EE & Skinner MK 2004 Kit ligand and basic fibroblast growth factor interactions in the induction of ovarian primordial to primary follicle transition. *Molecular and Cellular Endocrinology* **214** 19–25. (doi:10.1016/j.mce.2003.12.001)
- Nilsson E, Parrott JA & Skinner MK 2001 Basic fibroblast growth factor induces primordial follicle development and initiates folliculogenesis. *Molecular and Cellular Endocrinology* **175** 123–130. (doi:10.1016/S0303-7207(01)00391-4)
- Nilsson EE, Kezele P & Skinner MK 2002 Leukemia inhibitory factor (LIF) promotes the primordial to primary follicle transition in rat ovaries. *Molecular and Cellular Endocrinology* **188** 65–73. (doi:10.1016/S0303-7207(01)00746-8)
- Ogawa T, Yogo K, Ishida N & Takeya T 2003 Synergistic effects of activin and FSH on hyperphosphorylation of Rb and G1/S transition in rat primary granulosa cells. *Molecular and Cellular Endocrinology* **210** 31–38. (doi:10.1016/j.mce.2003.08.008)
- Oliver JE, Aitman TJ, Powell JF, Wilson CA & Clayton RN 1989 Insulin-like growth factor I gene expression in the rat ovary is confined to the granulosa cells of developing follicles. *Endocrinology* **124** 2671–2679. (doi:10.1210/endo-124-6-2671)
- Orisaka M, Tajima K, Mizutani T, Miyamoto K, Tsang BK, Fukuda S, Yoshida Y & Kotsuji F 2006 Granulosa cells promote differentiation of cortical stromal cells into theca cells in the bovine ovary. *Biology of Reproduction* **75** 734–740. (doi:10.1095/biolreprod.105.050344)
- Ortega S, Iltmann M, Tsang SH, Ehrlich M & Basilico C 1998 Neuronal defects and delayed wound healing in mice lacking fibroblast growth factor 2. *PNAS* **95** 5672–5677. (doi:10.1073/pnas.95.10.5672)
- O'Shea JD, Nightingale MG & Chamley WA 1977 Changes in small blood vessels during cyclical luteal regression in sheep. *Biology of Reproduction* **17** 162–177. (doi:10.1095/biolreprod17.2.162)
- O'Shea JD, Hay MF & Cran DG 1978 Ultrastructural changes in the theca interna during follicular atresia in sheep. *Journal of Reproduction and Fertility* **54** 183–187. (doi:10.1530/jrf.0.0540183)
- Otsuka F, Moore RK & Shimasaki S 2001 Biological function and cellular mechanism of bone morphogenetic protein-6 in the ovary. *Journal of Biological Chemistry* **276** 32889–32895. (doi:10.1074/jbc.M103212200)
- Padmanabhan V & McNeilly AS 2001 Is there an FSH-releasing factor? *Reproduction* **121** 21–30. (doi:10.1530/rep.0.1210021)
- Palermo R 2007 Differential actions of FSH and LH during folliculogenesis. *Reproductive Biomedicine Online* **15** 326–337. (doi:10.1016/S1472-6483(10)60347-1)
- Paradis F, Novak S, Murdoch GK, Dyck MK, Dixon WT & Foxcroft GR 2009 Temporal regulation of BMP2, BMP6, BMP15, GDF9, BMPR1A, BMPR1B, BMPR2 and TGFBR1 mRNA expression in the oocyte, granulosa and theca cells of developing preovulatory follicles in the pig. *Reproduction* **138** 115–129. (doi:10.1530/REP-08-0538)
- Park Y, Maizels ET, Feiger ZJ, Alam H, Peters CA, Woodruff TK, Unterman TG, Lee EJ, Jameson JL & Hunzicker-Dunn M 2005 Induction of cyclin D2 in rat granulosa cells requires FSH-dependent relief from FOXO1 repression coupled with positive signals from Smad. *Journal of Biological Chemistry* **280** 9135–9148. (doi:10.1074/jbc.M409486200)
- Parrott JA & Skinner MK 1997 Direct actions of kit-ligand on theca cell growth and differentiation during follicle development. *Endocrinology* **138** 3819–3827. (doi:10.1210/en.138.9.3819)
- Parrott JA & Skinner MK 1998 Thecal cell–granulosa cell interactions involve a positive feedback loop among keratinocyte growth factor, hepatocyte growth factor, and Kit ligand during ovarian follicular development. *Endocrinology* **139** 2240–2245. (doi:10.1210/en.139.5.2240)
- Parrott JA & Skinner MK 2000 Kit ligand actions on ovarian stromal cells: effects on theca cell recruitment and steroid production. *Molecular Reproduction and Development* **55** 55–64. (doi:10.1002/(SICI)1098-2795(200001)55:1<55::AID-MRD8>3.0.CO;2-L)
- Parrott JA, Vigne JL, Chu BZ & Skinner MK 1994 Mesenchymal–epithelial interactions in the ovarian follicle involve keratinocyte and hepatocyte growth factor production by thecal cells and their action on granulosa cells. *Endocrinology* **135** 569–575. (doi:10.1210/en.135.2.569)
- Peluso JJ, Downey MC & Gruenberg ML 1984 Role of LH pulse amplitude in controlling rat ovarian oestradiol-17 beta secretion *in vitro*. *Journal of Reproduction and Fertility* **71** 107–112. (doi:10.1530/jrf.0.0710107)
- Pollack SE, Furth EE, Kallen CB, Arakane F, Kiriakidou M, Kozarsky KF & Strauss JF III 1997 Localization of the steroidogenic acute regulatory protein in human tissues. *Journal of Clinical Endocrinology and Metabolism* **82** 4243–4251. (doi:10.1210/jc.82.12.4243)
- Poretzky L, Grigorescu F, Seibel M, Moses AC & Flier JS 1985 Distribution and characterization of insulin and insulin-like growth factor I receptors in normal human ovary. *Journal of Clinical Endocrinology and Metabolism* **61** 728–734. (doi:10.1210/jcem-61-4-728)
- Prodoehl MJ, Irving-Rodgers HF, Bonner WM, Sullivan TM, Micke GC, Gibson MA, Perry VE & Rodgers RJ 2009 Fibrillins and latent TGFbeta binding proteins in bovine ovaries of offspring following high or low protein diets during pregnancy of dams. *Molecular and Cellular Endocrinology* **307** 133–141. (doi:10.1016/j.mce.2009.03.002)
- Ravindranath N, Little-Ihrig L, Phillips HS, Ferrara N & Zeleznik AJ 1992 Vascular endothelial growth factor messenger ribonucleic acid expression in the primate ovary. *Endocrinology* **131** 254–260. (doi:10.1210/en.131.1.254)
- Redmer DA, Doraiswamy V, Bortnem BJ, Fisher K, Jablonka-Shariff A, Grazul-Bilska AT & Reynolds LP 2001 Evidence for a role of capillary pericytes in vascular growth of the developing ovine corpus luteum. *Biology of Reproduction* **65** 879–889. (doi:10.1095/biolreprod65.3.879)
- Ren Y, Cowan RG, Harman RM & Quirk SM 2009 Dominant activation of the hedgehog signaling pathway in the ovary alters theca development and prevents ovulation. *Molecular Endocrinology* **23** 711–723. (doi:10.1210/me.2008-0391)
- Rodgers RJ & Irving-Rodgers HF 2009 The morphological classification of bovine ovarian follicles. *Reproduction* **139** 309–318. (doi:10.1530/REP-09-0177)
- Roh JS, Bondestam J, Mazerbourg S, Kaivo-Oja N, Groome N, Ritvos O & Hsueh AJ 2003 Growth differentiation factor-9 stimulates inhibin production and activates Smad2 in cultured rat granulosa cells. *Endocrinology* **144** 172–178. (doi:10.1210/en.2002-220618)
- Ryan KE, Glistler C, Lonergan P, Martin F, Knight PG & Evans AC 2008 Functional significance of the signal transduction pathways Akt and Erk in ovarian follicles: *in vitro* and *in vivo* studies in cattle and sheep. *Journal of Ovarian Research* **1** 2. (doi:10.1186/1757-2215-1-2)
- Scaramuzzi RJ, Adams NR, Baird DT, Campbell BK, Downing JA, Findlay JK, Henderson KM, Martin GB, McNatty KP, McNeilly AS *et al.* 1993 A model for follicle selection and the determination of ovulation rate in the ewe. *Reproduction, Fertility, and Development* **5** 459–478. (doi:10.1071/RD9930459)
- Schallenger E, Schams D, Bullermann B & Walters DL 1984 Pulsatile secretion of gonadotrophins, ovarian steroids and ovarian oxytocin during prostaglandin-induced regression of the corpus luteum in the cow. *Journal of Reproduction and Fertility* **71** 493–501. (doi:10.1530/jrf.0.0710493)
- Schams D, Steinberg V, Steffl M, Meyer HH & Berisha B 2009 Expression and possible role of fibroblast growth factor family members in porcine antral follicles during final maturation. *Reproduction* **138** 141–149. (doi:10.1530/REP-09-0033)
- Senger DR, Van de Water L, Brown LF, Nagy JA, Yeo KT, Yeo TK, Berse B, Jackman RW, Dvorak AM & Dvorak HF 1993 Vascular permeability factor (VPF, VEGF) in tumor biology. *Cancer Metastasis Reviews* **12** 303–324. (doi:10.1007/BF00665960)
- Shi FT, Cheung AP, Huang HF & Leung PC 2009a Effects of endogenous growth differentiation factor 9 on activin A-induced inhibin B production in human granulosa-lutein cells. *Journal of Clinical Endocrinology and Metabolism* **94** 5108–5116. (doi:10.1210/jc.2009-1047)
- Shi J, Yoshino O, Osuga Y, Koga K, Hirota Y, Hirata T, Yano T, Nishii O & Taketani Y 2009b Bone morphogenetic protein-6 stimulates gene expression of follicle-stimulating hormone receptor, inhibin/activin beta subunits, and anti-Mullerian hormone in human granulosa cells. *Fertility and Sterility* **92** 1794–1798. (doi:10.1016/j.fertnstert.2009.05.004)
- Shikone T, Yamoto M & Nakano R 1992 Follicle-stimulating hormone induces functional receptors for basic fibroblast growth factor in rat granulosa cells. *Endocrinology* **131** 1063–1068. (doi:10.1210/en.131.3.1063)
- Shimasaki S, Zachow RJ, Li D, Kim H, Iemura S, Ueno N, Sampath K, Chang RJ & Erickson GF 1999 A functional bone morphogenetic protein system in the ovary. *PNAS* **96** 7282–7287. (doi:10.1073/pnas.96.13.7282)

- Shimasaki S, Moore RK, Otsuka F & Erickson GF 2004 The bone morphogenetic protein system in mammalian reproduction. *Endocrine Reviews* **25** 72–101. (doi:10.1210/er.2003-0007)
- Shimizu T, Jiang JY, Sasada H & Sato E 2002 Changes of messenger RNA expression of angiogenic factors and related receptors during follicular development in gilts. *Biology of Reproduction* **67** 1846–1852. (doi:10.1095/biolreprod.102.006734)
- Shimizu T, Kawahara M, Abe Y, Yokoo M, Sasada H & Sato E 2003 Follicular microvasculature and angiogenic factors in the ovaries of domestic animals. *Journal of Reproduction and Development* **49** 181–192. (doi:10.1262/jrd.49.181)
- Shukovski L, Dyson M & Findlay JK 1993 The effects of follistatin, activin and inhibin on steroidogenesis by bovine thecal cells. *Molecular and Cellular Endocrinology* **97** 19–27. (doi:10.1016/0303-7207(93)90207-Z)
- Smith AG, Nichols J, Robertson M & Rathjen PD 1992 Differentiation inhibiting activity (DIA/LIF) and mouse development. *Developmental Biology* **151** 339–351. (doi:10.1016/0012-1606(92)90174-F)
- Smith MF, Gutierrez CG, Ricke WA, Armstrong DG & Webb R 2005 Production of matrix metalloproteinases by cultured bovine theca and granulosa cells. *Reproduction* **129** 75–87. (doi:10.1530/rep.1.00381)
- Solovyeva EV, Hayashi M, Margi K, Barkats C, Klein C, Amsterdam A, Hsueh AJ & Tsafiri A 2000 Growth differentiation factor-9 stimulates rat theca-interstitial cell androgen biosynthesis. *Biology of Reproduction* **63** 1214–1218. (doi:10.1095/biolreprod63.4.1214)
- Souza CJ, Campbell BK, McNeilly AS & Baird DT 2002 Effect of bone morphogenetic protein 2 (BMP2) on oestradiol and inhibin A production by sheep granulosa cells, and localization of BMP receptors in the ovary by immunohistochemistry. *Reproduction* **123** 363–369. (doi:10.1530/rep.0.1230363)
- Spicer LJ & Chamberlain CS 1998 Influence of cortisol on insulin- and insulin-like growth factor 1 (IGF-1)-induced steroid production and on IGF-1 receptors in cultured bovine granulosa cells and thecal cells. *Endocrine* **9** 153–161. (doi:10.1385/ENDO:9:2:153)
- Spicer LJ, Voge JL & Allen DT 2004 Insulin-like growth factor-II stimulates steroidogenesis in cultured bovine thecal cells. *Molecular and Cellular Endocrinology* **227** 1–7. (doi:10.1016/j.mce.2004.08.003)
- Spicer LJ, Aad PY, Allen DT, Mazerbourg S, Payne AH & Hsueh AJ 2008 Growth differentiation factor 9 (GDF9) stimulates proliferation and inhibits steroidogenesis by bovine theca cells: influence of follicle size on responses to GDF9. *Biology of Reproduction* **78** 243–253. (doi:10.1095/biolreprod.107.063446)
- Spicer LJ, Sudo S, Aad PY, Wang LS, Chun SY, Ben-Shlomo I, Klein C & Hsueh AJ 2009 The hedgehog-patched signaling pathway and function in the mammalian ovary: a novel role for hedgehog proteins in stimulating proliferation and steroidogenesis of theca cells. *Reproduction* **138** 329–339. (doi:10.1530/REP-08-0317)
- Stewart RE, Spicer LJ, Hamilton TD & Keefer BE 1995 Effects of insulin-like growth factor I and insulin on proliferation and on basal and luteinizing hormone-induced steroidogenesis of bovine thecal cells: involvement of glucose and receptors for insulin-like growth factor I and luteinizing hormone. *Journal of Animal Science* **73** 3719–3731.
- Taylor PD, Hillier SG & Fraser HM 2004 Effects of GnRH antagonist treatment on follicular development and angiogenesis in the primate ovary. *Journal of Endocrinology* **183** 1–17. (doi:10.1677/joe.1.05685)
- Telfer EE, McLaughlin M, Ding C & Thong KJ 2008 A two-step serum-free culture system supports development of human oocytes from primordial follicles in the presence of activin. *Human Reproduction* **23** 1151–1158. (doi:10.1093/humrep/den070)
- Thomas FH, Armstrong DG & Telfer EE 2003 Activin promotes oocyte development in ovine preantral follicles *in vitro*. *Reproductive Biology and Endocrinology* **1** 76. (doi:10.1186/1477-7827-1-76)
- Tilly JL, Billig H, Kowalski KI & Hsueh AJ 1992 Epidermal growth factor and basic fibroblast growth factor suppress the spontaneous onset of apoptosis in cultured rat ovarian granulosa cells and follicles by a tyrosine kinase-dependent mechanism. *Molecular Endocrinology* **6** 1942–1950. (doi:10.1210/me.6.11.1942)
- Tomic D, Brodie SG, Deng C, Hickey RJ, Babus JK, Malkas LH & Flaws JA 2002 Smad 3 may regulate follicular growth in the mouse ovary. *Biology of Reproduction* **66** 917–923. (doi:10.1095/biolreprod66.4.917)
- Vallier L, Reynolds D & Pedersen RA 2004 Nodal inhibits differentiation of human embryonic stem cells along the neuroectodermal default pathway. *Developmental Biology* **275** 403–421. (doi:10.1016/j.ydbio.2004.08.031)
- Vallier L, Alexander M & Pedersen RA 2005 Activin/nodal GF pathways cooperate to maintain pluripotency of human embryonic stem cells. *Journal of Cell Science* **118** 4495–4509. (doi:10.1242/jcs.02553)
- Vallier L, Mendjan S, Brown S, Chng Z, Teo A, Smithers LE, Trotter MW, Cho CH, Martinez A, Rugg-Gunn P *et al.* 2009 Activin/nodal signalling maintains pluripotency by controlling Nanog expression. *Development* **136** 1339–1349. (doi:10.1242/dev.033951)
- Van Wezel IL, Dharmarajan AM, Lavranos TC & Rodgers RJ 1999 Evidence for alternative pathways of granulosa cell death in healthy and slightly atretic bovine antral follicles. *Endocrinology* **140** 2602–2612. (doi:10.1210/en.140.6.2602)
- Vernon RK & Spicer LJ 1994 Effects of basic fibroblast growth factor and heparin on follicle-stimulating hormone-induced steroidogenesis by bovine granulosa cells. *Journal of Animal Science* **72** 2696–2702.
- Walters DL & Schallenberger E 1984 Pulsatile secretion of gonadotrophins, ovarian steroids and ovarian oxytocin during the periovarian phase of the oestrous cycle in the cow. *Journal of Reproduction and Fertility* **71** 503–512. (doi:10.1530/jrf.0.0710503)
- Wandji SA, Pelletier G & Sirard MA 1992 Ontogeny and cellular localization of ¹²⁵I-labeled insulin-like growth factor-I, ¹²⁵I-labeled follicle-stimulating hormone, and ¹²⁵I-labeled human chorionic gonadotropin binding sites in ovaries from bovine fetuses and neonatal calves. *Biology of Reproduction* **47** 814–822. (doi:10.1095/biolreprod47.5.814)
- Wang J & Roy SK 2004 Growth differentiation factor-9 and stem cell factor promote primordial follicle formation in the hamster: modulation by follicle-stimulating hormone. *Biology of Reproduction* **70** 577–585. (doi:10.1095/biolreprod.103.023234)
- Wang H, Jiang JY, Zhu C, Peng C & Tsang BK 2006 Role and regulation of nodal/activin receptor-like kinase 7 signaling pathway in the control of ovarian follicular atresia. *Molecular Endocrinology* **20** 2469–2482. (doi:10.1210/me.2005-0446)
- Watson ED, Thomson SR & Howie AF 2000 Detection of steroidogenic acute regulatory protein in equine ovaries. *Journal of Reproduction and Fertility* **119** 187–192. (doi:10.1530/reprod/119.2.187)
- Welt C, Sidis Y, Keutmann H & Schneyer A 2002 Activins, inhibins, and follistatins: from endocrinology to signaling. A paradigm for the new millennium. *Experimental Biology and Medicine* **227** 724–752.
- van Wezel IL, Umaphysivam K, Tilley WD & Rodgers RJ 1995 Immunohistochemical localization of basic fibroblast growth factor in bovine ovarian follicles. *Molecular and Cellular Endocrinology* **115** 133–140. (doi:10.1016/0303-7207(95)03678-4)
- Wiater E & Vale W 2003 Inhibin is an antagonist of bone morphogenetic protein signaling. *Journal of Biological Chemistry* **278** 7934–7941. (doi:10.1074/jbc.M209710200)
- Wijgerde M, Ooms M, Hoogerbrugge JW & Grootegoed JA 2005 Hedgehog signaling in mouse ovary: Indian hedgehog and desert hedgehog from granulosa cells induce target gene expression in developing theca cells. *Endocrinology* **146** 3558–3566. (doi:10.1210/en.2005-0311)
- Wood JR & Strauss JF III 2002 Multiple signal transduction pathways regulate ovarian steroidogenesis. *Reviews in Endocrine & Metabolic Disorders* **3** 33–46. (doi:10.1023/A:1012748718150)
- Wrathall JH & Knight PG 1995 Effects of inhibin-related peptides and oestradiol on androstenedione and progesterone secretion by bovine theca cells *in vitro*. *Journal of Endocrinology* **145** 491–500. (doi:10.1677/joe.0.1450491)
- Wu X, Chen L, Brown CA, Yan C & Matzuk MM 2004 Interrelationship of growth differentiation factor 9 and inhibin in early folliculogenesis and ovarian tumorigenesis in mice. *Molecular Endocrinology* **18** 1509–1519. (doi:10.1210/me.2003-0399)
- Wulff C, Wiegand SJ, Saunders PT, Scobie GA & Fraser HM 2001a Angiogenesis during follicular development in the primate and its inhibition by treatment with truncated Flt-1-Fc (vascular endothelial growth factor Trap(A40)). *Endocrinology* **142** 3244–3254. (doi:10.1210/en.142.7.3244)
- Wulff C, Wilson H, Rudge JS, Wiegand SJ, Lunn SF & Fraser HM 2001b Luteal angiogenesis: prevention and intervention by treatment with vascular endothelial growth factor trap(A40). *Journal of Clinical Endocrinology and Metabolism* **86** 3377–3386. (doi:10.1210/jc.86.7.3377)
- Wulff C, Wilson H, Wiegand SJ, Rudge JS & Fraser HM 2002 Prevention of thecal angiogenesis, antral follicular growth, and ovulation in the primate by treatment with vascular endothelial growth factor Trap R1R2. *Endocrinology* **143** 2797–2807. (doi:10.1210/en.143.7.2797)

- Xia Y & Schneyer AL 2009 The biology of activin: recent advances in structure, regulation and function. *Journal of Endocrinology* **202** 1–12. (doi:10.1677/JOE-08-0549)
- Xu RH, Peck RM, Li DS, Feng X, Ludwig T & Thomson JA 2005 Basic FGF and suppression of BMP signaling sustain undifferentiated proliferation of human ES cells. *Nature Methods* **2** 185–190. (doi:10.1038/nmeth744)
- Xu WW, Kong XB, An LG & Zhang C 2009 Relationship between SnoN expression and mouse follicular development, atresia, and luteinization. *Zoological Science* **26** 66–73. (doi:10.2108/zsj.26.66)
- Yamamoto S, Konishi I, Nanbu K, Komatsu T, Mandai M, Kuroda H, Matsushita K & Mori T 1997 Immunohistochemical localization of basic fibroblast growth factor (bFGF) during folliculogenesis in the human ovary. *Gynecological Endocrinology* **11** 223–230. (doi:10.3109/09513599709152538)
- Yamamoto N, Christenson LK, McAllister JM & Strauss JF III 2002 Growth differentiation factor-9 inhibits 3′5′-adenosine monophosphate-stimulated steroidogenesis in human granulosa and theca cells. *Journal of Clinical Endocrinology and Metabolism* **87** 2849–2856. (doi:10.1210/jc.87.6.2849)
- Ying QL, Nichols J, Chambers I & Smith A 2003 BMP induction of Id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell* **115** 281–292. (doi:10.1016/S0092-8674(03)00847-X)
- Zhao J, Taverne MA, van der Weijden GC, Bevers MM & van den Hurk R 2001 Effect of activin A on *in vitro* development of rat preantral follicles and localization of activin A and activin receptor II. *Biology of Reproduction* **65** 967–977. (doi:10.1095/biolreprod65.3.967)
- Zsebo KM, Wypych J, McNiece IK, Lu HS, Smith KA, Karkare SB, Sachdev RK, Yuschenko VN, Birkett NC, Williams LR *et al.* 1990 Identification, purification, and biological characterization of hematopoietic stem cell factor from buffalo rat liver – conditioned medium. *Cell* **63** 195–201. (doi:10.1016/0092-8674(90)90300-4)

Received 16 February 2010

First decision 31 March 2010

Revised manuscript received 6 July 2010

Accepted 13 July 2010