The cholinergic system in rat testis is of non-neuronal origin

S U Schirmer^{1,2}, I Eckhardt², H Lau³, J Klein³, Y C DeGraaf¹, K S Lips^{2,4}, C Pineau⁵, I L Gibbins¹, W Kummer², A Meinhardt² and R V Haberberger¹

¹Anatomy and Histology, FMST, Flinders University, GPO Box 2100, 5001 Adelaide, South Australia, Australia,

²Department of Anatomy and Cell Biology, Justus-Liebig University, Aulweg 123, 35385 Giessen, Germany,

³Department of Pharmacology, College of Pharmacy, Goethe University, Max-von-Laue Strasse 9, 60438 Frankfurt, Germany, ⁴Laboratory for Experimental Trauma Surgery, Justus-Liebig University, Kerkrader Strasse 9, 35394 Giessen, Germany and ⁵Institut National de la Santé et de la Recherche Médicale, U625, Groupe d'étude de la reproduction chez l'homme et les mammifères, Institut Fédératif de Recherche 140, Campus de Beaulieu, 35042 Rennes, France

Correspondence should be addressed to R V Haberberger; Email: rainer.haberberger@flinders.edu.au

Abstract

The cholinergic system consists of acetylcholine (ACh), its synthesising enzyme, choline acetyltransferase (CHAT), transporters such as the high-affinity choline transporter (SLC5A7; also known as ChT1), vesicular ACh transporter (SLC18A3; also known as VAcHT), organic cation transporters (SLC22s; also known as OCTs), the nicotinic ACh receptors (CHRN; also known as nAChR) and muscarinic ACh receptors. The cholinergic system is not restricted to neurons but plays an important role in the structure and function of non-neuronal tissues such as epithelia and the immune system. Using molecular and immunohistochemical techniques, we show in this study that non-neuronal cells in the parenchyma of rat testis express mRNAs for *Chat*, *Slc18a3*, *Slc5a7* and *Slc22a2* as well as for the CHRN subunits in locations completely lacking any form of innervation, as demonstrated by the absence of protein gene product 9.5 labelling. We found differentially expressed mRNAs for eight α and three β subunits of CHRN in testis. Expression of the $\alpha 7$ -subunit of CHRN was widespread in spermatogonia, spermatocytes within seminiferous tubules as well as within Sertoli cells. Spermatogonia and spermatocytes also expressed the $\alpha 4$ -subunit of CHRN. The presence of ACh in testicular parenchyma (TP), capsule and isolated germ cells could be demonstrated by HPLC. Taken together, our results reveal the presence of a non-neuronal cholinergic system in rat TP suggesting a potentially important role for non-neuronal ACh and its receptors in germ cell differentiation.

Reproduction (2011) **142** 157–166

Introduction

The cholinergic system consists of acetylcholine (ACh), its synthesising enzymes, transporters and receptors. ACh is synthesised by choline acetyltransferase (CHAT). Ongoing ACh synthesis requires the uptake of choline into cholinergic cells via a high-affinity choline transporter (SLC5A7; also known as ChT1; Okuda *et al.* 2000). In neurons, ACh is transported into synaptic vesicles via the vesicular ACh transporter (SLC18A3; also known as VAcHT; Erickson *et al.* 1996, Parsons 2000) and released via exocytosis upon stimulation. In addition, ACh release can occur through bidirectional transport of ACh, utilising organic cation transporters (SLC22s; also known as OCTs) with SLC22A2 being the most likely candidate for non-neuronal transport (Wessler *et al.* 2001, Lips *et al.* 2005). ACh can stimulate five different G-protein-coupled muscarinic receptor subtypes (CHRM, also known as mAChR; Nathanson 2008) and an unknown variety of ionotropic nicotinic ACh receptors (CHRN; also known as nAChR) with

subtype-specific arrangements of nine α - and four β -subunits in mammals (Albuquerque *et al.* 2009). It is now known that the cholinergic system, traditionally associated with neurotransmission, is not restricted to neurons but plays an important role in the structure and function of non-neuronal tissues such as epithelia and the immune system (Fujii *et al.* 2008, Kummer *et al.* 2008, Wessler & Kirkpatrick 2008). For example, in epithelia, CHRNs are involved both in maintaining the integrity of the epithelial layer and in the development of neoplastic changes (Grando *et al.* 2003, Grozio *et al.* 2007, Paleari *et al.* 2009).

Several observations indicate the presence of a cholinergic system within the mammalian testis: functional ACh receptors are found on male germ cells and Sertoli cells; mice lacking CHRN subtypes or with reduced ACh levels reveal reduced sperm motility (Borges *et al.* 2001, Bray *et al.* 2005); and high nicotine levels in the blood lead to reduced sperm production and fertility (Dwivedi & Long 1989, Yamamoto *et al.* 1998). Published reports suggest that there is little, if any,

Figure 1 (A) Innervation within the testicular parenchyma (TP) was examined with an antibody against protein gene product 9.5 (PGP 9.5, red). α -smooth muscle actin antibody labels peritubular myoid cells and interstitial blood vessels (green). Nerve fibres were absent in rat TP, but PGP 9.5 appears to stain a spermatogonial subpopulation. Bar, 75 μ m. (B) The mesostructures showed a dense innervation with PGP 9.5 positive nerve fibres and nerve fibre bundles. Bar, 100 μ m. (C) Box plot showing the relative mRNA expression levels of choline acetyltransferase (*Chat*), high-affinity choline transporter (*Slc5a7*), vesicular ACh transporter (*Slc18a3*) and organic cation transporter 2 (*Slc22a2*) in TP. The ΔC_T values were subtracted from 50 showing higher values with higher expression.

cholinergic innervation of most testicular tissue suggesting the presence of a non-neuronal cholinergic system. However, neither the sites of testicular ACh synthesis nor the cells targeted by locally synthesised ACh are known. Therefore, we have used molecular and immunohistochemical techniques to determine the levels of ACh, and the expression patterns of CHAT, SLC5A7, SLC18A3, SLC22A2 and CHRN subunits in rat testicular tissue.

Results

The synthesising enzyme and the transporters of the cholinergic system are expressed in testicular parenchyma

No protein gene product (PGP) 9.5 positive nerve fibres could be detected in the parenchyma of rat testes, whereas non-neuronal basal cells of seminiferous tubules showed PGP 9.5 immunoreactivity (IR; Fig. 1). In contrast to the parenchyma, the mesostructures that connect testis with epididymis and ductus deferens were densely supplied with PGP 9.5 positive nerve fibres (Fig. 1). However, PCR revealed significant expression of mRNA for *Chat*, *Slc5a7*, *Slc18a3* and *Slc22a2* in testicular parenchyma (TP). After removal of the testicular capsule (TC), the relative mRNA expression profiles in the parenchyma had a rank order of *Chat* > *Slc5a7* > *Slc22a2* > *Slc18a3* (Fig. 1 and Table 1).

We further determined the expression of mRNAs for cholinergic elements in isolated populations of testicular cells. *Chat* mRNA was localised in cells of the seminiferous tubules (Fig. 2). High or moderate expression levels of mRNA for *Chat*, *Slc5a7*, *Slc18a3* and *Slc22a2* occurred in pachytene spermatocytes, round spermatids and, except for *Slc18a3*, in residual bodies (Fig. 2). In contrast, no product corresponding to *Chat*, *Slc5a7*, *Slc18a3* or *Slc22a2* mRNA could be

detected in Sertoli cells, peritubular cells and spermatogonia. Products corresponding to *Chat* and *Slc5a7* were present in total RNA from isolated testicular macrophages (TM). A PCR product corresponding to *Chat* mRNA could be detected in Leydig cells (Fig. 2).

CHAT and SLC5A7 proteins are present in seminiferous tubules

IR for CHAT, SLC5A7, SLC22A2 and SLC18A3 was detected mainly in the seminiferous epithelium. CHAT-IR was detected also in endothelial cells of small interstitial arteries. The labelling was absent after pre-absorption with the corresponding antigen. Strong IR for CHAT and SLC5A7 was detected in spermatids and in some basal cells of the seminiferous tubules, presumably spermatogonia (Fig. 3). Similar cells were also positive for SLC18A3 (Fig. 3), while only barely detectable SLC18A3 labelling was present in

Table 1 Relative expression levels for CHRN subunits, and *Chat*, *Slc5a7*, *Slc18a3* and *Slc22a2* in testicular parenchyma.

CHRN subunit	Samples (n)	Testicular parenchyma (mean $\pm$ s.d.)
$\alpha 1$	3	30.1 $\pm$ 0.17
$\alpha 2$	5	29.2 $\pm$ 0.54
$\alpha 3$	5	28.9 $\pm$ 0.92
$\alpha 4$	5	39.8 $\pm$ 0.19
$\alpha 5$	3	33.5 $\pm$ 0.42
$\alpha 6$	5	8.7 $\pm$ 11.94*
$\alpha 7$	3	34.6 $\pm$ 0.55
$\alpha 9$	5	32.9 $\pm$ 0.39
$\alpha 10$	3	29.0 $\pm$ 0.94
$\beta 1$	3	34.8 $\pm$ 0.40
$\beta 2$	3	35.8 $\pm$ 0.68
$\beta 3$	3	33.9 $\pm$ 0.19
$\beta 4$	5	0.0
<i>Slc22a2</i>	3	31.8 $\pm$ 0.28
<i>Chat</i>	3	37.2 $\pm$ 0.81
<i>Slc5a7</i>	7	35.0 $\pm$ 0.30
<i>Slc18a3</i>	3	27.7 $\pm$ 0.81

The asterisk indicates the absence of mRNA in three out of five replicates, while the other two replicates revealed very weak expression.

Figure 2 Ethidium bromide-stained agarose gels showing the PCR products for choline acetyltransferase (*Chat*, product length 183 bp), high-affinity choline transporter (*Slc5a7*, product length 150 bp), vesicular ACh transporter (*Slc18a3*, product length 149 bp) and organic cation transporter 2 (*Slc22a2*, product length 162 bp) in isolated cell populations as examined by RT-PCR. *Gapdh* (product length 320 bp) served as the reference gene. +, rat brain (positive control); -, water control (negative control) and M, marker.

other cell types of the seminiferous epithelium (Fig. 3). SLC22A2 IR was visible in Leydig cell clustered in the interstitium, but not within the seminiferous tubules (Fig. 3).

Multiple *CHRN* subunits are expressed in different cell types of the TP

To determine potential targets of ACh in the testis, we first analysed the relative mRNA expression patterns of *CHRN* subunits in TP. In the parenchyma, mRNA for 11 out of 13 *CHRN* subunits, including the supposedly 'muscle specific' $\alpha 1$ - and $\beta 1$ -subunits, was detected (Fig. 4 and Table 1). The mRNAs for *CHRN* $\alpha 6$ - and $\beta 4$ -subunits

could not be detected in the TP but were present in the TC (Fig. 4). The highest levels of expression were observed for mRNA for $\alpha 4$ -subunit followed by $\alpha 7$ -, $\alpha 5$ - and $\alpha 9$ -subunits and β -subunits 1–3 (Fig. 4 and Table 1).

The mRNAs for the $\alpha 7$ - and $\alpha 4$ -subunits of *CHRN* are expressed in the seminiferous tubule

As mRNAs for $\alpha 4$ - and $\alpha 7$ -subunits of *CHRN* were strongly expressed in the testis and $\alpha 7$ -subunit of *CHRN* has been reported to be associated with sperm motility (Bray *et al.* 2005), we investigated the mRNA expression patterns of both subunits in isolated cell preparations. PCR products corresponding to mRNAs of both subunits were present in peritubular and Leydig cells, whereas Sertoli cells contained mRNA for the $\alpha 7$ -, but not the $\alpha 4$ -subunit of *CHRN*. Spermatogonia as well as pachytene spermatocytes expressed mRNAs for $\alpha 4$ - and $\alpha 7$ -subunits; however, mRNA for $\alpha 7$ -subunit could not be detected in later stage spermatocytes. The PCR product corresponding to mRNA for the $\alpha 7$ -subunit was found in residual bodies, but not in round spermatids (Fig. 5). TM expressed mRNAs for both $\alpha 4$ - and $\alpha 7$ -subunits (Fig. 5). Since subtype-specific *CHRN* antisera were not available (Herber *et al.* 2004, Moser *et al.* 2007), it was not possible to further analyse the distribution of receptor subunit proteins. However, *in situ* hybridisation localised mRNA for the $\alpha 7$ -subunit of *CHRN* in pachytene spermatocytes (Fig. 6).

ACh is present in testis

ACh was measured in rat testis. Comparison of the ACh content in 16-day-old immature testes as well as in adult testes showed that ACh was present in all

Figure 3 Multiple labelling immunohistochemistry. (A, B and D) Choline acetyltransferase (CHAT, in red), (C and E) high-affinity choline transporter (SLC5A7, in red), (F) vesicular acetylcholine transporter (SLC18A3, in red), (G) organic cation transporter 2 (SLC22A2). CHAT and SLC5A7 are present in spermatids (A–C), with CHAT-antiserum additionally staining round spermatids and some meiotic cells (D), while SLC18A3 is present in spermatogonia (F, inset). SLC22A2 is present in interstitial cells (G). α -Smooth muscle actin was visualised with FITC-coupled antibody (green in A–G). Bars A, 100 μ m; B, C, F and G, 75 μ m; D and E, 20 μ m.

Figure 4 The mRNA expression level of the nicotinic ACh receptor (CHRN) subunits, α (A) and β (B), in testicular parenchyma (TP) was determined by RT-qPCR. The asterisk indicates the absence of mRNA in three out of five replicates, while the other two replicates revealed very weak expression. #, no mRNA detectable. (C and D) Comparison of the $\alpha 6$ - and $\beta 4$ -subunits of CHRN expression in testicular capsule (TC) and testicular parenchyma (TP). The ΔC_t values were subtracted from 50 showing higher values with higher expression.

samples. It was found at lower levels in TC and parenchyma of 16-day-old animals, whereas adult rats showed extremely high ACh levels in the parenchyma (3068.3 ± 764.94 vs 3.8 ± 1.62 pmol/mg protein, Table 2 and Fig. 7) and the ACh levels were similar in adults and 16-day-old rats in the capsule. Freshly isolated germ cells of adult rats also contained ACh; however, significantly higher levels were present in the supernatant (11.24 ± 3.65 vs 47.74 ± 23.79 pmol/mg protein, Table 2 and Fig. 7).

Discussion

ACh synthesising enzyme and transporters are present in TP

This study demonstrates that non-neuronal cells in the parenchyma of rat testis have all the components necessary for cholinergic signalling. We detected mRNAs for the ACh synthesising enzyme, transporters and ACh receptors in locations completely lacking any form of innervation as was demonstrated by the absence of PGP 9.5 labelling (see also Zhu *et al.* (1995)).

Cells of the germinal epithelium can synthesise ACh

Previous studies have demonstrated *Chat* mRNA and activity in human and rat spermatozoa (Sastry *et al.* 1981, Ibanez *et al.* 1991), while the ACh degrading enzyme ACh esterase, AChE, is expressed in rat spermatozoa with higher enzymatic activity in testis than in epididymis (Egbunike 1980). We now have found CHAT and SLC5A7 mRNAs and proteins in isolated germ cells *in vitro* as well as *in situ*. In addition, mRNAs for two cholinergic transporters, *Slc18a3* and *Slc22a2*, were present in isolated spermatocytes and spermatids. SLC5A7 is the rate-limiting factor in ACh synthesis (Ribeiro *et al.* 2006), whereas SLC18A3 and SLC22A2 are involved in vesicular and non-vesicular ACh release respectively (Erickson *et al.* 1996, Ribeiro *et al.* 2006). Taken together, these observations strongly support the possibility that the germinal epithelium has the capability to synthesise and release ACh. However, it cannot be excluded that the testicular ACh synthesising machinery is not functional. Lonnerberg & Ibanez (1999) described truncated non-functional forms of CHAT in rat testis. Our expression analysis demonstrated the presence of mRNA and protein for ACh synthesis, and data are supported by the detection of ACh in rat testis and in isolated germ cells. ACh was present in isolated cells at about 11 pmol/mg protein and at higher levels of about 48 pmol/mg protein in the supernatant. The levels are comparable to the ACh content determined in non-neuronal cells. ACh concentrations in mononuclear cells were 0.35 pmol/1 million cells (Fujii & Kawashima 2001), 1.21 pmol/1 million cells (Innis 2011) and 0.83 pmol/1 million cells (Hecker *et al.*

Figure 5 The presence of nicotinic ACh receptor (CHRN) subunits $\alpha 4$ (product length 137 bp) and $\alpha 7$ (product length 235 bp) was determined in isolated testicular cell populations using RT-PCR. *Gapdh* (product length 320 bp) served as the reference gene. +, positive control rat brain; –, negative water control; M, marker.

2009). In the absence of nerve fibres, our results clearly indicate a local non-neuronal source of ACh in rat testis. The two interesting observations in this study are the high ACh levels in the supernatant of germ cells and the strong increase in ACh levels in adult rat testis. The high level in the supernatant of freshly isolated germ cells may be related to constant ACh synthesis and release from the germ cells itself. This is supported by the fact that the molecule responsible for storage of ACh, SLC18A3, could not be detected in the germ cell population. This is similar to human and murine leukocytes and crypt cells in the distal rat colon, which also do not express SLC18A3, but contains or synthesises ACh (Kawashima & Fujii 2000, Yajima *et al.* 2010). In addition, vascular endothelial cells, TM and Sertoli cells could also be a potential source of endogenous ACh. Endothelial cells have been shown to contain the machinery responsible for ACh and release ACh (Haberberger *et al.* 2000, Kirkpatrick *et al.* 2003), and macrophages and Sertoli cells express *Chat* and *Slc5a7* mRNA. These cells could synthesise and release ACh in the absence of a functional ACh synthesis in germ cells.

Testicular cells can be targets for non-neuronal ACh

The TP contains a large variety of potential targets for non-neuronally released ACh. We found the mRNA for eight α and three β -subunits of CHRN in testicular cells, including mRNAs for the supposedly 'muscle specific' $\alpha 1$ - and $\beta 1$ -subunits. These subunits are also expressed in human skin, where their function remains elusive (Spies *et al.* 2006). Expression of the $\alpha 7$ -subunit of CHRN was widespread in spermatogonia, spermatocytes within seminiferous tubules as well as within Sertoli cells. Spermatogonia and spermatocytes also expressed the $\alpha 4$ -subunit of CHRN, potentially as part of the heteropentameric $\alpha 4\beta 2$ CHRN, which is one of the

major CHRN subtypes in neuronal tissue (Albuquerque *et al.* 2009).

The adverse effects of smoking on reproductive function such as preterm delivery and abortion are well established. In the male, nicotine, a major component of cigarette smoke, induced impairment of spermatogenesis and steroidogenesis, the latter probably by affecting steroidogenic acute regulatory protein (StAR), the rate-limiting factor in sex steroid synthesis (Gocze & Freeman 2000, Bose *et al.* 2007). However, the identification of the mRNAs for at least the $\alpha 4$ - and $\alpha 7$ -subunits of CHRN in Leydig cells in this study proposes an additional mechanism of suppression of androgen production. Effects on spermatogenesis and sperm function parameters can be explained by the presence of $\alpha 3$ -, $\alpha 5$ -, $\alpha 7$ -, $\alpha 9$ - and $\beta 4$ -subunits of CHRN in ejaculated human spermatozoa, which are indicative of an earlier production during germ cell development in the testis, a finding that is partially supported by our mRNA expression data. Functionally, mouse sperm deficient of the $\alpha 7$ -subunit of CHRN shows impaired motility (Bray *et al.* 2005) and the ACh-triggered acrosome

Figure 6 Localisation of $\alpha 7$ -subunit of CHRN mRNA expression was examined in rat testis using *in situ* hybridisation. (A) Sense and (B–D) antisense probes. Staining for the $\alpha 7$ -subunit of CHRN mRNA was mainly visible in primary spermatocytes, with only few round spermatids weakly labelled. Interstitial cells were negative. No staining was obtained using the sense probe (A). Bars, 20 μ m.

Table 2 Acetylcholine content in germ cell preparations and testicular capsule (TC) and testicular parenchyma (TP).

Source	Samples (n)	Ratio (%) ACh/choline	Protein ($\mu\text{g/ml}$)	ACh (pmol/mg protein)
Germ cells	5	67	425.00	11.24 ± 3.65
Germ cells supernatant	6	200	841.67	47.74 ± 23.79
TC d16	6	36	666.67	6.33 ± 2.41
TP d16	6	08	1662.50	3.61 ± 1.32
TC adult	2	18	5850.00	3.8 ± 1.62
TP adult	2	45	212.50	3068.3 ± 764.94

reaction was suppressed by antagonists of tyrosine phosphorylation (Kumar & Meizel 2005). Other CHRN subunits seem to have no direct impact on fertility since mice with deficiency in the subunits $\alpha 4$, $\beta 2$ (Marubio *et al.* 1999), $\alpha 5$ (Wang *et al.* 2002) and $\beta 4$ (Wang *et al.* 2003) were reported with no abnormalities with respect to litter size and fertility. As spermatozoa are exclusively transported in the luminal compartment of the male and female reproductive tract separated by epithelial cells from nerve endings that could release ACh, the studies of Bray *et al.* (2005) and Kumar & Meizel (2005) are a clear indication for the necessity of a non-neuronal cholinergic system in the male gonad and during fertilisation.

Albeit information from toxicological studies and transgenic mouse models have provided some insight in the pathophysiological functions of the cholinergic system, its role in the normal testis is still unclear. Observations from other systems suggest that CHRN can mediate effects on cell division, metabolism and motility also in testicular tissue. The proximity of cellular ACh synthesising and reception sites in the testis favours an autocrine or paracrine mode of regulating testicular function by the non-neuronal cholinergic system, comparable to what is observed in the lymphatic system and the skin (Kawashima & Fujii 2000, Kurzen *et al.* 2007). Taken together, our results are a clear indication for a functional non-neuronal cholinergic system in the testis and add important data to further understand the pathophysiological consequences of smoking on male reproductive function.

Materials and Methods

Tissue preparation

Specimens for RT-PCR were obtained from 8- to 10-week-old male Wistar-Furth rats. The animals were killed by a lethal dose of isoflurane, and the testis was removed and decapsulated. Approximately 30 mg TP and the complete capsule were snap-frozen separately in liquid nitrogen and stored at -80°C until required. For *in situ* hybridisation, the testis was snap frozen in liquid isopentane and fixed as required after sectioning. For immunohistochemistry, TP and capsule were immersion fixed in Zamboni's fixative (2% formaldehyde, 15% picric acid in 0.2 M phosphate buffer, pH 7), washed overnight in PBS,

cryoprotected in 18% sucrose and frozen in optimal cutting temperature compound. All animal procedures were performed in accordance with approval from the Flinders University Animal Welfare Committee.

Isolation of cells from the rat testis

Leydig cells, Sertoli cells, spermatogonia, pachytene spermatocytes, round spermatids and residual bodies were prepared from rat testes as described previously (Guillaume *et al.* 2001). TM were isolated from testes without any enzymatic treatment. The testes were decapsulated into 10 ml pre-warmed endotoxin-free DMEM-F12 medium (PAA Laboratories, Morningside, QLD, Australia). The seminiferous tubules were gently separated, and the volume was adjusted to 50 ml. After gentle stirring, the tubule fragments were allowed to settle for 5 min before the supernatant was centrifuged at 300 *g* for 10 min. The pellet containing interstitial cells and TM was resuspended in 5 ml DMEM-F12 and incubated at 32°C and 5% CO_2 for 30 min. TM and the remaining interstitial cells were separated by the more rapid adherence of TM to surfaces.

Figure 7 Detection of ACh in rat germ cells, testicular capsule and testicular parenchyma (A–C). Comparison of the ACh content in 16-day-old immature and adult testes showed that ACh was present at low levels in testicular capsule and testicular parenchyma of 16-day-old animals (A), while adult rats showed very high ACh levels in the parenchyma and low levels in the capsule (B). The box plots in C show the presence of ACh in freshly isolated germ cells and in the supernatant.

After 30 min, non-adherent cells were removed by washing with fresh medium. Subsequently, TM were washed a second time, by pipetting directly on the surface. Purity of TM (80%) was determined by immunolabelling using monoclonal antibodies ED1 and ED2 (AbD Serotec, Oxford, UK) directed against monocytes/macrophages.

RT-PCR

RT-PCR was used to detect mRNAs in isolated populations of testicular cells (PTC 200; Peqlab, Erlangen, Germany). Quantitative real-time PCR (RT-qPCR) was used to quantify relative expression levels of mRNAs using cDNA from TP and capsule (Corbett Roto-cycler, Sydney, NSW, Australia). Capsule and parenchyma were lysed in RLT lysis buffer (Qiagen) using a tissue lyser (Qiagen). RNA was extracted (RNeasy Mini Kit,

Qiagen), and quantity and quality of the RNA were determined. Isolated testicular cells were lysed in RLT buffer, homogenised with a 21 G syringe and subsequently used for RNA extraction according to the manufacturer's instructions (RNeasy Mini Kit, Qiagen). Total RNA from brain served as positive control. In negative controls, the reverse transcriptase was replaced by water (negative water control). The extraction was followed by DNase digestion and RT (iScript, Bio-Rad). The cDNA was used for subsequent PCR and RT-qPCR. RT-PCR for isolated testicular cells was performed as for standard PCR in a 25 µl reaction volume with 1 µl cDNA, 0.2 mM dNTP mix, 1 mM MgCl₂, 0.2 pmol each primer (Table 3), buffer and GoTaq Polymerase according to the manufacturer's protocol (Promega). *Gapdh* was used as the reference gene. All RT-qPCR were performed in duplicate or triplicate from three to six animals using a ready-to-use kit according to the

Table 3 Primer pairs used for RT-PCR.

Primer	Sequence (5' → 3') F – forward and R – reverse	Product length (bp)	Accession no.
<i>Chat</i>	F TGAACGCCTGCCTCCATTTCGGCCTGCTGA R GTGCCATCTCGGCCACCACGAAGTGA	272	XM_224626
<i>Chat</i>	F CAACCATCTTCTGGCACTGA R TAGCAGGCTCAATAGCCATT	183	XM_224626
<i>Slc18a3</i>	F GCCACATCGTTCACCTCTCTTG R CGGTTTCATCAAGCAACACATC	149	X80395
<i>Slc22a2</i>	F GCCTCCTGATCCTGGCTG R GGTGTCAGGTTCTGAAGAGAG	226	X98334
<i>Slc5a7</i>	F ATGGCTCTACCAGCCATTG R GGACATGACAGCAGCAGAAA	189	AB030947.1
<i>Slc5a7</i>	F CAAGACCAAGGAGGAAGCAC R GCAAACATGGAACCTTGTCGA	150	AB030947.1
<i>Rpl19</i>	F CATGGAGCACATCCCAAAC R CCATAGCCTGGCCACTATGT	216	NM_031103.1
<i>18S rRNA</i>	F CCGCAGCTAGGAATAATGGA R AGTCGGCATCGTTTATGGTC	245	M11188
<i>Gapdh</i>	F CATTGTTGCCATCAACGACC R TCACACCCATCACAACATG	320	NM_017008
$\alpha 1$	F AACTTCATGGAGAGCGGAGA R CAGCTCCACAATGACGAGAA	285	NM_024485.1
$\alpha 2$	F GGAGCAGATGGAGAGGACAG R AGCACAGTGAGGACGAGAT	216	NM_133420.1
$\alpha 3$	F GCCAACCTCACAAGAAGCTC R CCAGGATGAAAACCCAGAGA	208	NM_052805.2
$\alpha 4$	F GGACCCTGGTGACTACGAGA R CATAGAACAGGTGGGCCTTG	137	NM_024354
$\alpha 5$	F CACGTCGTGAAAGAGAACGA R TCCCAATGATTGACACCAGA	112	NM_017078.2
$\alpha 6$	F ACAGCTCTCCACACGCTCT R GAAGTCACCGACGGCATTAT	286	NM_057184.1
$\alpha 7$	F GGCTCTGCTGGTATTCTTGC R AAACCATGCACACAGTTCA	286	NM_012832.3
$\alpha 7$	F ACATTGACGTTTCGCTGGTTC R CTACGGCGCATGGTTACTGT	235	L31619
$\alpha 9$	F CGTGGGATCGAGACCAAGTAT R AAAGGTCAGGTTGCACTGCT	242	NM_022930.1
$\alpha 10$	F CTGCTGACTCTGGGAGAAAG R GGCTGACTCTAGTGGCTTGG	317	NM_022639.1
$\beta 1$	F CATCGAGTCTCTCCGTGTCA R TGCAATTCTGCCAGTCAAAG	206	NM_012528.1
$\beta 2$	F AAGCCTGAGGACTTCGACAA R TGCCATCATAGGAGACCACA	142	NM_019297.1
$\beta 3$	F CACTCTGCGCTTGAAAGGAA R GCGGACCCATTTCTGGTAAC	196	NM_133597.1
$\beta 4$	F CTCTGAACAAAACCCGGTA R ACCTCAATCTTGACGGCACT	371	NM_052806.2

Table 4 Primer used for generation of *in situ* hybridisation probes. T7 binding site TAATACGACTCACTATAGGG; SP6 binding site ATTTAGGTGACACTATAGAA.

Primer	Sequence (5' → 3') F – forward, R – reverse	Product length (bp)
Chat_F	TAATACGACTCACTATAGGGTGAACGCCTGCCTCCATTGGCCTGCTGA	312
Chat_R	ATTTAGGTGACACTATAGAAAGTGCCATCTCGGCCACCAACGAAGTCA	
$\alpha 7$ _F	TAATACGACTCACTATAGGGGGCTCTGCTGGTATTCTTGC	326
$\alpha 7$ _R	ATTTAGGTGACACTATAGAAAACCATGCACACCAGTTCA	

manufacturer's protocol (iQ SYBR green Supermix, Bio-Rad). Primers specific for mRNA sequences were designed using Blast (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>) (Table 3). All primers spanned introns except the primers for the $\alpha 2$ -subunit and *Slc18a3*. All PCR products were sequenced (Flinders University Sequencing Facility) and showed 100% homology with the predicted target sequence. Primers specific for the rat reference genes *Rpl19* (Adams *et al.* 2007) and *18S* rRNA were used for normalisation. The RT-qPCR data were normalised by subtracting the threshold cycle (C_T) levels between the genes of interest and the mean of *Rpl19* and *18S* rRNA (Livak & Schmittgen 2001). The ΔC_T values were subtracted from 50 so that higher values reflect higher expression levels.

In situ hybridisation

In situ hybridisation probes were prepared using digoxigenin (DIG) labelling (Roche). The primers contained binding sites for either the T7 or SP6 RNA polymerase (Table 4). Cryosections (12 μ m) were fixed in 4% paraformaldehyde (PFA), permeabilised with proteinase K (2 μ g/ml) for 7 min, fixed again in PFA and acetylated (0.1 M triethanolamine containing 0.25% (v/v) acetic anhydride). After prehybridisation, hybridisation was performed overnight at 65 °C. Subsequently, the sections were washed in decreasing concentrations of SSC. The DIG-labelled probes were detected using alkaline phosphate-conjugated anti-DIG antibody and subsequent visualisation using NBT/BCIP as substrate (Roche). Colour development was allowed to proceed in the darkness for 4–16 h. The reaction was terminated by immersion in tap water.

Immunohistochemistry

Rat testes were serially cryosectioned at a thickness of 12 μ m, fixed with methanol and subsequently preincubated for 1 h with PBS (145.4 mM NaCl (8.5 g/l), 7.54 mM Na_2HPO_4 (1.07 g/l), 2.5 mM $\text{NaH}_2\text{PO}_4\text{H}_2\text{O}$ (0.39 g/l), pH 7.1) containing 10% normal donkey serum, 0.1% BSA and 0.5% Tween 20. Indirect immunofluorescence was performed by overnight incubation with antisera directed against PGP 9.5, CHAT, SLC5A7 or SLC18A3 in combination with an FITC-conjugated anti-smooth muscle actin antibody (anti-SMA-FITC; Table 5) diluted in PBS with doubled concentration NaCl at room temperature followed by washing in PBS and subsequent incubation with appropriate combinations of secondary reagents (Table 5) for 1 h at room temperature. After incubation with the secondary reagents, the slides

were washed in PBS and coverslipped in carbonate-buffered glycerol at pH 8.6 and evaluated by fluorescence microscopy or sequential scanning using a confocal laser scanning microscope (TCS SP5, Leica, Bensheim, Germany). Specificity of the CHAT, SLC18A3 and SLC22A2 antisera was tested by pre-absorption with the corresponding antigen. In addition, the specificity of the SLC5A7 antibody has been shown in murine cochlea, brain and spinal cord (SLC5A7, Brandon *et al.* 2004, Bergeron *et al.* 2005).

ACh measurement

Freshly prepared tissue from testes was homogenised in ten volumes of a mixture of ice-cold acetone (85%) and 1 M formic acid (15%) as described previously (Klein *et al.* 1993). Cultured cells and cell media (supernatant) were taken in 96% ethanol. Homogenates were centrifuged at 10 000 *g*. Aliquots of the supernatant were taken to dryness in a vacuum centrifuge, taken in HPLC buffer and injected into a Eicom HTEC-500 microbore system coupled to a Shimadzu SIL-20AC auto-sampler. The buffer composition was 5 g KHCO_3 , 400 mg sodium decanesulfonate, and 50 mg EDTA in 1 l Aqua Dest. (pH 8.3). At a flow rate of 0.15 ml/7 min, retention times were 7.8 min for choline and 13.2 min for ACh. The limit of detection was 1–2 fmol analyte per 5 μ l injection volume. ACh chloride and choline chloride (purity >99% each) were purchased from Sigma-Aldrich.

Table 5 Primary and secondary antisera, nuclear staining.

Antibody	Host	Conjugate	Supplier	Dilution
Primary				
Anti-rat-SLC5A7	Polyclonal, rabbit		Chemicon	1:100
Anti-rat-CHAT	Polyclonal, sheep		Chemicon	1:2000
Anti-rat-SLC18A3	Polyclonal, goat		Chemicon	1:800
Anti-SMA	Monoclonal, mouse	FITC	Sigma	1:800
Anti-rat-PGP 9.5	Polyclonal, rabbit		Neuromics	1:500
Secondary				
Anti-rabbit-Ig	Donkey	Cy3	Jackson	1:100
Anti-sheep/goat-Ig	Donkey	Cy3	Jackson	1:100
Hoechst 333258			Mol. Probes	1:2000

Chemicon, Boronia, Australia; Neuromics, Medina, MN, USA; Jackson Immuno Research, West Grove, PA, USA; Molecular Probes (Invitrogen), Mulgrave, VIC, Australia.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by the Else Kröner-Fresenius-Stiftung and the Flinders University Mary Overton Neuroscience Fellowship.

Acknowledgements

The grant support of the Else Kröner-Fresenius-Stiftung and a travel grant of the DAAD are gratefully acknowledged. The helpful assistance of Sudhanshu Bhushan, Eva Schneider und Yongning Lu is gratefully acknowledged.

References

- Adams V, Linke A, Wisloff U, Döring C, Erbs S, Krankel N, Witt CC, Labeit S, Müller-Werdan U, Schuler G *et al.* 2007 Myocardial expression of *Murf-1* and *MAFbx* after induction of chronic heart failure: effect on myocardial contractility. *Cardiovascular Research* **73** 120–129. (doi:10.1016/j.cardiores.2006.10.026)
- Albuquerque EX, Pereira EF, Alkondon M & Rogers SW 2009 Mammalian nicotinic acetylcholine receptors: from structure to function. *Physiological Reviews* **89** 73–120. (doi:10.1152/physrev.00015.2008)
- Bergeron AL, Schrader A, Yang D, Osman AA & Simmons DD 2005 The final stage of cholinergic differentiation occurs below inner hair cells during development of the rodent cochlea. *Journal of the Association for Research in Otolaryngology* **6** 401–415. (doi:10.1007/s10162-005-0018-3)
- Borges MO, Abreu ML, Porto CS & Avellar MC 2001 Characterization of muscarinic acetylcholine receptor in rat Sertoli cells. *Endocrinology* **142** 4701–4710. (doi:10.1210/en.142.11.4701)
- Bose M, Debnath D, Chen Y & Bose HS 2007 Folding, activity and import of steroidogenic acute regulatory protein into mitochondria changed by nicotine exposure. *Journal of Molecular Endocrinology* **39** 67–79. (doi:10.1677/JME-07-0051)
- Brandon EP, Mellott T, Pizzo DP, Coufal N, D'Amour KA, Gobeske K, Lortie M, Lopez-Coviella I, Berse B, Gage FH *et al.* 2004 Choline transporter 1 maintains cholinergic function in choline acetyltransferase haploinsufficiency. *Journal of Neuroscience* **24** 5459–5466. (doi:10.1523/JNEUROSCI.1106-04.2004)
- Bray C, Son JH, Kumar P & Meizel S 2005 Mice deficient in *CHRNA7*, a subunit of the nicotinic acetylcholine receptor, produce sperm with impaired motility. *Biology of Reproduction* **73** 807–814. (doi:10.1095/biolreprod.105.042184)
- Dwivedi C & Long NJ 1989 Effect of cholinergic agents on human spermatozoa motility. *Biochemical Medicine and Metabolic Biology* **42** 66–70. (doi:10.1016/0885-4505(89)90042-X)
- Egbunike GN 1980 Changes in acetylcholinesterase activity of mammalian spermatozoa during maturation. *International Journal of Andrology* **3** 459–468. (doi:10.1111/j.1365-2605.1980.tb00134.x)
- Erickson JD, Weihe E, Schafer MK, Neale E, Williamson L, Bonner TI, Tao-Cheng JH & Eiden LE 1996 The *VACHT/ChAT*, "cholinergic gene locus": new aspects of genetic and vesicular regulation of cholinergic function. *Progress in Brain Research* **109** 69–82. (doi:10.1016/S0079-6123(08)62089-0)
- Fujii T & Kawashima K 2001 An independent non-neuronal cholinergic system in lymphocytes. *Japanese Journal of Pharmacology* **85** 11–15. (doi:10.1254/jjp.85.11)
- Fujii T, Takada-Takatori Y & Kawashima K 2008 Basic and clinical aspects of non-neuronal acetylcholine: expression of an independent, non-neuronal cholinergic system in lymphocytes and its clinical significance in immunotherapy. *Journal of Pharmacological Sciences* **106** 186–192. (doi:10.1254/jphs.FM0070109)
- Gocze PM & Freeman DA 2000 Cytotoxic effects of cigarette smoke alkaloids inhibit the progesterone production and cell growth of cultured MA-10 Leydig tumor cells. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* **93** 77–83. (doi:10.1016/S0301-2115(00)00254-2)
- Grando SA, Kawashima K & Wessler I 2003 Introduction: the non-neuronal cholinergic system in humans. *Life Sciences* **72** 2009–2012. (doi:10.1016/S0024-3205(03)00063-8)
- Grozio A, Catassi A, Cavalieri Z, Paleari L, Cesario A & Russo P 2007 Nicotine, lung and cancer. *Anti-Cancer Agents in Medicinal Chemistry* **7** 461–466. (doi:10.2174/187152007781058587)
- Guillaume E, Pineau C, Evrard B, Dupaix A, Moertz E, Sanchez JC, Hochstrasser DF & Jegou B 2001 Cellular distribution of translationally controlled tumor protein in rat and human testes. *Proteomics* **1** 880–889. (doi:10.1002/1615-9861(200107)1:7<880::AID-PROT880>3.0.CO;2-2)
- Haberberger RV, Bodenbenner M & Kummer W 2000 Expression of the cholinergic gene locus in pulmonary arterial endothelial cells. *Histochemistry and Cell Biology* **113** 379–387. (doi:10.1007/s004180000153)
- Hecker A, Mikulski Z, Lips KS, Pfeil U, Zakrzewicz A, Wilker S, Hartmann P, Padberg W, Wessler I, Kummer W *et al.* 2009 Pivotal advance: up-regulation of acetylcholine synthesis and paracrine cholinergic signaling in intravascular transplant leukocytes during rejection of rat renal allografts. *Journal of Leukocyte Biology* **86** 13–22. (doi:10.1189/jlb.1107722)
- Herber DL, Severance EG, Cuevas J, Morgan D & Gordon MN 2004 Biochemical and histochemical evidence of nonspecific binding of $\alpha 7$ nAChR antibodies to mouse brain tissue. *Journal of Histochemistry and Cytochemistry* **52** 1367–1376. (doi:10.1177/002215540405201013)
- Ibanez CF, Pelto-Huikko M, Soder O, Ritzén EM, Hersh LB, Hokfelt T & Persson H 1991 Expression of choline acetyltransferase mRNA in spermatogenic cells results in an accumulation of the enzyme in the postacrosomal region of mature spermatozoa. *PNAS* **88** 3676–3680. (doi:10.1073/pnas.88.9.3676)
- Innis SM, Davidson AG, Bay BN, Slack PJ & Hasman D 2011 Plasma choline depletion is associated with decreased peripheral blood leukocyte acetylcholine in children with cystic fibrosis. *American Journal of Clinical Nutrition* **93** 564–568. (doi:10.3945/ajcn.110.005413)
- Kawashima K & Fujii T 2000 Extraneuronal cholinergic system in lymphocytes. *Pharmacology & Therapeutics* **86** 29–48. (doi:10.1016/S0163-7258(99)00071-6)
- Kirkpatrick CJ, Bittiger F, Nozadze K & Wessler I 2003 Expression and function of the non-neuronal cholinergic system in endothelial cells. *Life Sciences* **72** 2111–2116. (doi:10.1016/S0024-3205(03)00069-9)
- Klein J, Gonzalez R, Köppen A & Löffelholz K 1993 Free choline and choline metabolites in rat brain and body fluids: sensitive determination and implications for choline supply to the brain. *Neurochemistry International* **22** 293–300. (doi:10.1016/0197-0186(93)90058-D)
- Kumar P & Meizel S 2005 Nicotinic acetylcholine receptor subunits and associated proteins in human sperm. *Reproduction of Biological Chemistry* **280** 25928–25935. (doi:10.1074/jbc.M502435200)
- Kummer W, Lips KS & Pfeil U 2008 The epithelial cholinergic system of the airways. *Histochemistry and Cell Biology* **130** 219–234. (doi:10.1007/s00418-008-0455-2)
- Kurzen H, Wessler I, Kirkpatrick CJ, Kawashima K & Grando SA 2007 The non-neuronal cholinergic system of human skin. *Hormone and Metabolic Research* **39** 125–135. (doi:10.1055/s-2007-961816)
- Lips KS, Volk C, Schmitt BM, Pfeil U, Arndt P, Miska D, Erment L, Kummer W & Koepsell H 2005 Polyspecific cation transporters mediate luminal release of acetylcholine from bronchial epithelium. *American Journal of Respiratory Cell and Molecular Biology* **33** 79–88. (doi:10.1165/rcmb.2004-0363OC)
- Livak KJ & Schmittgen TD 2001 Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method. *Methods* **25** 402–408. (doi:10.1006/meth.2001.1262)
- Lonnerberg P & Ibanez CF 1999 Novel, testis-specific mRNA transcripts encoding N-terminally truncated choline acetyltransferase. *Molecular Reproduction and Development* **53** 274–281. (doi:10.1002/(SICI)1098-2795(199907)53:3<274::AID-MRD3>3.0.CO;2-8)

- Marubio LM, del Mar Arroyo-Jimenez M, Cordero-Erausquin M, Léna C, Le Novère N, de Kerchove d'Exaerde A, Huchet M, Damaj MI & Changeux JP 1999 Reduced antinociception in mice lacking neuronal nicotinic receptor subunits. *Nature* **398** 805–810. (doi:10.1038/19756)
- Moser N, Mechawar N, Jones I, Gochberg-Sarver A, Orr-Urtreger A, Plomann M, Salas R, Molles B, Marubio L, Roth U *et al.* 2007 Evaluating the suitability of nicotinic acetylcholine receptor antibodies for standard immunodetection procedures. *Journal of Neurochemistry* **102** 479–492. (doi:10.1111/j.1471-4159.2007.04498.x)
- Nathanson NM 2008 Synthesis, trafficking, and localization of muscarinic acetylcholine receptors. *Pharmacology & Therapeutics* **119** 33–43. (doi:10.1016/j.pharmthera.2008.04.006)
- Okuda T, Haga T, Kanai Y, Endou H, Ishihara T & Katsura I 2000 Identification and characterization of the high-affinity choline transporter. *Nature Neuroscience* **3** 120–125. (doi:10.1038/72059)
- Paleari L, Sessa F, Catassi A, Servent D, Mourier G, Doria-Migletta G, Ognio E, Cilli M, Dominioni L, Paolucci M *et al.* 2009 Inhibition of non-neuronal alpha7-nicotinic receptor reduces tumorigenicity in A549 NSCLC xenografts. *International Journal of Cancer* **125** 199–211. (doi:10.1002/ijc.24299)
- Parsons SM 2000 Transport mechanisms in acetylcholine and monoamine storage. *FASEB Journal* **14** 2423–2434. (doi:10.1096/fj.00-0203rev)
- Ribeiro FM, Black SA, Prado VF, Rylett RJ, Ferguson SS & Prado MA 2006 The "ins" and "outs" of the high-affinity choline transporter ChT1. *Journal of Neurochemistry* **97** 1–12. (doi:10.1111/j.1471-4159.2006.03695.x)
- Sastry BV, Janson VE & Chaturvedi AK 1981 Inhibition of human sperm motility by inhibitors of choline acetyltransferase. *Journal of Pharmacology and Experimental Therapeutics* **216** 378–384.
- Spies M, Lips KS, Kurzen H, Kummer W & Haberberger RV 2006 Nicotinic acetylcholine receptors containing subunits alpha3 and alpha5 in rat nociceptive dorsal root ganglion neurons. *Journal of Molecular Neuroscience* **30** 55–56. (doi:10.1385/JMN:30:1:55)
- Wang N, Orr-Urtreger A, Chapman J, Rabinowitz R, Nachman R & Korcyn AD 2002 Autonomic function in mice lacking alpha5 neuronal nicotinic acetylcholine receptor subunit. *Journal of Physiology* **542** 347–354. (doi:10.1113/jphysiol.2001.013456)
- Wang N, Orr-Urtreger A, Chapman J, Rabinowitz R & Korcyn AD 2003 Deficiency of nicotinic acetylcholine receptor beta 4 subunit causes autonomic cardiac and intestinal dysfunction. *Molecular Pharmacology* **63** 574–580. (doi:10.1124/mol.63.3.574)
- Wessler I & Kirkpatrick CJ 2008 Acetylcholine beyond neurons: the non-neuronal cholinergic system in humans. *British Journal of Pharmacology* **154** 1558–1571. (doi:10.1038/bjp.2008.185)
- Wessler I, Roth E, Deutsch C, Brockerhoff P, Bittinger F, Kirkpatrick CJ & Kilbinger H 2001 Release of non-neuronal acetylcholine from the isolated human placenta is mediated by organic cation transporters. *British Journal of Pharmacology* **134** 951–956. (doi:10.1038/sj.bjp.0704335)
- Yajima T, Inoue R, Matsumoto M & Yajima M 2010 Non-neuronal release of ACh plays a key role in secretory response to luminal propionate in rat colon. *Journal of Physiology* **589** 953–962. (doi:10.1113/jphysiol.2010.199976)
- Yamamoto Y, Isoyama E, Sofikitis N & Miyagawa I 1998 Effects of smoking on testicular function and fertilizing potential in rats. *Urological Research* **26** 45–48. (doi:10.1007/s002400050022)
- Zhu BC, Chiochio SR, Suburo AM & Tramezzani JH 1995 Monoaminergic and peptidergic contributions of the superior and the inferior spermatic nerves to the innervation of the testis in the rat. *Journal of Andrology* **16** 248–258.

Received 9 July 2010

First decision 13 September 2010

Revised manuscript received 15 February 2011

Accepted 11 April 2011