Involvement of the oestrogenic receptors in superior mesenteric ganglion on the ovarian steroidogenesis in rat

Adriana Vega Orozco, Cristina Daneri, Gabriel Anesetti¹, Ricardo Cabrera^{2,3}, Zulema Sosa and Ana M Rastrilla

Laboratorio de Biología de la Reproducción (LABIR), Facultad de Química, Bioquímica y Farmacia, Universidad Nacional de San Luis, San Luis, Argentina, ¹Departamento de Histología y Embriología, Facultad de Medicina, Universidad de la República, General Flores 2125, CP 11800 Montevideo, Uruguay, ²Instituto de Ciencias Biomédicas, Facultad de Ciencias Médicas, Universidad de Mendoza, Mendoza, Argentina and ³Instituto de Medicina y Biología Experimental de Cuyo (IMBECU)-Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET), Mendoza, Argentina

Correspondence should be addressed to A Vega Orozco; Email: adriana.vega@um.edu.ar; Z Sosa; Email: zsosa@unsl.edu.ar

Abstract

Oestradiol (E₂) is a key hormone in the regulation of reproductive processes. The aims of this work were a) to examine the distributions of oestrogen receptor α (ER α) and ER β in the neurons of the superior mesenteric ganglion (SMG) in the oestrus stage by immuno-histochemistry, b) to demonstrate whether E₂ in the SMG modifies progesterone (P₄), androstenedione (A₂) and nitrite release in the ovarian compartment on oestrus day and c) to demonstrate whether E₂ in the ganglion modifies the activity and gene expression in the ovary of the steroidogenic enzymes β -hydroxysteroid dehydrogenase (β -HSD) and 20α -hydroxysteroid dehydrogenase (20α -HSD). The *ex vivo* SMG–ovarian nervous plexus–ovary system was used. E₂, tamoxifen (Txf) and E₂ plus Txf were added in the ganglion to measure ovarian P₄ release, while E₂ alone was added to measure ovarian A₂ and nitrites release. Immunohistochemistry revealed cytoplasmic ER α immunoreactivity only in the neural somas in the SMG. E₂ increased ovarian P₄ and A₂ release at 15, 30 and 60 min but decreased nitrites. The activity and gene expression of β -HSD increased, while the activity and gene expression of 20α -HSD did not show changes with respect to the control. Txf in the ganglion diminished P₄ release only at 60 min. E₂ plus Txf in the ganglion reverted the effect of E₂ alone and the inhibitory effect of Txf. The results of this study demonstrate that ER α activation in the SMG has an impact on ovarian steroidogenesis in rats, thus providing evidence for the critical role of peripheral system neurons in the control of ovarian functions under normal and pathological conditions.

Reproduction (2012) **143** 183–193

Introduction

There is ample evidence demonstrating a functional interaction between the peripheral nervous system and the reproductive system (Sosa *et al.* 2000, 2004, Casais *et al.* 2006). The ovary receives sympathetic innervation from the prevertebral ganglia by two routes: the ovarian nervous plexus (ONP) and the superior ovarian nerve (SON). The ONP fibres originate mostly in the neurons of the superior mesenteric ganglion (SMG; Baljet & Drukker 1979, Klein & Burden 1988a) and accompany the ovary artery in its entrance to the hilum. This pathway not only innervates the blood vessels but also the veins that form a net around the theca cells (Lawrence & Burden 1980). The SMG is included in the sympathetic prevertebral chain; it is constituted by principal neurons and interneurons (Eränkö 1978, Prud'Homme *et al.* 1999) and it has a profuse capillary plexus that constitutes a microcirculation among the different

ganglionic structures. The sympathetic ganglia possess a great variety of specific receptors and neurotransmitters, such as catecholamines (Klein & Burden 1988a) and neuropeptides (Dalsgaard *et al.* 1982). These ganglia are active points of concurrency for hormone signals, as revealed by the metabolic changes induced by steroids, as well as by the existence of putative receptors for some hormones such as oestradiol (E₂; Cardinali *et al.* 1983).

Oestrogens are important for the development of sexual phenotype and differentiation of the central nervous system (CNS). They play crucial roles in the development of the pituitary gonadal axis, ovarian folliculogenesis, ovulation and implantation, thereby acting on peripheral organs and sympathetic neurons associated with functions related to reproduction (Anesetti *et al.* 2009). Zoubina & Smith (2003) demonstrated that a small number of uterine-projecting neurons located in the SMG and other sympathetic ganglia expressed oestrogen receptors (ERs). Their

actions are mediated through two distinct receptors, ER α and ER β , which bind to specific DNA sequences regulating transcription. ER α is also localised in the plasma membrane and may help to elucidate the mechanisms through which rapid, 'non-genomic' oestrogen signalling occurs (Deecher *et al.* 2003, Doolan & Harvey 2003, Wang *et al.* 2005). ERs are regulated in a tissue- and/or cell-type specific manner and E₂ itself seems to play an important role in this regulation (Weihua *et al.* 2003).

A simple dose of E₂ valerate can modify the levels of expression of the enzyme tyrosine hydroxylase (TH), the rate-limiting enzyme in catecholamine biosynthesis, which induces changes in the nervous sympathetic activity in the ovary of mammals (Shinohara *et al.* 1998, Lara *et al.* 2000), provoking alterations in the follicular growth that lead to the development of polycystic ovaries (Lara *et al.* 2002, Rosa *et al.* 2003).

This enzyme, together with dopamine β -hydroxylase, which also participates in the biosynthesis pathway of catecholamines, colocalises with immunoreactive neuropeptides and choline acetyltransferase, the enzyme for the acetylcholine synthesis pathway (Lars-Gösta *et al.* 1997).

Indirect immunofluorescence studies have demonstrated the presence of the isoenzyme neuronal nitric oxide synthase (nNOS) in the SMG (Quinson *et al.* 2001). Berman *et al.* (1998) have demonstrated that ER α is predominantly expressed by nitric oxide synthase (NOS)-positive in parasympathetic pelvic ganglion neurons. There is evidence that NO decreases the steroidogenesis in different experimental schemes (Olson *et al.* 1996, Snyder *et al.* 1996), causing an inhibition of the enzymatic activities and gene expression of steroid synthesis-limiting enzymes such

as 3 β -hydroxysteroid dehydrogenase (3 β -HSD) in bovine luteal cells (Rekawiecki *et al.* 2005). Vega Orozco *et al.* (2006) have demonstrated in an *ex vivo* system that the ganglionic cholinergic stimulus modifies the ovarian steroids and NO release in the oestrus stage.

With these antecedents in mind and considering the neural influence from the SMG through the ONP on ovarian steroidogenesis, the aims of this work were a) to examine the distributions of ER α and ER β in neurons of the SMG during the oestrus (E) stage by immunohistochemistry, b) to demonstrate whether E₂ in the SMG modifies progesterone (P₄), androstenedione (A₂) and nitrite (NO) release in the ovarian compartment on oestrus day and c) to demonstrate whether E₂ in the ganglion modifies the activity and gene expression in the ovary of the steroidogenic enzymes 3 β -HSD and 20 α -HSD.

Results

Localisation of ERs in the SMG on oestrus day by immunohistochemistry

The histological study of the SMG showed a thin capsule of connective tissue and the parenchyma exhibited several neuronal profiles (Fig. 1A). ER α immunoreactivity was observed in most neuronal somas of the SMG, and its pattern was cytoplasmic and heterogeneous (Fig. 1B). Uterine control sections displayed the classical nuclear localisation of this receptor. ER β immunoreactivity was not observed in neuronal somas (Fig. 1C). Figure 1D shows a high magnification of Fig. 1C (40 $\times$). Rat ovary was used as the positive control for ER β (Fig. 1D'). No positive structures or cells were observed in the SMG sections when the primary antiserum was not added (negative control; Fig. 1D'').

Figure 1 (A–D) (A) Low magnification of the SMG showing that almost all neurons displayed immunoreactivity for ER α (4 $\times$). (B) High magnification of A (40 $\times$). Note the extensive immunoreactivity restricted to cytoplasm. No nuclear signal was detected. (B') Inset of positive control for ER α in rat uterus showing intense signal in glandular and stromal endometrial cells (40 $\times$). (C) Low magnification of the SMG showing absence of ER β reactivity (4 $\times$). (D) High magnification of C (40 $\times$). (D') Inset of positive control for ER β in rat ovary (40 $\times$). (D'') Inset of negative control for ER β in rat SMG.

Figure 2 Effect of oestradiol (E_2) and tamoxifen (Txf) in the superior mesenteric ganglion on the ovarian progesterone release in the superior mesenteric ganglion–ovarian nervous plexus–ovary system removed from oestrus day rat. The systems were incubated in a Krebs–Ringer buffer, plus ascorbic acid (0.1 mg/ml in Krebs–Ringer) solution, at 37 °C in an atmosphere of 95% O_2 to 5% CO_2 for 120 min without (control) and with the addition of E_2 (10^{-6} mol/l), Txf (10^{-6} mol/l) and E_2 plus Txf in the ganglion compartment (experimental group). E_2 , E_2 valerate. Values are the mean $\pm$ S.E.M. of six animals per experimental group. Columns with * and * indicate statistical significance of $*P < 0.001$ and $*P < 0.05$, respectively, compared with the control group (Student's t -test, ANOVA, Duncan's test).

Ganglionic effect of E_2 and tamoxifen on ovarian P_4 release on oestrus

To evaluate whether the presence of E_2 and its antagonist in the ganglion compartment modulated the release of ovarian P_4 in the *ex vivo* system and considering that this steroid has proved to be the most sensitive steroid to neural influence in *in vitro* studies, E_2 (10^{-6} mol/l), tamoxifen (Txf) only and E_2 plus Txf were added in the SMG. Txf was used since this antagonist is specific for $ER\alpha$ (Sun *et al.* 2010).

E_2 in the ganglion increased ovarian P_4 release at 15 min (0.008 ± 0.001 vs 0.03 ± 0.001), 30 min (0.023 ± 0.002 vs 0.048 ± 0.001) and 60 min (0.015 ± 0.006 vs 0.09 ± 0.004 ; $*P < 0.001$), whereas Txf alone decreased P_4 release only at 60 min (0.015 ± 0.002 vs 0.005 ± 0.001 ; $*P < 0.05$) with respect to the control group. The addition of E_2 (10^{-6} mol/l) plus Txf in the ganglion did not show changes in ovarian P_4 release in comparison with the control group (Fig. 2).

Effect of E_2 in the ganglion on the enzymatic activity and gene expression of the ovarian 3β -HSD and 20α -HSD steroidogenic enzymes on oestrus

To evaluate whether the addition of E_2 in the mesenteric ganglion modified the activity and gene expression of the P_4 synthesis enzymes in the oestrus stage, the ovaries of the system were obtained after 120 min of incubation.

P_4 in the ganglion increased the ovarian enzymatic activity and gene expression of 3β -HSD at 120 min of incubation ($*P < 0.001$; Figs 3a and 4a), whereas the activity and gene expression of 20α -HSD did not show significant changes at 120 min with respect to the control (Figs 3b and 4b).

Effect of E_2 in ganglion on ovarian A_2 release on oestrus

Considering that A_2 is synthesised by the ovarian interstitial cells and that ONP innervated these structures, A_2 release in the ovarian compartment by stimulation with E_2 at the ganglionic level was assessed.

E_2 in the ganglion increased the release of A_2 in the ovarian compartment at 15 min (0.024 ± 0.01 vs 0.354 ± 0.01), 30 min (0.595 ± 0.01 vs 0.92 ± 0.01 ; $*P < 0.001$) and 60 min (0.42 ± 0.01 vs 0.98 ± 0.01 ; $*P < 0.001$; Fig. 5).

Effect of E_2 in the ganglion on ovarian nitrite release on oestrus

The modulation of the release of nitrites by neural stimulus and their importance in the reproductive function has been demonstrated in our research group; therefore, we attempted to determine whether the addition of E_2 in the ganglion compartment modifies nitrite release on the ovarian compartment on oestrus day.

E_2 caused a significant decrease in nitrite release at all incubation times in comparison with the control group ($*P < 0.001$; Fig. 6).

Discussion

Given the importance of the neural regulation from the peripheral nervous system on ovarian physiology (Sosa *et al.* 2000, Casais *et al.* 2006, Vega Orozco *et al.* 2010) and taking into account that E_2 acts on peripheral organs and on the plasticity of sympathetic neurons associated with reproduction (Anesetti *et al.* 2009), the general aim of this work was to examine the distributions of $ER\alpha$ and $ER\beta$ in neurons of the SMG and to evaluate the importance of these receptors in ovarian steroidogenesis and their possible relation with NO release.

In this work, $ER\alpha$ was typified by immunohistochemistry in the SMG, but $ER\beta$ was not detected probably due to the fact that receptors of this type have been

Figure 3 Effect of oestradiol (E_2) in the ganglion compartment on the enzymatic activity of 3β -HSD (a) and 20α -HSD (b) in ovaries extracted from the superior mesenteric ganglion–ovarian nervous plexus–ovary system with and without the addition of E_2 (10^{-6} mol/l) in the mesenteric ganglion at 120 min of incubation on oestrus day. E_2 , E_2 valerate. Values are the mean $\pm$ S.E.M. of six animals in two experiments. Columns with * indicate statistical significance of $*P < 0.001$ compared with the control group (Student's t -test; ANOVA, Duncan's test).

Figure 4 Ganglionic effect of oestradiol on mRNA abundances of enzymes 3β-HSD (a) and 20α-HSD (b) obtained from rat ovary previously incubated in the superior mesenteric ganglion–ovarian nervous plexus–ovary system on oestrus day. Measurement by RT-PCR of the expression of 3β-HSD and cyclophilin A. Ethidium bromide fluorescence photograph of the gel electrophoresis of the amplification products. Expression of 3β-HSD relative to cyclophilin A. The gel photographs were quantified using Image J (NIH, Bethesda, MA, USA) and expressed as arbitrary units. Results are expressed as the mean $\pm$ S.E.M. ($n=3$). * $P<0.001$.

fundamentally found in prepubertal (Shinohara *et al.* 2000, Anesetti *et al.* 2009) and adult rats in other cycle stages (Zoubina & Smith 2002). The results obtained in this work differ from those reported by Zoubina & Smith (2002), who found, in a different experimental scheme, that the vast majority of sympathetic neurons express ERβ immunoreactive protein, whereas a smaller, presumably overlapping subset expresses the ERα. Papka *et al.* (2001) found that both receptors could potentially be affected by changes in endogenous oestrogen levels, which could explain why ERβ was not found in our study. Our findings showed that in SMG neurons, ERα was mainly distributed uniformly throughout the neuronal cytoplasm on oestrus day, but it was not observed in satellite or Schwann cells, in agreement with the findings reported by Anesetti *et al.* (2009) on lumbar paravertebral and prevertebral sympathetic ganglia in prepubertal animals. Moreover, immunocytochemical studies have demonstrated an extranuclear localisation for the ERs in neurons of rat brain (Blaustein 1992).

It is important to emphasise that in the case of neurons associated with the ovary, the studies are scanty and basically linked to long-term effects on the physiology of the organ, when oestrogens are applied in early stages of the development (Shinohara *et al.* 1998, 2000, Lara *et al.* 2000, Koszykowska *et al.* 2011). The functions of ERs in sympathetic ganglia are related to the protection of the neurons against damage and the regulation of the growth of the neuronal prolongations (Sohrabji *et al.* 1994a, 1994b). Both ERs have similar affinities for binding E₂

(Mosselman *et al.* 1996, Kuiper *et al.* 1997) and could exert their effect through different signalling paths. However, the tissue distribution and level of expression of ERα and ERβ vary in sympathetic ganglia (Zoubina & Smith 2002).

It has been demonstrated that besides the well-studied transcriptional effects of E₂, there are rapid effects, i.e. occurring within seconds or minutes after the addition of E₂ (Song *et al.* 2005, Song & Santen 2006, Warner & Gustafsson 2006). These rapid effects are probably produced through non-genomic mechanisms, as has been observed in several tissues (Heldring *et al.* 2007). However, there is no agreement yet as to the participation of the classical oestrogenic receptors (Levin 2001, Deecher *et al.* 2003, Razandi *et al.* 2004) or whether there is a distinct membrane-associated receptor, such as GPR30 (Doolan & Harvey 2003, Dun *et al.* 2009).

It has been reported that oestrogen rapidly (between 5 and 15 min) increases the activity of mitogen-activated protein kinase, which persists for at least 2 h (Singh *et al.* 1999). In concordance with the above results, the addition of E₂ in the SMG produces a rapid increase in ovarian P₄ and A₂ release at 15 min, reaching a maximum at 60 min and a decrease at 120 min, in agreement with Singh *et al.* (1999). This effect observed on ovarian P₄ release might be due to the classical ERα found in the SMG, considering that by immunohistochemistry the ERβ was not observed in the ganglion, or else, to some other type of oestrogenic receptor, probably GPR30, which has been found at the sympathetic ganglionic level (Dun *et al.* 2009), might be involved in the rapid responses obtained on the release of ovarian P₄.

The effect of E₂ stimulus in the ganglion on ovarian P₄ and A₂ released might be accounted for by an increase in cAMP levels in neurons that constitute the SMG and the consequent increase in ovarian noradrenaline (NA) as

Figure 5 Effect of oestradiol (E₂) on ovarian androstenedione release in the superior mesenteric ganglion–ovarian nervous plexus–ovary system removed from rats on oestrus day. The systems were incubated in Krebs–Ringer buffer, plus ascorbic acid (0.1 mg/ml in Krebs–Ringer) solution, at 37 °C in an atmosphere of 95% O₂ to 5% CO₂ for 120 min without (control) and with the addition of E₂ (10^{−6} mol/l) to the ganglion compartment (experimental group). E₂, E₂ valerate. Each bar represents the mean $\pm$ S.E.M. of six animals per experimental group. * $P<0.001$ compared with the control group (Student's *t*-test; ANOVA, Duncan's test).

Figure 6 Effect of oestradiol (E_2) in the ganglion compartment on nitric oxide (NO) release in the incubation liquid of the ovarian compartment in the superior mesenteric ganglion–ovarian nervous plexus–ovary system of rats on oestrus day. The system was incubated in Krebs–Ringer buffer, plus ascorbic acid (0.1 mg/ml in Krebs–Ringer) solution, at 37 °C in an atmosphere of 95% O_2 to 5% CO_2 for 120 min without (control) and with the addition of E_2 (10^{-6} mol/l) to the ganglion compartments (experimental group). E_2 , E_2 valerate. Values are the mean $\pm$ S.E.M. of six animals per experimental group. Columns with * indicate statistical significance of $*P < 0.001$ compared with the control group (Student's *t*-test; ANOVA, Duncan's test).

stated by [Lara et al. \(2002\)](#), with an increased capacity of the ovarian nerve terminals to incorporate and release NA. The effects of catecholamines in the ovary have been demonstrated in other experimental schemes ([Aguado & Ojeda 1984](#), [Barria et al. 1993](#)).

[Sosa et al. \(2004\)](#) demonstrated in an *ex vivo* system that the cholinergic stimulus in the coeliac ganglion caused changes in NA and P_4 release at the ovarian level, according to the stage of the oestrous cycle, and this constitutes one of the factors governing the secretory activity of the ovarian steroids ([Sosa et al. 2004](#)). Recent studies using the SMG–ONP–ovary system have determined an increase in NA release in the ovarian compartment on oestrus day by cholinergic ganglionic stimulus ([A Vega Orozco, C Daneri, D Bronzi, Z Sosa, AM Rastrilla](#), unpublished observations, 2010).

It has been demonstrated that the neuronal population in mesenteric ganglia is very heterogeneous, with numerous peptides and neurotransmitters. It has been shown in our research group that ganglionic stimulation in this *ex vivo* system releases not only catecholamines but also gonadotropin-releasing hormone (GnRH) and NO ([A Vega Orozco, C Daneri, D Bronzi, Z Sosa, AM Rastrilla](#), unpublished observations, 2010). [Garraza et al. \(2004\)](#) demonstrated that the effect of other neurotransmitters such as substance P (SP) and neuropeptide Y (NPY) depends on the oestric cycle and that they can exert effects on the postganglionic terminal that reaches the ovary. These authors demonstrated that the addition of NA alone in hemi-ovaries incubations increases P_4 release while the addition of NPY, SP and vasoactive intestinal peptide (VIP) alone and together with NA generally decreases P_4 release at short times and increases

it at long times. In addition, [Miyamoto et al. \(1993\)](#) reported that the addition of NPY, SP and VIP in cultures of luteal cells at short times and in the absence of luteinising hormone (LH) decreases the release of P_4 , although using a different experimental scheme. In our system, which is free of humoral influences, there was an increase in P_4 by oestrogenic stimulation at ganglionic level.

With regard to the enzymes involved in the synthesis and degradation of P_4 , the addition of E_2 in the ganglion increased the enzymatic activity and gene expression of 3β -HSD but not 20α -HSD in the ovary at 120 min of incubation in relation to the control. This result is surprising since there are no reports of enzymatic activity modulation at such short times. Variations in luteal cell cultures occurring at 36 h without neural influence and without the paracrine relations present in our *ex vivo* system have been reported ([Tellería et al. 1995](#), [Stocco & Deis 1996](#)). It is important to note that although in this case E_2 in the ganglion has a short time effect (15 min), [Singh et al. \(1999\)](#) report that E_2 acts between 5 and 120 min. For this reason, the enzymatic activity and gene expression of ovarian 3β -HSD were measured at 120 min of incubation.

Considering that NA released at the ovarian level by the oestrogenic ganglionic stimulus might be responsible for the increase in ovarian P_4 , the findings of this study are in agreement with postulations by [Hsueh et al. \(1984\)](#) and [Miszkiewicz et al. \(1999\)](#) that NA stimulates the activity of 3β -HSD. At the same time, P_4 produced by NA stimulus reduces the activity of monoaminooxidase (MAO) and catechol-*O*-methyl-transferase, the enzymes primarily responsible for the degradation of catecholamines ([Mannisto & Kaakkola 1999](#), [Eisenhofer et al. 2004](#)). Thus, in this way, P_4 prolongs the half-life of NA and the duration of its stimulatory influence on P_4 synthesis.

Also, taking into account that [Kotwica et al. \(2004\)](#) and [Rekawiecki et al. \(2008\)](#) have postulated the autoregulation of P_4 synthesis in the corpus luteum, which affects the transcription of the genes encoding steroidogenic enzymes, it can be assumed that the values of P_4 can be due to the increase in the activity and gene expression of 3β -HSD caused by the same P_4 released throughout time. The accumulation of P_4 in ovarian steroidogenic cells is also influenced by the participation of the enzyme 20α -HSD, a P_4 degradation enzyme, which has been determined, cloned and sequenced in the corpus luteum of rats ([Mao et al. 1997](#)). In our case, this enzyme did not suffer variations at the studied times.

When the ERs antagonist was added, Txf alone in the ganglion inhibited ovarian P_4 release only at 60 min. Possibly, the effect of Txf occurs at the ganglionic level. In fact, [Paech et al. \(1997\)](#) and [Kuiper et al. \(1999\)](#) have demonstrated that the effect of Txf varies with the type of tissue and has action mechanisms that are different compared with traditional receptors ([Paige et al. 1999](#), [Zou et al. 1999](#)). It is possible that, due to the E_2 level at this stage, the levels of the ER β in the ganglion are too low

to be detected by immunohistochemistry. Both receptors are probably co-expressed at the ganglionic level, where ER β antagonises the actions that depend on ER α as has been observed in other studies (Paech *et al.* 1997). Thus, the addition of an antagonist as Txf in the ganglion caused an inhibition of ovarian P₄ release at 60 min, an opposite to that of E₂ alone at the ganglionic level. The effects of E₂ and Txf alone were reverted by the addition of E₂ plus Txf, taking the P₄ levels to control values.

On the other hand, we postulate that the inhibitory effect of Txf alone in the ganglion might be through the ER β . Webb *et al.* (1995) have demonstrated that anti-oestrogens such as Txf act as agonists of an alternative pathway (AP-1) to the classical ERs one, which is more potent in the presence of ER β . Thus, it has been demonstrated that the ER β potentially increases the transcription dependent on this alternative way in the presence of anti-oestrogens, but it is not activated in the presence of oestrogens, which might explain an inhibitory effect of Txf through this receptor that apparently acts as a repressor (Navarro Despaigne 2001). Thus, this effect is reverted by the addition of E₂ plus Txf at the ganglionic level. This issue will be the subject of future research.

It must be noted that in previous work, we demonstrated in the same SMG–ONP–ovary system that, besides catecholamines, other neurotransmitters such as NO may be released from the ONP by way of the ganglionic stimulus (Vega Orozco *et al.* 2010). Evidence has been reported that supports the involvement of NO in ovarian physiology (Fridén *et al.* 2000, Motta *et al.* 2001). In addition, it has been demonstrated that oestrogen up-regulates neuronal NOS expression through ER α in parasympathetic pelvic ganglion neurons (Berman *et al.* 1998). However, other investigators have shown that the various ERs on the cell surface are coupled to constitutive NOS-derived NO release (Stefano *et al.* 2003).

Our research group has demonstrated that the cholinergic agent in the coeliac ganglion stimulates NO release through NOS in prepubertal rat ovaries (Delgado *et al.* 2004). Studies with antagonists of the neuronal and inducible NO enzyme, both at the ganglionic and ovarian levels, showed that NO can reach the ovary by neural ways or be synthesised at the ovarian level (Delgado *et al.* 2004, Casais *et al.* 2006). In addition, Vega Orozco *et al.* (2006, 2010) have demonstrated that NO is involved in the formation and maintenance of the corpus luteum during the oestrous cycle through the ONP pathway.

Considering the above antecedents, it was decided to evaluate whether NO release is modified at the ovarian level by ganglionic oestrogenic stimulus in the *ex vivo* system and whether its presence is involved in ovarian steroidogenesis. Nitrites were found to decrease in the ovary compartment, which might be attributed to the direct action of oestrogen on the activity of one or more isoforms of the NO synthesis enzyme present in the ganglion, as was also observed by other authors (Stefano *et al.* 2003).

On the other hand, it is important to note that NO has been reported to decrease P₄ and A₂ release in *in vitro* cellular cultures (Olson *et al.* 1996, Dunnam *et al.* 1999, Fridén *et al.* 2000). Motta *et al.* (2001) have shown in rat that endogenous NO increased the production of glutathione and P₄ in corpora lutea in the middle stage of development. This result shows a dual, protective or pro-oxidising effect according to the stage of the oestrous cycle during which the tissue was obtained (Dong *et al.* 1999). Therefore, NO may be a luteolytic and/or luteotrophic factor on the corpus luteum, depending on the stage of the oestrous cycle (Olson *et al.* 1996, Fridén *et al.* 2000, Motta *et al.* 2001). In addition, Yamamoto *et al.* (1993) have demonstrated that NO might be involved in the modulation of the output and synthesis of NA, which is in agreement with the postulation that NA might be responsible for P₄ increase.

E₂ action in the ganglion observed in the present work probably favours the observed ovarian P₄ increase, having a trophic effect on the corpus luteum. On the basis of the results we obtained, we can propose that oestrogenic stimulus on the SMG would act in favour of the events that lead to the ovulation and later formation of the corpus luteum.

There is ample evidence suggesting that the sympathetic nervous system is involved in pathologies associated with the reproductive system. The role of oestrogen at the ganglionic level on ovarian steroidogenesis remains unclear and controversial. Anesetti *et al.* (2009) demonstrated that elevated levels of exogenous oestrogen during the prepubertal period in postganglionic sympathetic neurons increased sympathetic coeliac neuronal size and modified the expression of p^{75NTR} (p75 neurotrophin receptor), inducing a significant increase in catecholaminergic innervation of the ovary.

Polycystic ovary syndrome (PCOS) is associated with an abnormal activation of the sympathetic nervous system, showing an impaired metabolism of NA in adolescents suffering from this disease (Garcia-Rudaz *et al.* 1998).

The results of this study demonstrate that ER α activation in the SMG has an impact on ovarian steroidogenesis in rats at the ovarian level, thus providing evidence for the critical role of peripheral system neurons in the control of ovarian functions under normal and possibly under pathological conditions such as PCOS and ovarian cancer.

This work is expected to contribute new evidence for the clinical importance of these receptors in the therapy and prevention of ovarian pathology.

Materials and Methods

Animals

Virgin Holtzman strain adult female rats on oestrus (E) day weighing 250 g were used in all the experiments. The rats were kept in a light- (lights on from 0700 to 1900 h) and

temperature-controlled room ($24 \pm 2^\circ\text{C}$). Animals had free access to food (Cargill, SACI, Saladillo, Buenos Aires, Argentina) and tap water was available *ad libitum*. Vaginal smears were taken daily, and only the rats exhibiting at least two 4-day consecutive oestrous cycles were used. Groups of six animals were used for each group utilised in the experimental procedure. The experiments were performed per duplicate according to the procedures approved in the UFAW Handbook on the Care and Management of Laboratory Animals (Poole 1999). The experimental protocol was approved by the University of San Luis Animal Care and Use Committee (number protocol: B17/04, ordinance CD 006/02).

Chemicals

The compounds E_2 valerate (E_2), Txf, ascorbic acid, BSA fraction V, sulphanilamide and N-(1-naphthyl)ethylenediamine were purchased from Sigma-Aldrich Co. Other reagents and chemicals were of analytical grade. $1,2,6,7\text{-}[^3\text{H}]\text{-P}_4$ (107.0 Ci/mmol), $1,2,6,7\text{-}[^3\text{H}]\text{-androst 4-ene-3,17 dione}$ (115.0 Ci/mmol) and $17\text{-}\beta\text{-}2,4,6,7\text{-}[^3\text{H}]\text{-estradiol}$ (102.0 Ci/mmol) were provided by New England Nuclear Products (Boston, MA, USA).

Experimental procedures

Extraction of SMG fixed in situ for localisation of oestrogenic receptors by immunohistochemistry

Rats on oestrus day were anaesthetised under ether bell. The SMG and nearby ganglia were fixed *in situ* with Bouin solution for subsequent extraction and application of immunohistochemistry.

Extraction of the SMG–ONP–ovary system

Groups of six animals on oestrus day were used. The procedure was carried out between 1500 and 1600 h taking into account previous descriptions of the anatomical trajectory of this neural pathway as a guide (Lawrence & Burden 1980, Klein & Burden 1988a, 1988b, Vega Orozco *et al.* 2006).

The piece of tissue utilised in this work corresponds to the whole SMG where the principal neurons are located, whose axons constitute the ONP that reaches the ovary, without the presence of the coeliac ganglion (Vega Orozco *et al.* 2006).

The anatomical localisation of the SMG and coeliac ganglia (CG) is different, as well as the neuron type and size. The SMG was extracted from the area proximal to the CG, proximal to the renal artery and abdominal aorta artery. It is important to clarify immediately before and after incubation (120 min) that the presence of the SMG was confirmed in the system by the application of routine histological techniques. The presence of the ganglion was verified in the removed system before incubation in similar conditions to the systems that were going to be used later.

Rats were anaesthetised with ether under a bell, and the *ex vivo* system was immediately removed by dissection.

Each system consisted of the ovary, the fibres constituting the ONP, parallel to the ovarian artery, and the SMG, surrounded by some small ganglia.

The total surgical procedure was completed in 1–2 min. The strip of tissue was carefully dissected, avoiding contact between the surgical instruments, the nerve fibres and the ganglion to prevent spontaneous depolarisations of the nerves, which might have been caused by inappropriate contact.

The extracted systems were washed with incubation solution and immediately placed in a cuvette with two compartments. Each cuvette contained 2 ml of work solution, Krebs–Ringer bicarbonate buffer, pH 7.4, in the presence of dextrose (0.1 mg/ml) and BSA (0.1 mg/ml) as described in previous studies (Sosa *et al.* 2000, Vega Orozco *et al.* 2006).

The ganglion was placed in one compartment and the ovary in the other, both joined by the ONP, which was maintained humid with the work solution. The system was immediately preincubated in a metabolic bath at 37°C in a saturated O_2/CO_2 (95–5%) atmosphere for 15 min to achieve the stabilisation of the system as described in a previous study (Vega Orozco *et al.* 2006).

After 15 min of preincubation (time 0 of incubation), the Krebs–Ringer solution was changed in both compartments, and ascorbic acid (0.1 mg/ml in Krebs–Ringer) was added as an antioxidant agent (Koh & Hille 1997) in the ganglion compartment whereas the Krebs–Ringer solution was added in the ovary compartment. The P_4 , A_2 and nitrite values released under these conditions were considered as control values (control group).

The experimental groups proposed were a) SMG–ONP–ovary system with the addition of E_2 (10^{-6} mol/l) in the ganglion compartment, b) SMG–ONP–ovary system with addition of Txf (10^{-6} mol/l) in the ganglion compartment and c) SMG–ONP–ovary system with addition of Txf (10^{-6} mol/l) and E_2 (10^{-6} mol/l) 5 min later in the ganglion compartment. Groups b and c were only used for P_4 release determination and for evaluating whether Txf acts as an oestrogenic antagonist at ganglionic level. The control group was SMG–ONP–ovary system without treatment (control). The different substances were dissolved in equal concentration (10^{-6} mol/l) and volumes (2 ml) of Krebs–Ringer buffer plus ascorbic acid.

For the system standardisation, the dose/response curve was obtained for this *ex vivo* system and at the working times used. The curve shows the highest P_4 increase with the used dose (10^{-6} mol/l) compared with other doses used.

Liquid samples from the ovary compartment (250 μl) were collected at 15, 30, 60 and 120 min and kept at -20°C until the determination of P_4 and A_2 by RIA and NO using the Griess method.

The results were expressed as the nanogram of P_4 and picogram of A_2 per milligram of ovarian tissue per millilitre and nitrites as nanomol of nitrites per milligram of ovarian tissue per millilitre (nmol/mg ovary/ml) all against the incubation time. The respective corrections were made in all cases considering the volume extracted in each tested period. When the system incubation was finished (120 min), the ovaries were kept at -80°C until determination of $3\beta\text{-HSD}$ and $20\alpha\text{-HSD}$ activities and gene expression.

Immunohistochemistry

After the experimental treatment, the SMGs were fixed in 4% paraformaldehyde, included in paraffin and sectioned at 5 μm .

For immunohistochemistry, sections were deparaffinised, rehydrated and rinsed in distilled water. After antigen retrieval using microwave treatment at 160 W in 10 mM sodium citrate, pH 6.0, for 30 min, sections were washed in PBS and incubated for 20 min in 0.3% hydrogen peroxide in PBS to inhibit endogenous peroxidases. Then, sections were permeabilised by 30-min incubation in PBS containing 0.5% Triton X-100 and blocked in PBS containing 0.5% Triton X-100 and 2% BSA for 1 h. Sections were incubated overnight at 4 °C with the appropriate primary antibody diluted in blocking solution. The primary antibodies used were as follows: a polyclonal rabbit anti-ER α (ER α , MC-20; SC-542, Santa Cruz Biotechnology, Santa Cruz, CA, USA; 1:400) and a rabbit polyclonal anti-serum against ER β (ER β , H-150; SC-8974 Santa Cruz Biotechnology; 1:100). Subsequently, sections were rinsed in PBS, incubated with biotinylated secondary antibody (1:300 in blocking solution) for 1 h, washed in PBS and incubated for 1 h with avidin-biotin complex (Vector Laboratories, Burlingame, CA, USA). After several washes in PBS, sections were exposed to a 0.02% 3,3'-diaminobenzidine (Sigma-Aldrich Co.) solution. The reaction was stopped in tap water and some sections were counterstained in haematoxylin (20 s), dehydrated in ethanol, cleared in xylene, coverslipped and examined on a Nikon photomicroscope. Each immunoreaction was performed on at least three slides from each set of ganglia. Sections of uterus and ovary were processed in parallel under identical conditions and used as positive controls; staining in these tissues showed strong, predominantly nuclear localisation with both antibodies. To confirm the specificity of the staining, selected sections were incubated without primary or secondary antibodies with positive and negative controls being included.

Nitrite assay

Levels of nitrite, a water-soluble metabolite of NO, were measured spectrophotometrically by the Griess method and were expressed in nanomole of nitrite per milligram of ovarian tissue per millilitre (nmol/mg ovary per ml; Egami & Taniguchi 1974). The samples (50 μ l) were immediately mixed with Griess reagent (sulphanilamide with N-(1-naphthyl) ethylenediamine/HCl). After 10 min of incubation at room temperature, it was read for absorbance at 540 nm, and nanomoles of nitrite were determined using a standard curve. The assay sensitivity was <2.5 nmol/ml. The intra-assay coefficient of variation for all the assays was <10.0%.

P₄ and A₂ assay

Steroids were measured in duplicate by RIA. The anti-sera to P₄ were kindly provided by Dr R Deis (Laboratorio de Reproducción y Lactancia, Mendoza, Argentina). P₄ was measured in nanogram per milligram ovary per millilitre and assay sensitivity was <5 ng P₄/ml. A₂ was expressed as picogram per milligram of ovarian tissue per millilitre (pg/ml ovary per ml), all against incubation time on the systems. The assay sensitivity was <10 pg A₂/ml. The inter- and intra-assay coefficients of variation in all the assays were <10.0%. The result of each experiment is expressed with respect to the corresponding control group.

Enzymatic activity

The activities of 3 β -HSD and 20 α -HSD were measured according to Kawano *et al.* (1988) with a slight modification. The ovary from each animal was homogenised in 0.7 ml of 0.1 M Tris-HCl, 1 mM EDTA (pH 8) at 0 °C with a glass homogeniser. The homogenates were centrifuged at 105 000 *g* for 60 min. The supernatant fluids were used for the assay of 20 α -HSD activity. The precipitates were rehomogenised with 0.7 ml of 0.25 M sucrose and centrifuged at 800 *g* for 5 min. The supernatants were used as the enzyme solution for the assay of 3 β -HSD activity. The substrates for 3 β -HSD and 20 α -HSD were pregnenolone (5 μ g) and 20 α -hydroxypregn-4-en-3-one (12.5 μ g) respectively. Both enzyme activities were assayed spectrophotometrically, depending on the increase in NADH or NADPH in 1 min at 37 °C, and the values were expressed as mU/mg protein/min. The method of Lowry *et al.* (1951) was used for protein determination with BSA as the standard.

RNA isolation and RT-PCR analysis

Once the ovaries were defrosted, the total RNA was extracted using the TRIzol-reagent method (Invitrogen Life Technologies), following the manufacturer's instructions for the RNA extraction (Chomczynski 1993).

Two micrograms of total RNA were reverse transcribed using a Moloney murine leukaemia virus (MMLV) RT and random primers. First, a prereverse transcription (pre-RT) was carried out, using random primers, by incubating for 10 min at room temperature and then 45 min at 42 °C. The buffer RT, the enzyme MMLV-RT (Invitrogen Life Technologies) and a mixture of dNTPs were added to the total pre-RT product, and it was carried to a final volume of 50 μ l. The mixture was incubated for 60 min at 42 °C, then 15 min at 70 °C and finally the reverse transcriptase activity was inactivated by incubating the tubes in an ice bath.

For amplification of the RT products, the reaction mixture consisted of 10 μ l Go Taq Green reaction buffer, 0.2 mM dNTPs, 0.5 μ M specific oligonucleotide primers and 1.25 U Go Taq DNA polymerase (Promega, Inc.) in a final volume of 50 μ l. Amplification was carried out for 30 cycles using 93 °C for denaturation (1 min), 59 °C for annealing (1 min) and 72 °C for extension (15 min) in an Eppendorf thermal cycler.

Specific primers for 3 β -HSD were 5'-CTGCCTGGTGACAG-GAGCAGG-3' and 5'-GCCAGCACTGCCTTCTCGGCC-3' and for 20 α -HSD were 5'-TTCGAGCAGAACTCATGGCTA-3' and 5'-CAACCAGGTAGAATGCCATCT-3'. Each reaction also included primers (5'-CAAGACTGAGTGGCTGGATGG-3' and 5'-ACTTGAAGGGGAATGAGGAAAA-3') to amplify protein cyclophilin A as an internal control. The predicted sizes of the PCR-amplified products were 489 bp for 3 β -HSD, 440 pb for 20 α -HSD and 293 pb for cyclophilin A. Reaction products were electrophoresed on 1.5% agarose gels, visualised with ethidium bromide (5.5 mg/ml) and examined by u.v. transillumination. Band intensities of RT-PCR products were quantified using ImageJ (NIH, Bethesda, MA, USA; <http://rsb.info.nih.gov/ij/>) and expressed as arbitrary units.

Statistical analysis

All data are presented as the mean $\pm$ S.E.M. in each group of six rats. Differences between two groups were analysed with the Student's *t*-test. ANOVA followed by the Tukey's test was used for between-group comparison. A value of $P < 0.05$ was considered statistically significant (Snedecor & Cochran 1976).

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by Grant 9302 from Universidad Nacional de San Luis, Argentina.

Acknowledgements

We thank Susana Rezzano for English language assistance. We also acknowledge Carlos Pellarin for his technical assistance. This work is part of the doctoral thesis of biochemist Adriana Vega Orozco, funded by Secretaria de Ciencia y Técnica, Universidad Nacional de San Luis.

References

- Aguado LI & Ojeda SR 1984 Prepubertal ovarian function is finely regulated by direct adrenergic influences: role of noradrenergic innervation. *Endocrinology* **114** 1845–1853. (doi:10.1210/endo-114-5-1845)
- Anesetti G, Lombide P & Chavez-Genaro R 2009 Prepubertal exposure modifies neurotrophin receptor expression in celiac neurons and alters ovarian innervation. *Autonomic Neuroscience* **1**–2 35–43. (doi:10.1016/j.autneu.2008.10.021)
- Baljet B & Drukker J 1979 The extrinsic innervation of the abdominal organs in the female rat. *Acta Anatomica* **104** 243–267. (doi:10.1159/000145073)
- Barria A, Leyton V, Ojeda SR & Lara HE 1993 Ovarian steroidal response to gonadotropins and beta-adrenergic stimulation is enhanced in polycystic ovary syndrome: role of sympathetic innervation. *Endocrinology* **133** 2696–2703. (doi:10.1210/en.133.6.2696)
- Berman JR, McCarthy MM & Kyprianou N 1998 Effect of estrogen withdrawal on nitric oxide synthase expression and apoptosis in the rat vagina. *Urology* **51** 650–656. (doi:10.1016/S0090-4295(97)00683-3)
- Blaustein JD 1992 Cytoplasmic estrogen receptors in rat brain immunocytochemical evidence using three antibodies with distinct epitopes. *Endocrinology* **131** 1336–1342. (doi:10.1210/en.131.3.1336)
- Cardinali DP, Vacas MI, Gejman PV, Pisarev MA, Barontin M & Boado RJ 1983 The sympathetic superior cervical ganglia as "little neuroendocrine brains". *Acta Physiologica Latino Americana* **33** 205–221.
- Casais M, Delgado SM, Sosa ZY, Telleria CM & Rastrilla AM 2006 The celiac ganglion modulates LH-induced inhibition of androstenedione release in late pregnant rat ovaries. *Reproductive Biology and Endocrinology* **4** 66–72. (doi:10.1186/1477-7827-4-66)
- Chomczynski P 1993 A reagent for the single-step simultaneous isolation of RNA, DNA and proteins from cell and tissue samples. *Biotechniques* **15** 532–534, 536–537.
- Dalsgaard CJ, Vincent SR, Hökfelt T, Lundberg JM, Dahlström A, Schultzberg M, Dockray GJ & Cuello AC 1982 Coexistence of cholecystokinin- and substance P-like peptides in neurons of the dorsal root ganglia of the rat. *Neuroscience Letters* **33** 159–163. (doi:10.1016/0304-3940(82)90244-0)
- Deecher DC, Swiggard P, Frail DE & O'Connor LT 2003 Characterization of a membrane-associated estrogen receptor in a rat hypothalamic cell line (D12). *Endocrine* **22** 211–223. (doi:10.1385/ENDO:22:3:211)
- Delgado SM, Sosa Z, Dominguez NS, Casais M, Aguado L & Rastrilla AM 2004 Effect of the relation between neural cholinergic action and nitric oxide on ovarian steroidogenesis in prepubertal rats. *Journal of Steroid Biochemistry and Molecular Biology* **91** 139–145. (doi:10.1016/j.jsbmb.2004.04.004)
- Dong YL, Gangula PR, Fang L & Yallampalli C 1999 Nitric oxide reverses prostaglandin-induced inhibition in ovarian progesterone secretion in rats. *Human Reproduction* **14** 27–32. (doi:10.1093/humrep/14.1.27)
- Doolan CM & Harvey BJA 2003 Galphas protein-coupled membrane receptor, distinct from the classical oestrogen receptor, transduces rapid effects of oestradiol on $[Ca^{2+}]_i$ in female rat distal colon. *Molecular and Cellular Endocrinology* **199** 87–103. (doi:10.1016/S0303-7207(02)00303-9)
- Dun SL, Brailoiu GC, Gao X, Brailoiu E, Arterburn JB, Prossnitz ER, Oprea TI & Dun NJ 2009 Expression of estrogen receptor GPR30 in the rat spinal cord and in autonomic and sensory ganglia. *Journal of Neuroscience Research* **87** 1610–1619. (doi:10.1002/jnr.21980)
- Dunnam RC, Hill MJ, Lawson M & Dunbar JC 1999 Ovarian hormone secretory response to gonadotropins and nitric oxide following chronic oxide deficiency in the rat. *Biology of Reproduction* **60** 959–963. (doi:10.1095/biolreprod60.4.959)
- Egami F & Taniguchi S 1974 Nitrate. In, *Methods of Enzymatic Analysis*, 2nd edn, pp 2260–2265. Ed HU Bergmeyer. New York, NY, USA: Academic Press.
- Eisenhofer G, Kopin IJ & Goldstein DS 2004 Catecholamine metabolism: a contemporary view with implications for physiology and medicine. *Pharmacological Reviews* **56** 331–349. (doi:10.1124/pr.56.3.1)
- Eränkő O 1978 Small intensely fluorescent (SIF) cells and nervous transmission in sympathetic ganglia. *Annual Review of Pharmacology and Toxicology* **18** 417–430. (doi:10.1146/annurev.pa.18.040178.002221)
- Fridén B, Runesson E, Hahlin M & Brannstrom M 2000 Evidence for nitric oxide acting as a luteolytic factor in the human corpus luteum. *Molecular Human Reproduction* **6** 397–405. (doi:10.1093/molehr/6.5.397)
- Garcia-Rudaz C, Armando I, Levin G, Escobar ME & Barontini M 1998 Peripheral catecholamine alterations in adolescents with polycystic ovary syndrome. *Clinical Endocrinology* **49** 221–228. (doi:10.1046/j.1365-2265.1998.00511.x)
- Garraza MH, Aguado LI & De Bortoli MA 2004 *In vitro* effect of neuropeptides on ovary or celiac ganglion affects the release of progesterone from ovary in the rat. *Medical Science Monitor* **10** 440–446.
- Heldring N, Pike A, Andersson S, Matthews J, Cheng G, Hartman J, Tujague M, Ström A, Treuter E, Warner M *et al.* 2007 Estrogen receptors: how do they signal and what are their targets. *Physiological Reviews* **87** 905–931. (doi:10.1152/physrev.00026.2006)
- Hsueh AJ, Adashi EY, Jones PBC & Welsh TH Jr 1984 Hormonal regulation of the differentiation of cultured ovarian granulosa cells. *Endocrine Reviews* **5** 76–127. (doi:10.1210/edrv-5-1-76)
- Kawano T, Okamura H, Tajima C, Fukuma K & Katabuchi H 1988 Effect of RU 486 on luteal function in the early pregnant rat. *Journal of Reproduction and Fertility* **83** 279–285. (doi:10.1530/jrf.0.0830279)
- Klein CM & Burden HW 1988a Anatomical localization of afferent and postganglionic sympathetic neurons innervating the rat ovary. *Neuroscience Letters* **85** 217–222. (doi:10.1016/0304-3940(88)90354-0)
- Klein CM & Burden HW 1988b Substance P and vasoactive intestinal polypeptide (VIP) immunoreactive nerve fibers in relation to ovarian postganglionic perikarya in para- and prevertebral ganglia: evidence from combined retrograde tracing and immunocytochemistry. *Cell and Tissue Research* **252** 403–410. (doi:10.1007/BF00214383)
- Koh DS & Hille B 1997 Modulation by neurotransmitters of catecholamine secretion from sympathetic ganglion neurons detected by amperometry. *PNAS* **94** 1506–1511. (doi:10.1073/pnas.94.4.1506)
- Koszykowska M, Calka J, Sz wajca P & Jana B 2011 Long-term estradiol-17 β administration decreases the number of neurons in the caudal mesenteric ganglion innervating the ovary in sexually mature gilts. *Journal of Reproduction and Development* **57** 62–71. (doi:10.1262/jrd.10-0615)
- Kotwica GJ, Rekawiecki R & Duras M 2004 Stimulatory influence of progesterone on its own synthesis in bovine corpus luteum. *Bulletin of the Veterinary Institute in Pulawy* **48** 139–145.

- Kuiper GG, Carlsson B, Grandien K, Enmark E, Häggblad J, Nilsson S & Gustafsson JA 1997 Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors alpha and beta. *Endocrinology* **138** 863–870. (doi:10.1210/en.138.3.863)
- Kuiper GG, Van den Bemd GJ & Van Leeuwen JP 1999 Estrogen receptor and the SERM concept. *Journal of Endocrinological Investigation* **22** 594–603.
- Lara HE, Disson GA, Leyton V, Paredes A, Fuenzalida H, Fiedler JL & Ojeda SR 2000 An increased intraovarian synthesis of nerve growth factor and its low affinity receptor is a principal component of steroid-induced polycystic ovary in the rat. *Endocrinology* **141** 1059–1072. (doi:10.1210/en.141.3.1059)
- Lara HE, Dorfman M, Venegas M, Luza SM, Luna SL, Mayerhofer A, Guimaraes MA, Rosa ESA & Ramirez VD 2002 Changes in sympathetic nerve activity of the mammalian ovary during a normal estrous cycle and in polycystic ovary syndrome: studies on norepinephrine release. *Microscopic Research and Technique* **59** 495–502. (doi:10.1002/jemt.10229)
- Lars-Gösta E, Holmberg K, Emson P, Schemmann M & Hokfelt T 1997 Nitric oxide synthase, choline acetyltransferase, catecholamine enzymes and neuropeptides and their colocalization in the anterior pelvic ganglion and hypogastric nerve of the male guinea pig. *Anatomical Record* **35** 68–76.
- Lawrence IE Jr & Burden HW 1980 The origin of the extrinsic adrenergic innervation to the ovary. *Anatomical Record* **196** 51–59. (doi:10.1002/ar.1091960106)
- Levin ER 2001 Cell localization, physiology, and nongenomic actions of estrogen receptors. *Journal of Applied Physiology* **91** 1860–1867.
- Lowry OH, Rosebrough HJ, Farr AL & Randall RJ 1951 Protein measurement with the folin phenol reagent. *Journal of Biological Chemistry* **193** 265–275.
- Mannisto PT & Kaakkola S 1999 Catechol-O-methyltransferase (COMT): biochemistry, molecular biology, pharmacology, and clinical efficacy of the new selective COMT inhibitors. *Pharmacological Reviews* **51** 593–628.
- Mao J, Duan RW, Zhong L, Gibori G & Azhar S 1997 Expression, purification and characterization of the rat luteal 20 α -hydroxysteroid dehydrogenase. *Endocrinology* **1** 182–190. (doi:10.1210/en.138.1.182)
- Miszkiel G, Skarzynski D, Bogacki M & Kotwica J 1999 Concentrations of catecholamines, ascorbic acid, progesterone and oxytocin in the corpora lutea of cyclic and pregnant cattle. *Reproduction, Nutrition, Development* **39** 509–516. (doi:10.1051/rnd:19990410)
- Miyamoto A, Brückmann A, von Lütow H & Schams D 1993 Multiple effects of neuropeptide Y, substance P and vasoactive intestinal polypeptide on progesterone and oxytocin release from bovine corpus luteum *in vitro*. *Journal of Endocrinology* **138** 451–458. (doi:10.1677/joe.0.1380451)
- Mosselman S, Polman J & Dijkema R 1996 ERbeta: identification and characterization of a novel human estrogen receptor. *FEBS Letters* **392** 49–53. (doi:10.1016/0014-5793(96)00782-X)
- Motta AB, Estevez A, Tognetti T, Gimeno MA & Franchi AM 2001 Dual effects of nitric oxide in functional and regressing rat corpus luteum. *Molecular Human Reproduction* **7** 43–47. (doi:10.1093/molehr/7.1.43)
- Navarro Despaigne D 2001 Moduladores selectivos del receptor estrógeno. Su utilidad en la mujer posmenopáusica. *Rev. Cubana. Endocrinología* **12** 124–127.
- Olson L, Jones BCh & Jablonka-Shariff A 1996 Nitric oxide decreases estradiol synthesis of rats luteinized ovarian cells: possible role for nitric oxide in functional luteal regression. *Endocrinology* **137** 3531–3536. (doi:10.1210/en.137.8.3531)
- Paech K, Webb P, Kuiper GG, Nilsson S, Gustafsson J, Kushner PJ & Scanlan TS 1997 Differential ligand activation of estrogen receptors ERalpha and ERbeta at AP1 sites. *Science* **277** 1508–1510. (doi:10.1126/science.277.5331.1508)
- Paige LA, Christensen DJ, Grøn H, Norris JD, Gottlin EB, Padilla KM, Chang CY, Ballas LM, Hamilton PT, McDonnell DP *et al.* 1999 Estrogen receptor (ER) modulators each induce distinct conformational changes in ER alpha and ER beta. *PNAS* **96** 3999–4004. (doi:10.1073/pnas.96.7.3999)
- Papka RE, Estorey-Workley M, Shughrue PJ, Merchenthaler I, Kollins JJ, Usip S, Saunders PT & Shupnik M 2001 Estrogen receptor-alpha and beta-immunoreactivity and mRNA in neurons of sensory and autonomic ganglia and spinal cord. *Cell and Tissue Research* **2** 193–214. (doi:10.1007/s004410100363)
- Poole T 1999 UFAW hand book of the care and management of Laboratory animals, vol 2, 7th edn. Oxford, UK: Blackwell Publishing.
- Prud'Homme MJ, Houdeau E, Serghini R, Tillet Y, Schemmann M & Rousseau JP 1999 Small intensely fluorescent cells of the rat paracervical ganglion synthesize adrenaline, receive afferent innervation from postganglionic cholinergic neurones, and contain muscarinic receptors. *Brain Research* **821** 141–149. (doi:10.1016/S0006-8993(99)01094-X)
- Quinson N, Robbins HL, Clark MJ & Furness JB 2001 Locations and innervation of cell bodies of sympathetic neurons projecting to the gastrointestinal tract in the rat. *Archives of Histology and Cytology* **64** 281–294. (doi:10.1679/aohc.64.281)
- Razandi M, Pedram A, Merchenthaler I, Greene GL & Levin ER 2004 Plasma membrane estrogen receptors exist and functions as dimers. *Molecular Endocrinology* **18** 2854–2865. (doi:10.1210/me.2004-0115)
- Rekawiecki R, Nowik M & Kotwica J 2005 Stimulatory effect of LH, PGE₂ and progesterone on StAR protein, cytochrome P450 cholesterol side chain cleavage and 3 β -hydroxysteroid dehydrogenase gene expression in bovine luteal cells. *Prostaglandins & Other Lipid Mediators* **78** 169–184. (doi:10.1016/j.prostaglandins.2005.06.009)
- Rekawiecki R, Kowalik MK, Slonina D & Kotwica J 2008 Regulation of progesterone synthesis and action in bovine corpus luteum. *Journal of Physiology and Pharmacology* **59** 75–89.
- Rosa ESA, Guimaraes MA, Padmanabhan V & Lara HE 2003 Prepubertal administration of estradiol valerate disrupts cyclicity and leads to cystic ovarian morphology during adult life in the rat: role of sympathetic innervation. *Endocrinology* **144** 4289–4297. (doi:10.1210/en.2003-0146)
- Shinohara Y, Matsumoto A & Mori T 1998 Effects of prenatal exposure to diethylstilbestrol on the sympathetic nervous system in the rat ovary. *Neuroscience Letters* **255** 123–126. (doi:10.1016/S0304-3940(98)00681-8)
- Shinohara Y, Matsumoto A, Hayashi S & Mori T 2000 Prenatal exposure to diethylstilbestrol decreases the number of estrogen receptor alpha-containing neurons innervating the ovary in rat celiac ganglion. *Neuroscience* **101** 779–783. (doi:10.1016/S0306-4522(00)00395-X)
- Singh M, Sétáló G, Guan X, Wuaren M & Toran-Allerand CD 1999 Estrogen-induced activation of mitogen-activated protein kinase in cerebral cortical explants: convergence of estrogen and neurotrophin signalling pathways. *Journal of Neuroscience* **4** 1179–1188.
- Snedecor WG & Cochran WG 1976 *Statistical Methods*, Ames, Iowa: The Iowa State University Press.
- Snyder GD, Holmes RW, Bates JN & Van Voorhis BJ 1996 Nitric oxide inhibits aromatase activity: mechanisms of action. *Journal of Steroid Biochemistry and Molecular Biology* **58** 63–69. (doi:10.1016/0960-0760(96)00008-8)
- Sohrabji F, Greene LA, Miranda RC & Toran-Allerand CD 1994a Reciprocal regulation of estrogen and NGF receptors by their ligands in PC12 cells. *Journal of Neurobiology* **25** 974–988. (doi:10.1002/neu.480250807)
- Sohrabji F, Miranda RC & Toran-Allerand CD 1994b Estrogen differentially regulates estrogen and nerve growth factor receptor mRNAs in adult sensory neurons. *Journal of Neuroscience* **14** 459–471.
- Song RX & Santen RJ 2006 Membrane initiated estrogen signaling in breast cancer. *Biology of Reproduction* **75** 9–16. (doi:10.1095/biolreprod.105.050070)
- Song RX, Zhang Z & Santen RJ 2005 Estrogen rapid action via protein complex formation involving ERalpha and Src. *Trends in Endocrinology and Metabolism* **16** 347–353. (doi:10.1016/j.tem.2005.06.010)
- Sosa ZY, Casais M, Rastrilla AM & Aguado LI 2000 Adrenergic influences on coeliac ganglion affect the release of progesterone from cycling ovaries. Characterisation of an *in vitro* system. *Journal of Endocrinology* **166** 307–318. (doi:10.1677/joe.0.1660307)
- Sosa Z, Delgado SM, Casais M, Aguado L & Rastrilla AM 2004 Release of ovarian progesterone during the rat oestrous cycle by ganglionic cholinergic influence. The role of norepinephrine. *Journal of Steroid Biochemistry and Molecular Biology* **91** 179–184. (doi:10.1016/j.jsmb.2004.03.119)

- Stefano G, Cadet P, Mantione K, Cho J, Jones D & Zhu W** 2003 Estrogen signalling at the cell surface coupled to nitric oxide release in *Mytilus edulis* nervous system. *Endocrinology* **144** 1234–1240. (doi:10.1210/en.2002-220967)
- Stocco CO & Deis RP** 1996 Luteolytic effect of LH: inhibition of 3β -hydroxysteroid dehydrogenase and stimulation of 20α -hydroxysteroid dehydrogenase luteal activities in late pregnant rats. *Journal of Endocrinology* **150** 423–429. (doi:10.1677/joe.0.1500423)
- Sun L, Wen L, Shao X, Qian H, Jin Y, Liu W & Fu Z** 2010 Screening of chemicals with anti-estrogenic activity using *in vitro* and *in vivo* vitellogenin induction responses in zebrafish (*Danio rerio*). *Chemosphere* **78** 793–799. (doi:10.1016/j.chemosphere.2009.12.020)
- Tellería CM, Stocco CO, Stati AO, Rastrilla AM, Carrizo DG, Aguado LI & Deis RP** 1995 Dual regulation of luteal progesterone production by androstenedione during spontaneous and RU486-induced luteolysis in pregnant rats. *Journal of Steroid Biochemistry and Molecular Biology* **55** 385–393. (doi:10.1016/0960-0760(95)00190-5)
- Vega Orozco A, Sosa Z, Fillipa V, Mohamed F & Rastrilla AM** 2006 The cholinergic influence on the mesenteric ganglion affects the liberation of ovarian steroids and nitric oxide in oestrus day rats: characterization of an *ex vivo* system. *Journal of Endocrinology* **191** 587–598. (doi:10.1677/joe.1.06859)
- Vega Orozco A, Sosa Z, Delgado S, Casais M & Rastrilla AM** 2010 Involvement of ganglionic cholinergic receptors on the steroidogenesis in the luteal phase in rat. *Journal of Steroid Biochemistry and Molecular Biology* **120** 45–52. (doi:10.1016/j.jsbmb.2010.03.040)
- Wang Z, Zhang X, Shen P, Loggie BW, Chang Y & Deuel TF** 2005 Identification, cloning, expression of human estrogen receptor- α 36 a novel variant of human estrogen receptor- α 66. *Biochemical and Biophysical Research Communications* **366** 1023–1027. (doi:10.1016/j.bbrc.2005.08.226)
- Warner M & Gustafsson JA** 2006 Nongenomic effects of estrogen: why all the uncertainty? *Steroids* **71** 91–96. (doi:10.1016/j.steroids.2005.09.001)
- Webb P, Lopez GN, Uht RM & Kushner PJ** 1995 Tamoxifen activation of the estrogen receptor/AP-1 pathway: potential origin for the cell-specific estrogen-like effects of antiestrogens. *Molecular Endocrinology* **9** 443–456. (doi:10.1210/me.9.4.443)
- Weihua Z, Andersson S, Cheng G, Simpson ER, Warner M & Gustafsson JA** 2003 Update on estrogen signaling. *FEBS Letters* **546** 17–24. (doi:10.1016/S0014-5793(03)00436-8)
- Yamamoto R, Wada A, Asada Y, Niina H & Sumiyoshi A** 1993 N omega-nitro-larginine, an inhibitor of nitric oxide synthesis, decreases noradrenaline outflow in rat isolated perfused mesenteric vasculature. *Naunyn-Schmiedeberg's Archives of Pharmacology* **347** 238–240. (doi:10.1007/BF00169274)
- Zou A, Marschke KB, Arnold KE, Berger EM, Fitzgerald P, Mais DE & Allegretto EA** 1999 Estrogen receptor beta activates the human retinoic acid receptor α -1 promoter in response to tamoxifen and other estrogen receptor antagonists, but not in response to estrogen. *Molecular Endocrinology* **13** 418–430. (doi:10.1210/me.13.3.418)
- Zoubina EV & Smith PG** 2002 Distributions of estrogen receptors α and β in sympathetic neurons of female rats: enriched expression by uterine innervation. *Journal of Urology* **169** 382–385. (doi:10.1016/S0022-5347(05)64132-8)
- Zoubina EV & Smith PG** 2003 Expression of oestrogen receptors α and β by sympathetic ganglion neurons projecting to the proximal urethra of female rats. *Journal of Urology* **1** 382–385.

Received 10 March 2011

First decision 7 April 2011

Revised manuscript received 6 November 2011

Accepted 11 November 2011