Distribution of GFRA1-expressing spermatogonia in adult mouse testis

Margherita Grasso, Andrea Fuso¹, Lisa Dovere, Dirk G de Rooij^{2,3}, Mario Stefanini, Carla Boitani and Elena Vicini

Fondazione Pasteur Cenci Bolognetti, Section of Histology and Medical Embryology, Department of Anatomical, Histological, Forensic and Orthopaedic Sciences, 'La Sapienza' University of Rome, Via Antonio Scarpa 14, 00161 Rome, Italy, ¹Department of Surgery, 'La Sapienza' University of Rome, Rome, Italy, ²Department of Endocrinology, Faculty of Science, Utrecht University, Utrecht, The Netherlands and ³Center for Reproductive Medicine, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

Correspondence should be addressed to E Vicini; Email: elena.vicini@uniroma1.it

Abstract

In mice and other mammals, spermatogenesis is maintained by spermatogonial stem cells (SSCs), a cell population belonging to undifferentiated type A spermatogonia. In the accepted model of SSC self-renewal, Asingle (As) spermatogonia are the stem cells, whereas paired (Apaired (Apr)) and chained (Aaligned (Aal)) undifferentiated spermatogonia are committed to differentiation. This model has been recently challenged by evidence that As and chained (Apr and Aal), undifferentiated spermatogonia are heterogeneous in terms of gene expression and function. The expression profile of several markers, such as GFRA1 (the GDNF co-receptor), is heterogeneous among As, Apr and Aal spermatogonia. In this study, we have analysed and quantified the distribution of GFRA1-expressing cells within the different stages of the seminiferous epithelial cycle. We show that in all stages, GFRA1 + chained spermatogonia (Apr to Aal) are more numerous than GFRA1 + As spermatogonia. Numbers of chained GFRA1 + spermatogonia are sharply reduced in stages VII–VIII when Aal differentiate into A1 spermatogonia. GFRA1 expression is regulated by GDNF and in cultures of isolated seminiferous tubules, we found that GDNF expression and secretion by Sertoli cells is stage-dependent, being maximal in stages II–VI and decreasing thereafter. Using qRT-PCR analysis, we found that GDNF regulates the expression of genes such as *Tex14*, *Sohlh1* and *Kit* (*c-Kit*) known to be involved in spermatogonial differentiation. Expression of *Kit* was upregulated by GDNF in a stage-specific manner. Our data indicate that GDNF, besides its crucial role in the self-renewal of stem cells also functions in the differentiation of chained undifferentiated spermatogonia.

Reproduction (2012) **143** 325–332

Introduction

In mice and other mammals, spermatogenesis is maintained by spermatogonial stem cells (SSCs), a subpopulation of the undifferentiated type A spermatogonia. These undifferentiated spermatogonia can be found as single cells (Asingle (As)) or as chains of two (Apaired (Apr)) or up to 16–32 cells (Aaligned (Aal)) connected by intercellular bridges. According to the 'As model', As spermatogonia are the stem cells of spermatogenesis, whereas longer clones of cells are committed to differentiation (Huckins 1971b, Oakberg 1971, Lok *et al.* 1982). The As either self-renew by forming single cells or generate pairs of cells (Apr), which are connected by an intercellular bridge as a result of incomplete cytokinesis. Therefore, As always undergo symmetric cell division, and during the steady state, the ratio between self-renewal and differentiation is close to 1 (de Rooij 2001).

Spermatogenesis is a cyclic process that in the mouse is divided into 12 stages, each representing a unique

association of germ cells at different steps of differentiation. The relationship between spermatogenic stages and the kinetics of undifferentiated spermatogonia proliferation has been analysed in different mammalian species (Phillips *et al.* 2010). In the mouse, morphometric analysis performed on whole mounts of seminiferous tubules indicates that As and Apr numbers remain relatively constant throughout the stages of spermatogenesis while the density of Aals fluctuates cyclically (Tegelenbosch & de Rooij 1993). All classes of undifferentiated spermatogonia randomly divide primarily in stages X–II, and subsequently, during stages VII and VIII, almost all of the larger chains (Aal4–Aal32) differentiate into A1 spermatogonia (Schrans-Stassen *et al.* 1999). This event marks the transition from the randomly cycling population of undifferentiated spermatogonia to the rigidly controlled series of mitotic divisions of differentiating spermatogonia, which are highly synchronised. All subsequent divisions of A1, A2, A3, A4, In, B spermatogonia and primary spermatocytes

occur in a stage-specific fashion (de Rooij 2001). The transition between undifferentiated and differentiating spermatogonia is a finely regulated step that is marked by a reduction in cell cycle length (Huckins 1971a, 1971b, Lok & de Rooij 1983, Lok *et al.* 1983) and a gene expression profile switch (Schrans-Stassen *et al.* 1999, Beumer *et al.* 2000). Undifferentiated spermatogonia express the transcriptional repressor ZBTB16 (PLZF) while its expression is downregulated thereafter (Buaas *et al.* 2004, Costoya *et al.* 2004). Initiation of the expression of KIT (c-KIT) marks the transition from Aal to A1 spermatogonia (Schrans-Stassen *et al.* 1999).

The 'As model' has recently been challenged by evidence, from our and other laboratories, which indicates that As and chained undifferentiated spermatogonia are heterogeneous in terms of gene expression and function. The expression profile of CDH1, LIN28, GFRA1, NANOS2, NANOS3 and NEUROG3 (NGN3) is heterogeneous among As, Apr and Aal spermatogonia (Grisanti *et al.* 2009, Suzuki *et al.* 2009, Zheng *et al.* 2009, Nakagawa *et al.* 2010); moreover, different subsets of As cells may be generated through an asymmetric division (Grisanti *et al.* 2009, Suzuki *et al.* 2009). Data from transgenic animal models were interpreted to mean that two functionally distinct stem cell compartments exist: the actual and the potential stem cell compartments, which may play different roles in different conditions. Furthermore, *in vivo* time-lapse observations suggest that occasionally longer clones of Aal spermatogonia undergo fragmentation, opening the possibility that pinched cells become new As or Apr cells (Nakagawa *et al.* 2010). Altogether, these studies encourage further studies into the validity of the 'As model'. Furthermore, elucidation of the lineage relationship within undifferentiated spermatogonia compartments deserves further investigation.

To gain more insight into the lineage relationship of different subsets of undifferentiated spermatogonia, we have analysed and quantified the distribution of GFRA1-expressing cells within the subsequent stages of the seminiferous epithelial cycle. GFRA1 is the co-receptor for *gdnf*, a Sertoli cell-derived factor that controls the balance between self-renewal and differentiation of SSCs (Meng *et al.* 2000). The distribution of GFRA1-expressing cells, along with the gene expression profiles of candidate genes after GDNF *in vitro* treatment of intact expressing seminiferous tubules, suggests that, apart from regulating stem cell self-renewal, surprisingly GDNF also stimulates cells to progress along the differentiation pathway.

Results

Distribution of GFRA1+ A spermatogonia during the seminiferous epithelial cycle

To analyse the expression pattern of GFRA1 within undifferentiated spermatogonia, intact seminiferous tubules were co-stained for GFRA1 and ZBTB16,

a marker of all types of undifferentiated spermatogonia (Buaas *et al.* 2004, Costoya *et al.* 2004). Figure 1 shows the frequency of GFRA1+ and GFRA1- clones for each class of undifferentiated spermatogonia obtained by morphometric analysis of mixed seminiferous tubules. In agreement with published data, our results show that the frequency of GFRA1+ spermatogonia decreases as spermatogonia progress from As to Aal (Nakagawa *et al.* 2010). In addition to single cells and chains of two, four, eight and 16 cells, a few groups of three ZBTB16+/GFRA1+ or GFRA1- cells were also observed. As no chains of six cells were observed during this study, we assumed that the groups of three were composed of a single and a pair, and they were counted as such. Possibly, these groups of three descended from false pairs, i.e. singles that did not yet migrate away from each other while one of the cells had already divided into a pair.

Next, we identified and quantified the numbers of GFRA1+ and GFRA1- cells in three groups of stages of the seminiferous epithelial cycle: stages II–VI, VII–VIII and IX–XI. Figure 2 (A–C) shows representative images of GFRA1 and ZBTB16-stained tubules for each group. The total number of ZBTB16+ cells was similar in stages II–VI and VII–VIII but significantly decreased in stages IX–XI ($P < 0.05$). Likewise, total numbers of GFRA1- undifferentiated spermatogonia were also similar in stages II–VI and VII–VIII but significantly decreased in stages IX–XI ($P < 0.05$). Conversely, the total number of GFRA1+ cells was similar in stages II–VI and VII–VIII but significantly increased in stages IX–XI ($P < 0.05$; Fig. 3A). To further investigate the distribution of each population of undifferentiated spermatogonia, we counted clones of As, Apr and Aal (4–8) expressing GFRA1, in each group of stages (Fig. 3B). Statistical analysis showed that the number of GFRA1+ As and Apr clones did not significantly differ among the groups of stages. However, GFRA1+ clones composed of 4 or 8 were relatively numerous in stages IX–XI, decreased in

Figure 1 Frequency of spermatogonia clones positive or negative for GFRA1 expression. The number of clones examined in three different experiments is shown above each column.

Figure 2 The distribution of GFRA1-expressing undifferentiated spermatogonia in the different stages of seminiferous epithelium. (A–C) Intact seminiferous tubules isolated from adult testis and staged were stained with anti-ZBTB16 (green), anti-GFRA1 (red) and TO-PRO-3 (blue). Pictures illustrate merged Z sections. Representative pictures for each group of stages are shown: II–VI (A), VII–VIII (B) and IX–XI (C). Solid arrows, As spermatogonia; empty arrows, Apr spermatogonia; arrowheads, Aal4; asterisk, Aal8 spermatogonia. Bar, 20 μ m.

stages II–VI and further reduced in stages VII–VIII ($P < 0.05$). Even though clones of ZBTB16 + Aal8 were found in all stages (Supplementary Figure 1, see section on supplementary data given at the end of this paper), clones of GFRA1 + Aal8 were only found in stages IX–XI. The large numbers of GFRA1 + chains of 4 and 8 cause the significant increase of GFRA1 + cells in stages IX–XI compared with the other stages (Fig. 3A). To further understand this point, for each group of stages, we calculated the numbers of As cells and the total number of cells in chains and expressed them as a percentage (single vs chained cells; Fig. 3C). Results show that in each group of stages, the number of positive cells in chains (Apr to Aal4–Aal8) outnumbered positive single cells (As), with a maximum in stages IX–XI (81% of chained cells vs 19% of As; Fig. 3C).

GFRA1 expression and secretion during the cycle of the seminiferous epithelium

GFRA1 is the specific co-receptor for GDNF. In the seminiferous epithelium GDNF is produced and secreted by Sertoli cells, a population of non-dividing somatic cells (Tadokoro *et al.* 2002). We addressed the question whether specific changes in *gdnf* expression occurred during the seminiferous epithelial cycle. Using real-time PCR we found that GDNF mRNA expression was higher in stages II–VI compared with stages VII–VIII ($P < 0.05$) and with stages IX–XI ($P < 0.01$) while there was no difference between stages VII–VIII and stages IX–XI (Fig. 4A).

To determine whether mRNA levels correlated to secreted protein levels, staged seminiferous tubules were maintained in culture for 24 h and GDNF released into the culture medium was measured by ELISA. As shown in Fig. 4B, secretion of GDNF was stage-dependent and paralleled mRNA levels, being maximal at stages II–VI.

GDNF regulates genes expressed by spermatogonia

Following the observations that i) GFRA1-expressing spermatogonia appear more frequently in chains than as single cells (As) and ii) GDNF levels are maximal at

stages II–VI when undifferentiated spermatogonia are arrested in G0–G1 and are going to differentiate into A1 spermatogonia (Schrans-Stassen *et al.* 1999), we hypothesised that GDNF may control genes involved in spermatogonial differentiation. We analysed the expression level of known GDNF target genes, as well as genes involved in spermatogonial differentiation, by qRT-PCR after *in vitro* GDNF treatment of intact seminiferous tubules.

Microarray analysis, performed on SSC lines, showed that GDNF upregulates several genes, including *Bcl6b* and, intriguingly, its own receptor *Gfra1*; in contrast, GDNF downregulates *Neurog3* (Oatley *et al.* 2006). We asked whether similar regulation by GDNF takes place

Figure 3 Morphometric analysis of GFRA1-labelled cell distribution in the different stages of the seminiferous epithelium. (A) Quantification of ZBTB16+, GFRA1+ and GFRA1– spermatogonia expressed as number of cells/1000 Sertoli cells. (B) Clone numbers of As, Apr, Aal4 and Aal8 for each group of stages. (C) Percentage of single cells (As) vs cells in chains (Apr–Aal) in each group of stages. Results are expressed as the mean $\pm$ S.E.M.; $n = 3$ experiments. a Significant difference vs d and b; f significant difference vs d and e; i significant difference vs g and h; l significant difference vs j and k ($P < 0.05$). #Indicates that no Aal8 clones were present in stages II–VIII. Aal8 clones were present only in stages IX–XI.

Figure 4 *Gdnf* expression in different groups of stages. (A) Real-time PCR analysis of GDNF mRNA in staged, freshly isolated seminiferous tubules. Data are expressed as arbitrary units normalised to β -actin $\pm$ s.e.m. Results are from $n=3$ experiments measured in duplicate. (B) ELISA quantification of GDNF released into medium by staged seminiferous tubules after 24 h of *in vitro* culture. Results are expressed as the mean $\pm$ s.e.m.; $n=4$ experiments. * $P<0.05$ and ** $P<0.01$.

in intact seminiferous tubules. Mixed seminiferous tubules from adult mice were cultured in the absence or presence, of increasing GDNF concentrations (from 0 to 100 ng/ml), and gene expression was assessed by qRT-PCR analysis (Fig. 5A). The results indicate that GDNF upregulates *Gfra1* and *Bcl6b* mRNA levels and downregulates *Neurog3* mRNA levels in a dose-dependent manner (Fig. 5A).

Next, we investigated the impact of GDNF treatment on the expression level of genes involved in undifferentiated spermatogonia renewal/differentiation, specifically *Tex14*, *Sohlh1*, *Kit* and *Pou5f1* (*Oct4*). Expression levels were measured in seminiferous tubules that were untreated or treated with 100 ng/ml GDNF. *Bcl6b*, *Gfra1* and *Neurog3* were included as internal controls (Fig. 5B). Interestingly, GDNF is able to upregulate *Tex14*, *Sohlh1* and *Kit* mRNA levels and to downregulate *Pou5f1* mRNA levels compared with controls (Fig. 4B). Because initiation of *Kit* expression marks the differentiation of Aal to A1 spermatogonia (Schrans-Stassen *et al.* 1999), we analysed the impact of GDNF treatment on *Kit* expression levels in the different stages of the seminiferous epithelial cycle. In untreated seminiferous tubules, *Kit* mRNA is significantly higher

in stages VII–VIII compared with other stages ($P<0.05$). GDNF treatment affects *Kit* mRNA in a stage-dependent manner, *Kit* is upregulated in stages II–VIII ($P<0.05$) but is unaffected in stages IX–XI (Fig. 5C). We next asked whether GDNF treatment altered the expression of target genes at the protein level. To this end, mixed seminiferous tubules from adult mice were cultured in the absence, or presence, of 100 ng/ml GDNF and protein extracts were analysed by Western blotting. Neither *Kit* nor *Sohlh1* was found to be upregulated in GDNF-treated samples compared with controls (Supplementary Figure 2, see section on supplementary data given at the end of this article).

Discussion

In this study, we have shown that the number of undifferentiated spermatogonia, identified by the specific marker ZBTB16, changes during the cycle of the seminiferous epithelium, being lower in stages IX–XI, compared with other stages. In contrast, in stages IX–XI, the number of ZBTB16+/GFRA1+ cells was significantly higher than in other stages. The analysis of clone distribution revealed that while the numbers of clones of 1 and 2 GFRA1+ spermatogonia remain constant in all stages, clones of 4 and 8 GFRA1+ spermatogonia are differentially distributed with a significant decrease of GFRA1+ Aal4 in stages VII–VIII and relatively large numbers of GFRA1+ Aal4 and Aal8 clones in stages IX–XI. This further supports the concept of heterogeneity of gene expression among undifferentiated spermatogonia during the epithelial cycle (Grisanti *et al.* 2009, Nakagawa *et al.* 2010).

Interestingly, levels of GDNF mRNA and secreted protein also display a stage-specific profile, being maximal at stages II–VI and reaching a minimum at stages IX–XI. These data are in line with a stage-dependent expression of GDNF in the rat seminiferous tubules (Johnston *et al.* 2011). The finding that GDNF levels are high when the mitotic activity of undifferentiated spermatogonia is low (from stages II–III until stage X (de Rooij 2001)) suggests that GDNF might have a role in the differentiation of GFRA1+ Aal. In line with this, the numbers of clones of GFRA1+ Aal4 were markedly reduced and GFRA1+ Aal8 were even absent in stages VII–VIII. Since under physiological conditions apoptosis takes place in differentiating spermatogonia but not in undifferentiated spermatogonia (de Rooij & Janssen 1987, de Rooij & Lok 1987) we interpret this as a consequence of the progression of GFRA1+ Aal clones into the next differentiation compartment. In addition, we found that *Kit* mRNA levels were significantly upregulated by GDNF during stages II–VIII. Since *Kit* expression marks the transition of undifferentiated to differentiating type A spermatogonia (Schrans-Stassen *et al.* 1999), our observation supports the idea that undifferentiated spermatogonial differentiation into A1

Figure 5 Regulation of spermatogonia-expressed genes by GDNF. (A) Real-time PCR analysis of *Gfra1*, *Neurog3* and *Bcl6b* mRNAs in total seminiferous tubules treated with different concentrations of GDNF. $n=6$, $**P<0.01$ vs control (GDNF, 0 ng/ml; one-way ANOVA, Bonferroni *post-hoc*). (B) Real-time PCR analysis of *Tex14*, *Sohlh1*, *Kit*, *Pou5f1*, *Neurog3*, *Gfra1* and *Bcl6b* mRNAs in seminiferous tubules treated, or untreated, with GDNF 100 ng/ml. $n=3$, $*P<0.05$ and $**P<0.01$ vs control (GDNF, 0 ng/ml; *t*-test). (C) Real-time PCR analysis of *Kit* mRNA in different stages of seminiferous tubules treated, or untreated, with GDNF, 100 ng/ml $n=3$, $*P<0.05$ and $**P<0.01$. In A and B, data are expressed as 'fold increase' with control indicated as $1 \pm \text{S.E.M.}$; in C data are expressed as arbitrary units normalised to β -actin $\pm \text{S.E.M.}$.

spermatogonia might be facilitated by GDNF. An additional factor, involved in translational regulation, will be necessary to induce the actual synthesis of the KIT protein.

A list of GDNF-regulated genes has been generated through microarray analysis performed on freshly

isolated mouse pup GFRA1-expressing type A spermatogonia or mouse SSC lines (Hofmann *et al.* 2005, Oatley *et al.* 2006). For instance, GDNF mediates the upregulation of several transcription factor-encoding genes (*N-Myc*, *Egr2*, *Egr3*, *Bcl6b*, *Lhx1*, etc.) and intriguingly *Gfra1*. In contrast, several genes were found to be downregulated by GDNF including the differentiation marker *Neurog3*. Our cultures of isolated seminiferous tubules, revealed a similar upregulation of *Gfra1*, *Bcl6b* and a downregulation of *Neurog3*. Interestingly, we observed a GDNF-dependent upregulation of *Tex14*, *Sohlh1* and *Kit* and concomitant downregulation of *Pou5f1* mRNAs. *TEX14* is an essential component of germ cell intercellular bridges. In its absence, spermatogenesis is disrupted due to enhanced apoptosis of spermatocytes (Greenbaum *et al.* 2006). Inactivation of *SOHLH1*, a basic helix–loop–helix transcription factor that is expressed in germ cells, causes infertility by blocking differentiation of spermatogonia into spermatocytes (Ballow *et al.* 2006). *Pou5f1* is a homeobox transcription factor required for the maintenance of totipotency of embryonic stem cells (Nichols *et al.* 1998, Niwa *et al.* 2000), is expressed in undifferentiated spermatogonia (Pesce *et al.* 1998) and is required for SSC maintenance in culture and for colonisation activity following testis germ cell transplantation (Dann *et al.* 2008). In the microarray analysis performed on mouse SSC lines *Tex14*, *Sohlh1*, *Kit* and *Pou5f1* mRNA levels were unaffected by *gdnf* treatment (data accessible at NCBI GEO datasets, accession GDS2112 (Oatley *et al.* 2006). This discrepancy may well be related to the different culture conditions of the GDNF target cells in the two experimental models. However, since conditions of target cells in intact seminiferous tubules are more similar to the *in vivo* situation, it is likely that our data closely reflect the effect of GDNF *in vivo*. In future experiments, it will be interesting to perform a microarray analysis on cultures of intact seminiferous tubules to gain a more comprehensive list of GDNF-regulated genes. We attempted to analyse whether upregulation of mRNA levels of our candidate genes results in increased protein levels. By western blot analysis of KIT and SOHLH1 we could not detect any significant difference between control and GDNF-treated seminiferous tubules. However, since GDNF target cells in the adult seminiferous tubules represent a very small population compared with all the other cell types, it is likely that an increase in protein level in target cells is hard to detect by Western blot analysis.

GDNF is critical for self-renewal maintenance of SSCs, both *in vivo* and *in vitro*. In the mouse, disruption of one GDNF allele leads to spermatogonial depletion and the appearance of Sertoli-cell-only tubules in the adult. In contrast, overexpression of GDNF causes an accumulation of clusters of undifferentiated spermatogonia (Meng *et al.* 2000). GDNF is also required for the initiation and maintenance of SSCs in

culture (Kanatsu-Shinohara *et al.* 2003, Kubota *et al.* 2004). In this study, we show that GDNF also induces the transcription of genes involved in the differentiation of Aal into A1 spermatogonia. This differentiation becomes visible during stages VII and VIII, but the start of the process may be much earlier, for example in stages II–III when these cells stop proliferating and become quiescent. Our results show that the co-receptor for GDNF, GFRA1, is expressed in Aal spermatogonia in stages IX–XI and that the expression of this receptor decreases to below the immunohistochemical detection level thereafter. Maybe the high levels of GDNF during stages II–VI can compensate for the low levels of GFRA1 in Aal spermatogonia and still regulate transcription levels in these cells. Interestingly, in cultures of rat spermatogonia, GDNF was also shown to stimulate development of Aal (Hamra *et al.* 2007). How can GDNF impact both self-renewal and differentiation programmes in target cells? GDNF signal transduction in germ cells is elaborate and may lead to the activation of many signalling pathways, including the SRC family of tyrosine kinase, phosphatidylinositol-3-kinase/AKT (PI3K/AKT) and mitogen-activated protein kinase (MAPK) (Braydich-Stolle *et al.* 2007, Lee *et al.* 2007, Oatley *et al.* 2007, He *et al.* 2008). Work in other models has shown that GDNF acts through two different mechanisms, *in cis* or *in trans*, leading to the activation of different pathways. During activation *in cis*, the coreceptor RET interacts with the GFRA1–GDNF complex inside lipid rafts. However, *in trans* RET binds a soluble GFRA1–GDNF complex outside lipid rafts (Airaksinen & Saarma 2002). In addition, GDNF may function in a RET-independent fashion (Sariola & Saarma 2003). Therefore, GDNF may activate different transduction pathways in As vs chained cells eliciting divergent patterns of gene expression.

In conclusion, our data suggest that GDNF is not only involved in the regulation of the self-renewal of stem/progenitor cells, but also in further progression of chained undifferentiated spermatogonia.

Materials and Methods

Animals

C57BL/6 mice were used in all experiments. Animals were housed in a standard animal facility with free access to food and water, in accordance with the guidelines for animal care at the Sapienza University of Rome. All procedures were approved by the Department of Health Animal Care and Use Committee.

Reagents and antibodies

L-Glutamine, non-essential amino acids, HEPES, gentamicin, donkey serum and foetal bovine serum were acquired from Sigma–Aldrich. TO-PRO-3 fluorescent dye and TRIzol were acquired from Gibco, Invitrogen. Recombinant rat GDNF was

acquired from R&D Systems (Minneapolis, MN, USA). Transcriptor Reverse Transcriptase was acquired from Roche Applied Science (Milano, Italy). FluoCycle SybrGreen Kit was acquired from Euroclone (Milano, Italy). The antibodies used were goat anti-rat GFRA1 (Neuromics, Herford, Germany), mouse anti-human ZBTB16 (2A9; Calbiochem, Milano, Italy), Cy3-conjugated donkey anti-goat and FITC-conjugated donkey anti-mouse (Jackson Laboratories, Suffolk, UK). Vectashield mounting medium was acquired from Vector (Milano, Italy).

Preparation of stage-specific seminiferous tubules

Adult mice were killed, and the testes were removed and placed in a Petri dish containing PBS. The testes were decapsulated and transferred to a new Petri dish containing PBS. The seminiferous tubules were viewed on a trans-illuminating dissection microscope. Using forceps, the tubules were gently pulled apart, avoiding damaging them by squeezing or shaking. The stages of the seminiferous epithelial cycle were identified by the light absorption pattern (Parvinen & Vanha-Perttula 1972) and the tubules were subdivided into three groups: stages II–VI (the strong spot), stages VII–VIII (the dark zone) and stages IX–XI (the pale zone) as described previously (Morena *et al.* 1995). Seminiferous tubules were either fixed for whole-mount immunofluorescence experiments or cultured for quantitative real-time PCR, ELISA and Western blot experiments as described later.

Whole-mount immunofluorescence

Whole-mount immunofluorescence on adult seminiferous tubules was performed as described previously (Corallini *et al.* 2006). Briefly, both mixed tubules and each group of stages were fixed in 4% v/v paraformaldehyde at 4 °C for 3 h. Tubules were incubated overnight at 4 °C with mouse anti-ZBTB16 and goat anti-GFRA1 antibodies (1:100) followed by an incubation with species-specific secondary antibodies conjugated to FITC and Cy3 for 1 h at room temperature (RT). Nuclear staining was performed with TO-PRO-3 (0.2 mg/ml). Tubules were mounted with Vectashield medium and observed under a confocal microscope. Confocal photomicrographs of double-immunostained tubules were acquired with a Leica TCS SP2 (Leica, Milano, Italy) equipped with three laser lines (argon 488, 543 and 633 nm). Each channel was acquired separately using specific laser lines to avoid bleed-through of the fluorochromes. Photomicrographs were made using LAS AF Software (Leica). Stacks of longitudinal Z sections, representing different layers of a tubule, were acquired.

Quantification of type As, Apr and Aal spermatogonia in whole-mount immunofluorescence

Stained tubules from three different experiments were observed using a confocal microscope with a 40× oil immersion objective. Cell counts were performed on stored microphotographs using LAS AF Software by placing a frame of 36 100 µm² on the domed centre of the tubule. The total number of ZBTB16+ spermatogonia, including type As, Apr and Aal spermatogonia, their corresponding GFRA1 staining

status and Sertoli cell nuclei were counted within the grid on randomly selected tubules. A ZBTB16+ cell was described as type As if surrounding cells within 25 µm lacked ZBTB16 staining and exhibited a dissimilar nuclear morphology; furthermore, doublets of cells were designated type Apr when the internuclear distance was below 25 µm (Tegelenbosch & de Rooij 1993, Grisanti *et al.* 2009). In three different experiments, we analysed and counted 56 frames for stages II–VI corresponding to 737 ZBTB16+ cells, 44 for stages VII–VIII corresponding to 564 ZBTB16+ cells and 102 for stages IX–XI corresponding to 931 ZBTB16+ cells. The number of cells (single and chained) found in each group of stages were expressed per 1000 Sertoli cells because these cells do not proliferate in the adult testis and their number does not vary among stages.

Semi-quantitative real-time PCR

Thirty millimetres of each group of stages or total tubules were cultured *in vitro* for 16 h in 24-well culture plates with DMEM supplemented with antibiotics, L-glutamine, non-essential amino acids, HEPES and gentamicin. Afterwards, tubules were untreated or treated for 6 h with GDNF at 100 ng/ml. To quantify relative expression levels of selected genes, total RNA from the tubules was purified using TRIzol reagent and quantified. Gene expression was evaluated by real-time PCR analysis. Total RNA/sample (1 µg) was used for cDNA synthesis with Transcriptor Reverse Transcriptase. The resulting cDNA was used for real-time PCR, where analysis was performed in triplicate for each sample. cDNA was mixed with 0.2 µM forward and reverse primers and 10 µl 2× master mix to a final reaction volume of 20 µl. Reactions were performed on Opticon2 DNA Engine (MJ Research, Waltham, MA, USA). Primers used to detect target genes were designed and evaluated using the Primer3 Software (<http://frodo.wi.mit.edu/primer3>) and ordered from Sigma-Genosys Oligo Synthesis (Supplementary Table 1, see section on supplementary data given at the end of this article). The amplification efficiency for each primer pair was determined by amplification of a linear standard curve of total cDNA (from 0.32 to 200 ng for *Gfra1*, *Bcl6b*, *Neurog3*, *Tex14*, *Sohlh1* and *Pou5f1*; from 1.6 to 1000 ng for *Kit*). All primer pairs showed good amplification linearity and efficiency (>95%). Total cDNA levels were normalised against a β-actin control and presented either as the fold increase over the basal control or as arbitrary units. Expression levels of interest genes were also normalised against two other reference genes, *Gapdh* and *Rn18s* (18S), giving similar results (data not shown). Each real-time PCR assay was repeated at least two times in three different experiments and each sample was analysed in triplicate in each assay; mean values with S.E.M. were used for comparison.

GDNF secretion

Hundred millimetres of pieces of seminiferous tubules of each group of stages were seeded in 24-well culture plates with DMEM supplemented with antibiotics, L-glutamine, non-essential amino acids, HEPES pH 7.7, gentamicin and 100 ng/ml FSH. GDNF production in the culture medium

was measured at 24 h, using the GDNF Emax ImmunoAssay System Kit (Promega, Milano, Italy) according to the manufacturer's instructions. The ELISA reaction was carried out in 96-well plates, and the optical densities were recorded by the Microplate reader at 450 nm wavelength. Results are from four independent experiments and are expressed as means ± S.E.M.

Statistical analysis

Data were analysed to define the significance of the differences between the two groups using a *t*-test or, to compare many groups, using a one-way ANOVA followed by a *post-hoc* Bonferroni test or Dunn test.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/REP-11-0385>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This study was supported by the Ministero Istruzione Università Ricerca (#20083CBLPY_002), Agenzia Spaziale Italiana (#I/065/08/0) and Sapienza University (#C26V10RPR2).

Acknowledgements

We are grateful to Stefania Fera and Stefania De Grossi for their excellent technical assistance.

References

- Airaksinen MS & Saarma M 2002 The GDNF family: signalling, biological functions and therapeutic value. *Nature Reviews. Neuroscience* **3** 383–394. (doi:10.1038/nrn812)
- Ballow D, Meistrich ML, Matzuk M & Rajkovic A 2006 *Sohlh1* is essential for spermatogonial differentiation. *Developmental Biology* **294** 161–167. (doi:10.1016/j.ydbio.2006.02.027)
- Beumer TL, Ropers-Gajadien HL, Gademan IS, Lock TM, Kal HB & de Rooij DG 2000 Apoptosis regulation in the testis: involvement of Bcl-2 family members. *Molecular Reproduction and Development* **56** 353–359. (doi:10.1002/1098-2795(200007)56:3<353::AID-MRD4>3.0.CO;2-3)
- Braydich-Stolle L, Kostereva N, Dym M & Hofmann MC 2007 Role of Src family kinases and N-Myc in spermatogonial stem cell proliferation. *Developmental Biology* **304** 34–45. (doi:10.1016/j.ydbio.2006.12.013)
- Buaas FW, Kirsh AL, Sharma M, McLean DJ, Morris JL, Griswold MD, de Rooij DG & Braun RE 2004 Plzf is required in adult male germ cells for stem cell self-renewal. *Nature Genetics* **36** 647–652. (doi:10.1038/ng1366)
- Corallini S, Fera S, Grisanti L, Falcatori I, Muciaccia B, Stefanini M & Vicini E 2006 Expression of the adaptor protein m-Numb in mouse male germ cells. *Reproduction* **132** 887–897. (doi:10.1530/REP-06-0062)
- Costoya JA, Hobbs RM, Barna M, Cattoretti G, Manova K, Sukhwani M, Orwig KE, Wolgemuth DJ & Pandolfi PP 2004 Essential role of Plzf in maintenance of spermatogonial stem cells. *Nature Genetics* **36** 653–659. (doi:10.1038/ng1367)

- Dann CT, Alvarado AL, Molyneux LA, Denard BS, Garbers DL & Porteus MH 2008 Spermatogonial stem cell self-renewal requires OCT4, a factor downregulated during retinoic acid-induced differentiation. *Stem Cells* **26** 2928–2937. (doi:10.1634/stemcells.2008-0134)
- Greenbaum MP, Yan W, Wu MH, Lin YN, Agno JE, Sharma M, Braun RE, Rajkovic A & Matzuk MM 2006 TEX14 is essential for intercellular bridges and fertility in male mice. *PNAS* **103** 4982–4987. (doi:10.1073/pnas.0505123103)
- Grisanti L, Falcatori I, Grasso M, Doveire L, Fera S, Muciaccia B, Fuso A, Berno V, Boitani C, Stefanini M *et al.* 2009 Identification of spermatogonial stem cell subsets by morphological analysis and prospective isolation. *Stem Cells* **27** 3043–3052. (doi:10.1002/stem.206)
- Hamra FK, Chapman KM, Nguyen D & Garbers DL 2007 Identification of neuregulin as a factor required for formation of aligned spermatogonia. *Journal of Biological Chemistry* **282** 721–730. (doi:10.1074/jbc.M608398200)
- He ZP, Jiang JJ, Kokkinaki M, Golestaneh N, Hofmann MC & Dym M 2008 GDNF upregulates c-fos transcription via the Ras/ERK1/2 pathway to promote mouse spermatogonial stem cell proliferation. *Stem Cells* **26** 266–278. (doi:10.1634/stemcells.2007-0436)
- Hofmann MC, Braydich-Stolle L & Dym M 2005 Isolation of male germ-line stem cells; influence of GDNF. *Developmental Biology* **279** 114–124. (doi:10.1016/j.ydbio.2004.12.006)
- Huckins C 1971a Cell cycle properties of differentiating spermatogonia in adult Sprague-Dawley rats. *Cell and Tissue Kinetics* **4** 139–154.
- Huckins C 1971b The spermatogonial stem cell population in adult rats. I. Their morphology, proliferation and maturation. *Anatomical Record* **169** 533–557. (doi:10.1002/ar.1091690306)
- Johnston DS, Olivas E, Dicandloro P & Wright WW 2011 Stage-specific changes in GDNF expression by rat Sertoli cells: a possible regulator of the replication and differentiation of stem spermatogonia. *Biology of Reproduction* **85** 763–769. (doi:10.1095/biolreprod.110.087676)
- Kanatsu-Shinohara M, Ogonuki N, Inoue K, Miki H, Ogura A, Toyokuni S & Shinohara T 2003 Long-term proliferation in culture and germline transmission of mouse male germline stem cells. *Biology of Reproduction* **69** 612–616. (doi:10.1095/biolreprod.103.017012)
- Kubota H, Avarbock MR & Brinster RL 2004 Culture conditions and single growth factors affect fate determination of mouse spermatogonial stem cells. *Biology of Reproduction* **71** 722–731. (doi:10.1095/biolreprod.104.029207)
- Lee J, Kanatsu-Shinohara M, Inoue K, Ogonuki N, Miki H, Toyokuni S, Kimura T, Nakano T, Ogura A & Shinohara T 2007 Akt mediates self-renewal division of mouse spermatogonial stem cells. *Development* **134** 1853–1859. (doi:10.1242/dev.003004)
- Lok D & de Rooij DG 1983 Spermatogonial multiplication in the Chinese hamster. I. Cell cycle properties and synchronization of differentiating spermatogonia. *Cell and Tissue Kinetics* **16** 7–18.
- Lok D, Weenk D & de Rooij DG 1982 Morphology, proliferation, and differentiation of undifferentiated spermatogonia in the Chinese hamster and the rat. *Anatomical Record* **203** 83–99. (doi:10.1002/ar.1092030109)
- Lok D, Jansen MT & de Rooij DG 1983 Spermatogonial multiplication in the Chinese hamster. II. Cell cycle properties of undifferentiated spermatogonia. *Cell and Tissue Kinetics* **16** 19–29.
- Meng X, Lindahl M, Hyvonen ME, Parvinen M, de Rooij DG, Hess MW, Raatikainen-Ahokas A, Sainio K, Rauvala H, Lakso M *et al.* 2000 Regulation of cell fate decision of undifferentiated spermatogonia by GDNF. *Science* **287** 1489–1493. (doi:10.1126/science.287.5457.1489)
- Morena AR, Boitani C, de Grossi S, Stefanini M & Conti M 1995 Stage and cell-specific expression of the adenosine 3',5' monophosphate-phosphodiesterase genes in the rat seminiferous epithelium. *Endocrinology* **136** 687–695. (doi:10.1210/en.136.2.687)
- Nakagawa T, Sharma M, Nabeshima Y, Braun RE & Yoshida S 2010 Functional hierarchy and reversibility within the murine spermatogenic stem cell compartment. *Science* **328** 62–67. (doi:10.1126/science.1182868)
- Nichols J, Zevnik B, Anastasiadis K, Niwa H, Klewe-Nebenius D, Chambers I, Scholer H & Smith A 1998 Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. *Cell* **95** 379–391. (doi:10.1016/S0092-8674(00)81769-9)
- Niwa H, Miyazaki J & Smith AG 2000 Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nature Genetics* **24** 372–376. (doi:10.1038/74199)
- Oakberg EF 1971 Spermatogonial stem-cell renewal in the mouse. *Anatomical Record* **169** 515–531. (doi:10.1002/ar.1091690305)
- Oatley JM, Avarbock MR, Telaranta AI, Fearon DT & Brinster RL 2006 Identifying genes important for spermatogonial stem cell self-renewal and survival. *PNAS* **103** 9524–9529. (doi:10.1073/pnas.0603332103)
- Oatley JM, Avarbock MR & Brinster RL 2007 Glial cell line-derived neurotrophic factor regulation of genes essential for self-renewal of mouse spermatogonial stem cells is dependent on src family kinase signaling. *Journal of Biological Chemistry* **282** 25842–25851. (doi:10.1074/jbc.M703474200)
- Parvinen M & Vanha-Perttula T 1972 Identification and enzyme quantitation of the stages of the seminiferous epithelial wave in the rat. *Anatomical Record* **174** 435–449. (doi:10.1002/ar.1091740404)
- Pesce M, Wang XY, Wolgemuth DJ & Scholer H 1998 Differential expression of the Oct-4 transcription factor during mouse germ cell differentiation. *Mechanisms of Development* **71** 89–98. (doi:10.1016/S0925-4773(98)00002-1)
- Phillips BT, Gassei K & Orwig KE 2010 Spermatogonial stem cell regulation and spermatogenesis. *Philosophical Transactions of the Royal Society of London. Series B, Biological sciences* **365** 1663–1678. (doi:10.1098/rstb.2010.0026)
- de Rooij DG 2001 Proliferation and differentiation of spermatogonial stem cells. *Reproduction* **121** 347–354. (doi:10.1530/rep.0.1210347)
- de Rooij DG & Janssen JM 1987 Regulation of density of spermatogonia in the seminiferous epithelium of the Chinese hamster: I. Undifferentiated spermatogonia. *Anatomical Record* **217** 124–130. (doi:10.1002/ar.1092170203)
- de Rooij DG & Lok D 1987 Regulation of density of spermatogonia in the seminiferous epithelium of the Chinese hamster: II. Differentiating spermatogonia. *Anatomical Record* **217** 131–136. (doi:10.1002/ar.1092170204)
- Sariola H & Saarma M 2003 Novel functions and signalling pathways for GDNF. *Journal of Cell Science* **116** 3855–3862. (doi:10.1242/jcs.00786)
- Schrans-Stassen BH, van de Kant HJ, de Rooij DG & van Pelt AM 1999 Differential expression of c-kit in mouse undifferentiated and differentiating type A spermatogonia. *Endocrinology* **140** 5894–5900. (doi:10.1210/en.140.12.5894)
- Suzuki H, Sada A, Yoshida S & Saga Y 2009 The heterogeneity of spermatogonia is revealed by their topology and expression of marker proteins including the germ cell-specific proteins Nanos2 and Nanos3. *Developmental Biology* **336** 222–231. (doi:10.1016/j.ydbio.2009.10.002)
- Tadokoro Y, Yomogida K, Ohta H, Tohda A & Nishimune Y 2002 Homeostatic regulation of germinal stem cell proliferation by the GDNF/FSH pathway. *Mechanisms of Development* **113** 29–39. (doi:10.1016/S0925-4773(02)00004-7)
- Tegelenbosch RA & de Rooij DG 1993 A quantitative study of spermatogonial multiplication and stem cell renewal in the C3H/101 F1 hybrid mouse. *Mutation Research* **290** 193–200. (doi:10.1016/0027-5107(93)90159-D)
- Zheng K, Wu X, Kaestner KH & Wang PJ 2009 The pluripotency factor LIN28 marks undifferentiated spermatogonia in mouse. *BMC Developmental Biology* **9** 38. (doi:10.1186/1471-213X-9-38)

Received 12 October 2011

First decision 31 October 2011

Accepted 5 December 2011