Connecting *cis*-elements and *trans*-factors with mechanisms of developmental regulation of mRNA translation in meiotic and haploid mammalian spermatogenic cells

Kenneth C Kleene

Department of Biology, University of Massachusetts Boston, 100 Morrissey Boulevard, Boston, Massachusetts 02125-3393, USA

Correspondence should be addressed to K C Kleene; Email: kenneth.kleene@umb.edu

Abstract

mRNA-specific regulation of translational activity plays major roles in directing the development of meiotic and haploid spermatogenic cells in mammals. Although many RNA-binding proteins (RBPs) have been implicated in normal translational control and sperm development, little is known about the keystone of the mechanisms: the interactions of RBPs and microRNAs with *cis*-elements in mRNA targets. The problems in connecting factors and elements with translational control originate in the enormous complexity of post-transcriptional regulation in mammalian cells. This creates confusion as to whether factors have direct or indirect and large or small effects on the translation of specific mRNAs. This review argues that gene knockouts, heterologous systems, and overexpression of factors cannot provide convincing answers to these questions. As a result, the mechanisms involving well-studied mRNAs (*Ddx4/Mvh*, *Prm1*, *Prm2*, and *Sycp3*) and factors (DICER1, CPEB1, DAZL, DDX4/MVH, DDX25/GRTH, translin, and ELAV1/HuR) are incompletely understood. By comparison, mutations in elements can be used to define the importance of specific pathways in regulating individual mRNAs. However, few elements have been studied, because the only reliable system to analyze mutations in elements, transgenic mice, is considered impractical. This review describes advances that may facilitate identification of the direct targets of RBPs and analysis of mutations in *cis*-elements. The importance of upstream reading frames in the developmental regulation of mRNA translation in spermatogenic cells is also documented.

Reproduction (2013) **146** R1–R19

Introduction

Spermatogenesis, the process of cell renewal and differentiation that produces male gametes, is divided into three main phases, each with distinct cell type and function, shown in Fig. 1 (Russell *et al.* 1990). The first is the mitotic phase in which diploid spermatogonia proliferate in the mitotic cycle, greatly amplifying the number of cells. This is followed by the meiotic phase in which two copies of the diploid chromosome set in spermatocytes are shuffled and reduced to a haploid set. During the final phase, spermiogenesis, haploid spermatids differentiate into highly polar cells specialized for motility and fertilization, spermatozoa. Spermatogenesis is extremely complex and occurs in many developmental stages (Fig. 1). Spermatocytes pass through six stages, distinguished by the morphology of the meiotic chromosomes, and mouse spermatids pass through 16 stages, known as steps, distinguished by the morphology of the developing acrosome, nucleus, and sperm tail (Russell *et al.* 1990).

Translational regulation plays an important role in controlling the development of spermatocytes and spermatids (reviewed in Paronetto & Sette (2010),

Nguyen-Chi & Morello (2011) and Idler & Yan (2012)). The most familiar example originates in chromatin remodeling in which histones are replaced by transition proteins and protamines during late spermiogenesis (reviewed in Meistrich *et al.* (2003)). Round spermatids in steps 1–8 have round transcriptionally active nuclei. The nuclei of elongating spermatids in steps 9 through early 12 change from round to elongated accompanied by a drastic reduction in transcriptional activity (Kierszenbaum & Tres 1975). As a result, elongated spermatids in steps 12–16 differentiate without new mRNA transcripts. Many mRNAs are transcribed in spermatocytes and round spermatids and stored in a translationally inactive state in the absence of the encoded protein for several days to more than 2 weeks, before the protein is first detected in elongating and elongated spermatids (Fig. 1). In addition, translational control directs the beginning and end of the first meiotic prophase, and all mRNAs in pachytene spermatocytes and round spermatids are partially repressed by an unknown global mechanism (discussed below).

mRNA-specific translational regulation in eukaryotic cells is usually mediated by *cis*-elements that bind

trans-factors, RNA-binding proteins (RBPs), or small non-coding RNAs, sncRNAs (Jackson *et al.* 2010). By binding elements, factors promote or block the recruitment of other factors and ribosomes to mRNAs, thereby decreasing, negative control, or stimulating, positive control, the translation of mRNA-coding sequences into protein. However, one type of *cis*-element discussed below, upstream reading frames, does not necessarily involve the binding of factors to elements.

The importance of translational regulation in spermatogenesis is further demonstrated by mutations in a large number of genes, mostly knockout models, which cause abnormal sperm development, reductions in male fertility, and failure to express proteins at the normal developmental stage (Table 1). Although many reviews discussing these and related subjects cited here affirm translational control as an important problem in

spermatogenesis, little is known about the basic features of the mechanisms: the interactions of factors with elements and the magnitude of the change in translational activity mediated by factors and elements. This review focuses on systems in which elements and factors have and have not been connected, strategies that may facilitate the identification and validation of elements, and important controversies to be resolved in future work.

Elucidating mechanisms of translational control with gene knockouts, overexpression of factors, heterologous systems, and mutations in elements

The challenges in connecting mRNA targets with regulatory factors can be traced to a major insight: post-transcriptional regulation of gene expression in

Figure 1 Developmental expression of mRNAs and proteins during spermatogenesis. Spermatogenesis begins with primary germ cells (PGCs) *in utero* and continues with mitotic, meiotic, and haploid phases after birth. The stages of meiosis are denoted PL, preleptotene; L, leptotene; Z, zygotene; P, pachytene; D, diplotene; and M, meiotic divisions 1 and 2. Spermiogenesis is subdivided into three main phases in mice: round spermatids, steps 1–8; elongating spermatids, steps 9–early 12; and elongated spermatids, steps 12–16. The diagrams of spermatogenic cells display the gross morphology, and the black nuclei symbolize the highly condensed chromatin of elongated spermatids and spermatozoa (adapted from Russell *et al.* (1990)). The decrease in transcriptional activity in elongating spermatids is based on the study of Kierszenbaum & Tres (1975). The duration of each phase of meiosis and steps of spermiogenesis have been rounded to the nearest number of days (Russell *et al.* 1990). The stages of expression of mRNAs and proteins are depicted respectively by blue and brown lines. The thickness of the lines symbolizes the levels of expression, and the arrows on the left ends of the lines depicting the expression of DAZL and DDX4/MVH indicate that both proteins are expressed *in utero*. The sources of information concerning the stages of expression of mRNAs and proteins are as follows: *Dbil5* and *Pgk2* mRNAs and proteins, cited in the text; *Prm1*, *Prm2*, and *Tnp1* mRNAs and proteins (Mali *et al.* 1989, Meistrich *et al.* 2003); *Smcp* mRNA and protein (Shih & Kleene 1992, Cataldo *et al.* 1996); *DDX4/MVH* (Onohara *et al.* 2010); *GRTH/DDX25* (Onohara & Yokota 2012); *DAZL* (Brook *et al.* 2009); *SYCP3* (Cohen *et al.* 2006); *YBX2/MSY2* and *YBX3/MSY4* (Oko *et al.* 1996, Davies *et al.* 2000, Giorgini *et al.* 2001); *ELAVL1/HuR* (Nguyen Chi *et al.* 2009); and *KHDRBS1/SAM68* (Paronetto *et al.* 2009).

eukaryotic cells is much more complex than anyone could have predicted 10 years ago (reviewed in Bartel (2009), Kishore *et al.* (2010) and Morris *et al.* (2010)). Post-transcriptional gene regulation in mammalian cells is realized at the levels of alternative splicing,

polyadenylation site selection, mRNA export, translation, stability, and localization, and individual mRNAs are often regulated at multiple levels. These processes are mediated by the activities of RBPs, and sncRNAs, the best studied of which are microRNAs,

Table 1 Factors implicated in translational control in spermatocytes and spermatids.

Factor ^a	Knockout phenotype ^b	mRNA targets (evidence) ^c	Positive/negative ^d	Reference ^e
ADAR1/TENR (dsRBD)	Infertile, oligospermia, spermiation, nuclear shape, flagellum, motility	Unknown	Unknown	Connolly <i>et al.</i> (2005)
Adenosine deaminase				
Boll/Boule (ssRNA, RRM)	Infertile, azoospermia, total block step 6, acrosome	Unknown	Pos?	VanGompel & Xu (2010)
CELF3/BRUNOL1 (ssRNA, RRM)	Fertile, mild oligospermia, no morphological defects	Unknown	?	Dev <i>et al.</i> (2007)
CELF1/CUGBP1 (ssRNA, RRM)	Normal to infertile, step 7 block, apoptotic round spermatids	Unknown	?	Kress <i>et al.</i> (2007)
CPEB1 (ssRNA, RRM, and ZFD)	Subfertile, oligospermia, pachytene apoptosis, partial block leptotene–zygotene?, chromatin fragmentation, motility	<i>Sycp3</i> (IG, UV) <i>Ddx4/Mvh?</i>	Pos and Neg?	Tay & Richter (2001)
DAZL (ssRNA, RRM)	Infertile, azoospermia, total block leptotene–zygotene	<i>Sycp3</i> , <i>Ddx4/Mvh</i> , <i>Cdc25c</i> (<i>Cdc25</i>) (IG, IP, UV)	Pos	Saunders <i>et al.</i> (2003) and Reynolds <i>et al.</i> (2005, 2007)
DDX4/MVH (ssRNA?, DEAD-box RNA helicase)	Infertile, azoospermia, total block near leptotene–zygotene	Unknown	Pos?	Tanaka <i>et al.</i> (2000)
DDX25/GRTH (DEAD-box RNA helicase)	Infertile, azoospermia, massive meiotic metaphase I apoptosis, total block step 8, nuclear elongation	<i>Prm1</i> , <i>Prm2</i> , <i>Tnp1</i> , and <i>Tnp2</i> histone <i>H4</i> , many others (IG, IP, MP)	Pos	Tsai-Morris <i>et al.</i> (2004, 2012)
EIF4G3 (binds EIF4E and PABPC?)	Infertile, azoospermia, total block before metaphase I	<i>Hsp2a</i> (IG)	Pos	Sun <i>et al.</i> (2010)
ELAVL1/HuR (ssRNA, RRM)	Infertile, azoospermia, apoptosis? at or before meiotic division, nuclear elongation	<i>Hsp2a</i> (IG), many	Pos	Chi <i>et al.</i> (2011)
miRNA469 (microRNA)	Not studied	<i>Tnp2</i> , <i>Prm2</i> (IG in 3T3 cells)	Neg	Dai <i>et al.</i> (2011)
YBX2/MSY2 (ssRNA, CSD)	Infertile, azoospermia, minor pachytene and massive step 8 apoptosis, nuclear elongation and condensation, flagella, spermiation, mRNA degradation	Unknown	Neg	Yang <i>et al.</i> (2007)
YBX3/MSY4/MSY3L (ssRNA, CSD)	Infertile or fertile (line dependent), pachytene apoptosis, no morphological defects	Unknown	Neg	Lu <i>et al.</i> (2006)
PABPC1 (ssRNA, RRM)	Not studied	All?	Neg and Pos	Yanagiya <i>et al.</i> (2010)
PAIP2a (binds PABPC1)	Infertile, severe oligospermia, round spermatid apoptosis, nuclear elongation flagella, spermiation	<i>Prm1</i> , <i>Prm2</i> , <i>Tnp1</i> (IG)	Pos and Neg	Yanagiya <i>et al.</i> (2010)
KHDRBS1/SAM68 (ssRNA, KH)	Infertile, azoospermia or severe oligospermia, pachytene and round spermatid apoptosis, flagella, nuclei	<i>Spag16</i> , <i>Nedd1</i> , <i>Spdy</i> (IG, IP)	Pos	Paronetto <i>et al.</i> (2009)
STRBP/SPNR (dsRNA, dsRBP)	Subfertile, oligospermia, 'degenerated cells throughout', flagella	Unknown	?	Pires-daSilva <i>et al.</i> (2001)
TARBP2/PRBP (dsRNA, dsRBP)	Infertile, severe oligospermia, round spermatid apoptosis, spermiation, nuclear elongation and condensation	<i>Prm1</i> , <i>Prm2</i> (IG, indirect?)	Pos	Zhong <i>et al.</i> (1999)
PAPβ/TPAP (unknown)	Infertile, azoospermia, total step 8 block pachytene apoptosis, step 7 block	Unknown	Pos?	Kashiwabara <i>et al.</i> (2002)
TSN/translin (ssRNA)	Fertile, mild oligospermia, pachy apoptosis, no morphological defects	<i>Prm2</i> , <i>Gapdhs?</i> , many? (RRL)	Neg?	Chennathuzukuzhi <i>et al.</i> (2003)

^aThe official name of the protein is according to the Mouse Genome Informatics website and common synonyms. The information in parentheses lists the preference for binding single-stranded or double-stranded RNA and the type of RNA-binding domain. The RNA-binding domains include RRM, RNA recognition motif; CSD, cold shock domain; KH, KH-domain; dsRBP, double-stranded RNA-binding domain; ZFD, zinc-finger domain. ^bIt is difficult to summarize the phenotypes of the knockouts accurately within the space limitations of a table, since the phenotypes are influenced by genetic background and exhibit different effects on the differentiation of various organelles and biases due to differences in the depth of analysis. The descriptions of the phenotypes in the primary literature have been reproduced above, but question marks have been inserted when the developmental abnormalities have not been described precisely. For knockouts that cause partial blocks, the organelles or developmental processes that exhibit abnormalities are listed. ^cThe mRNAs listed are proposed to be direct targets of the knocked-out factor. The information in parentheses describes the evidence: IG, *in vivo* genetic effects based on gene knockouts; u.v., u.v. cross-linking assays; IP, co-immunoprecipitation; IPPM, immunoprecipitation of polysomes and microarray analysis; RRL, rabbit reticulocyte. ^dInferences that a factor activates mRNA translation, positive control, or represses mRNA translation, negative control, based on the effects of knockouts. ^eThe references of the articles containing supporting information.

miRNAs. It is estimated that the mammalian genome encodes about 400 miRNAs and 600 RBPs recognized by various types of RNA-binding domains (RBDs) (Bartel 2009, Ascano *et al.* 2012). However, the number of RBPs is potentially much greater because the genome encodes an additional ~1400 uncharacterized proteins with domains that can bind RNA and/or DNA (Morris *et al.* 2010, Ascano *et al.* 2012). In addition, polyadenylated mRNAs in somatic cells bind several hundred proteins that lack recognizable RBDs (Baltz *et al.* 2012, Castello *et al.* 2012).

The current paradigm is that individual RBPs and miRNAs interact with many, sometimes thousands of, mRNAs and that the expression of individual mRNAs is regulated cooperatively by multiple RBPs and miRNAs (Bartel 2009, Kishore *et al.* 2010, Morris *et al.* 2010). However, the functions of RBPs and miRNAs in post-transcriptional regulation in somatic mammalian cells differ significantly. miRNAs base-pair with target mRNAs and guide mRNA degradation, translational repression, or both, but the decrease in the expression of individual proteins is usually slight, 'fine-tuning' (Bartel 2009, Fabian & Sonenberg 2012). In contrast, RBPs increase or decrease mRNA degradation and translation, by magnitudes ranging from small to large.

Another striking difference between RBPs and miRNAs is that RBPs often coordinate the expression of proteins with related functions accompanying the differentiation of specific cell types or changes in the physiological state (Kishore *et al.* 2010, Morris *et al.* 2010). By comparison, miRNAs usually fine-tune the expression of large numbers of proteins with unrelated functions (Bartel 2009, Fabian & Sonenberg 2012) and more rarely the expression of proteins with closely related functions (Mestdagh *et al.* 2010).

About 20 factors have been implicated in translational regulation by gene knockouts, most of which have deleterious effects on sperm development (see Table 1). Pinpointing the defects in post-transcriptional regulation that result from knockouts in spermatogenic cells is remarkably difficult, because of problems in distinguishing between mRNA targets that are regulated directly and indirectly by a factor. Specifically, many factors regulate groups of target mRNAs directly, but some of the direct mRNA targets encode factors that regulate secondary targets, creating confusion as to whether the effect of the knockout is direct or indirect. This problem is magnified by the interactions of factors with very large numbers of mRNAs and functions in multiple post-transcriptional processes.

The problems in relating phenotypic effects to direct and indirect targets are illustrated by two RBPs with important functions in spermatogenesis, YBX2/MSY2 and ELAVL1/HuR (Yang *et al.* 2007, Chi *et al.* 2011). YBX2/MSY2 belongs to the family of Y-box proteins that interacts with thousands of, and potentially all, mRNAs in somatic cells and functions in transcription, splicing,

positive and negative regulation of translation, stability, and mRNP structure (Skabkin *et al.* 2006). Similarly, ELAVL1/HuR binds U-rich elements in the 3' UTRs and introns of thousands of mRNAs in cultured mammalian cells, enhancing stability and translation by mechanisms that are sometimes coupled to pre-mRNA processing (Lebedeva *et al.* 2011, Mukherjee *et al.* 2011). As some of the most responsive mRNAs to siRNA knockdown of ELAVL1/HuR encode proteins with functions in regulating gene expression, ELAVL1/HuR has been described as 'a regulator of regulators' (Mukherjee *et al.* 2011).

Distinguishing between the direct and indirect effects of knockouts is especially difficult in spermatogenic cells, because RBPs that are necessary for spermatogenesis are usually expressed for more than a week, a period during which the cells undergo profound developmental changes (Fig. 1). In fact, the earliest phenotype of knockouts is often increased apoptosis, followed days later by morphological abnormalities (examples are shown in Table 1). These compound phenotypes could be produced by defective regulation of two sets of genes, early and late, with independent or dependent functions. Obviously, a knockout that blocks sperm development before the stage at which an mRNA is normally activated or repressed in wild-type mice precludes inferring that the knocked-out factor directly regulates that mRNA.

Knockouts often produce different abnormalities in different cells at the same stage in a single testis (Zhong *et al.* 1999, Dass *et al.* 2007, Reynolds *et al.* 2007) or different phenotypes on different genetic backgrounds (Connolly *et al.* 2005, Reynolds & Cooke 2005). These phenotypes are the hallmarks of incomplete and variable expressivity and penetrance, indicative of modifying factors that compensate for or intensify the phenotype created by the absence of a factor.

Studies of the overexpression of factors are also difficult to interpret because elevated levels of a factor can produce regulatory effects that differ from the effects of normal levels. This is clearly illustrated by findings that high levels of PABPC1 repress mRNA translation in the Krebs cell-free translation lysate and round spermatids, in contrast to the usual function of lower levels of PABPC1 in stimulating translation (Yanagiya *et al.* 2010). Similarly, depletion of ELAV1/HuR with a conditional knockout decreases *Hspa2* mRNA translation in late spermatocytes (Chi *et al.* 2011), in agreement with a well-established function of this RBP in translational activation in somatic cells (Lebedeva *et al.* 2011). In contrast, the overexpression of ELAV1/HuR in spermatogenic cells in transgenic mice also decreases *Hspa2* mRNA translation, consistent with direct or indirect functions in translational repression (Chi *et al.* 2011). In addition, the overexpression of Y-box protein YBX3L/MSY4 in round and elongated spermatids produces deformed spermatozoa (Giorgini *et al.* 2002), whereas the *Ybx3/My4* knockout does not (Lu *et al.* 2006).

The functions of factors in translational regulation in spermatogenic cells have also been inferred from studies in cultured somatic cells, *Xenopus* oocytes and the rabbit reticulocyte cell-free lysate (RRL). Heterologous systems can produce interpretable results when the interaction between a factor and an element is sufficient to regulate translation, sometimes with the assistance of factors conveniently present in the heterologous cells (Ling *et al.* 2002, Collier *et al.* 2005). However, this approach is particularly risky with spermatogenic cells, which express many factors that differ qualitatively and quantitatively from those in somatic cells and oocytes (references in Kleene (2003), Nguyen-Chi & Morello (2011) and Idler & Yan (2012)). In some cases, the addition of a factor at a superphysiological level may alter translational activity only at excessive levels (see above) or the activity of the factor may depend on post-translational modifications that are not fulfilled in the heterologous milieu. There are several reports of misleading results in the RRL (Kleene & Smith 1994, Svitkin *et al.* 2009). A striking example is that TARBP2 is a non-specific translational repressor in the RRL, while the gene knockout implicates TARBP2 in a pathway that specifically activates *Prm1* and *Prm2* mRNA translation (Lee *et al.* 1996, Zhong *et al.* 1999). In general, heterologous systems cannot elucidate the magnitude of the regulatory effects of elements and factors on mRNA translation in spermatogenic cells.

Numerous elements in flies, worms, *Xenopus* oocytes and cultured mammalian cells have been identified with mutations that abrogate translational control *in vivo*. Mutations in elements eliminate uncertainties concerning the identity of the target mRNA and have often led to evidence that a short mRNA sequence is a translational control element (TCE) that binds a specific factor. Mutations have also provided evidence that elements mediate distinctive mechanisms involving the length of the poly(A) tail, recruitment of PABP without a poly(A) tail, a block to initiation of translation at the level of free mRNPs, subunit joining or reinitiation, regulation in a specific cell type or developmental stage, translation of upstream open reading frames (uORFs), or spatial relations between elements (Mueller & Hinnebusch 1986, McGrew *et al.* 1989, Shyu *et al.* 1989, Levy *et al.* 1991, Ostareck *et al.* 2001, Collier *et al.* 2005, Lyabin *et al.* 2011). As mutations in elements that inactivate individual regulatory pathways can usually be created, they can provide decisive evidence whether a specific factor has a large or small effect on the translational activity of a single mRNA and about the relative importance of multiple pathways in regulating a single mRNA and the importance of the position of elements (Shyu *et al.* 1989, Levy *et al.* 1991, Lyabin *et al.* 2011).

Studies of mutations in elements can also be problematic. They are less likely than knockouts to produce phenotypes. Deletions can produce misleading phenotypes by altering the positions of elements

(see comments regarding the position of the *Prm1* Y-box recognition sequence (YRS) below) or by abrogating the binding of multiple factors. The huge number of factors involved in post-transcriptional regulation makes it likely that replacing a UTR with a heterologous UTR will introduce extraneous elements with unpredictable regulatory effects. In addition, elements may be present in multiple copies in individual UTRs, some or all of which must be mutated to understand their function, and the effects of mutating one element may only be realized in the presence of other elements in the same or a different UTR (Mehta *et al.* 2006, Bartel 2009). In general, mutations that produce strong phenotypes are easy to analyze experimentally, and mutations in small numbers of bases minimize the chance of abrogating the binding of multiple factors.

It is also desirable to identify mutations in elements that result in the total loss of negative or positive translational control of an individual mRNA at a specific stage of spermatogenesis. Mutations of positive control elements are yet to be studied.

Notably, mutations in negative control elements do not result in the total loss of translational repression in round spermatids. In somatic mammalian cells, >85% of fully active mRNAs sediment with polysomes in sucrose gradients (Schmidt *et al.* 1999, Mathews *et al.* 2007). In contrast, mutation or deletion of negative TCEs in the 3' UTRs of two mRNAs that are strongly repressed in round spermatids, *Prm1* and *Smcp*, results in relatively low levels of polysomal mRNA, 35–55% (Schmidt *et al.* 1999, Zhong *et al.* 2001, Hawthorne *et al.* 2006, Bagarova *et al.* 2010). In addition, each of the more than 60 active mRNAs in pachytene spermatocytes and spermatids that have been analyzed with sucrose gradients exhibit 25–55% polysomal mRNA in total testis (Kleene 2003, Paronetto *et al.* 2009, Kleene *et al.* 2010, Chi *et al.* 2011). Many of these partially repressed mRNAs, exemplified by the *Ldhc* mRNA, exhibit a similar polysomal loading in pachytene spermatocytes and round spermatids (Kleene 2003). Thus, the strong initial repression of developmentally regulated mRNAs appears to be superimposed on a ubiquitous background of partial global repression. The factors that are potentially involved in the partial global repression include YBX2/MSY2, PABPC1, EIF4E, EIF4BP1, and ribosomal protein SP6 (Schmidt *et al.* 1999, Messina *et al.* 2010, Yanagiya *et al.* 2010). At present, the contribution of each of these factors to the partial repression of all mRNAs and the strong initial repression of developmentally regulated mRNAs is unknown.

Unfortunately, few mutations in regulatory elements in spermatogenic cells have been analyzed, because cell culture and cell-free translation systems that reproduce the patterns of translational control in spermatogenic cells have not been authenticated. Mutations in transgenic mice have rarely been studied, probably because the approach is expensive, laborious, and prone

Table 2 Translational control *cis*-elements in spermatogenic cells.

RBP: element ^a	Sequence ^b	Approach ^c	Reference ^d
CPEB1: <i>Sycp3</i> 3' UTR	UUUUAAU-24 nt-AAUAAA	u.v. cross-link	Tay & Richter (2001)
DAZL: <i>Mvh</i> 3' UTR	UUUUUUGUUUUUU	Filter binding, RNA-pull down	Reynolds <i>et al.</i> (2005)
DAZL: <i>Sycp3</i> 3' UTR	UUUUUUUUGUUUUUUUUUU	Filter binding	Reynolds <i>et al.</i> (2005, 2007)
DAZL: <i>Mvh</i> and <i>Sycp3</i> 3' UTRs	GUU	Quantification of RNA binding and point mutations	Jenkins <i>et al.</i> (2011)
Unknown: <i>Prm1</i> 3' UTR TCE	GAACAAUGCCACCUGUC	Mutation in transgenic mice	Zhong <i>et al.</i> (2001)
YBX2/MSY2, YBX3/MSY4: <i>Prm1</i> YRS	UCCAUCA	Mutation in transgenic mice	Giorgini <i>et al.</i> (2001)
YBX2/MSY2, YBX3/MSY4: <i>Prm1</i> YRS	[AUC][AC]CA[UC]A[ACU]	RNA-EMSA with point mutations	Giorgini <i>et al.</i> (2001)
YBX2/MSY2, YBX3/MSY4: <i>Prm1</i> YRS	N[AC]CA[UC]AN	Yeast three hybrid with point mutations	Giorgini <i>et al.</i> (2001)
YBX2/MSY2, YBX3/MSY4: <i>Smcp</i> and <i>Tnp1</i> 3' UTR YRS	AACACUU	u.v. cross-link with point mutations	Chowdhury & Kleene (2012)
YBX2/MSY2: all	Non-specific	RNA-EMSA and filter binding	Kwon <i>et al.</i> (1993)
TSN-TSNAX: <i>Prm2</i> 3' UTR Y-element	CUGAGCCCUGAGCU	RNA-EMSA and deletion	Kwon & Hecht (1991)
TSN-TSNAX: <i>Prm2</i> 3' UTR H-element	GCCACGAGAUCCUGAGUAC	RNA-EMSA and deletion	Kwon & Hecht (1991)
TSN-TSNAX: mutated <i>Prm2</i> Y-H	UUUGUGUGUUUUUUUUUUUUUUUUUUUU	EMSA with point mutations	Li & Baraban (2004)
TSN-TSNAX: synthetic homopolymer	AGAGAGAGAGAGAGAGAGAGAGAGA	Quantification of RNA binding with point mutations	Lluis <i>et al.</i> (2010)

^aThe name of the element according to the 3' UTR in which it is found and the factor that it binds if known. ^bThe sequence of the element. ^cThe experimental approach used to determine the sequence of the element. Binding affinities were determined with RNA-EMSA, point or deletion mutations, and quantification of the binding of RNA oligonucleotides to recombinant proteins. ^dThe reference of the article in which the data have been reported.

to negative results. Potential solutions to these problems are described below.

Tables 1 and 2 list factors and elements that have been implicated in translational regulation in mammalian spermatogenesis by studies of gene knockouts, heterologous systems, and transgenic mice. Significantly, most of these factors have not been connected with targets. The following sections scrutinize the small number of systems in which elements have been implicated in translational control in meiotic and haploid spermatogenic cells.

Elements in the *Prm1* 3' UTR: TCE and YRS

The *Prm1* mRNA is first detected in step 7 spermatids and is repressed in free mRNPs until the PRM1 protein is first detected in step 10 spermatids, see Fig. 1 (Kleene 1989, Mali *et al.* 1989, Meistrich *et al.* 2003). In classic studies using the *Prm1* promoter and 5' UTR and human GH (hGH) reporter, Braun *et al.* (1989) demonstrated that the timing of *Prm1* mRNA translation is regulated by the 3' UTR in transgenic mice. Subsequent studies of deletion and point mutations identified two elements in the *Prm1* 3' UTR that repress translation in early spermatids.

The first is two imperfect copies of a TCE, GAAAAAUGCCACCUGUC and GAACAAUGCCACCUGUC, 41 and 0 nts upstream of the AAUAAA poly(A) signal (Zhong *et al.* 2001). Mutation of both the TCEs in the full-length *Prm1* 3' UTR results in a precocious detection of hGH in step 7 round spermatids in transgenic mice. Although it has been suggested that the TCEs are sufficient for the initial repression of *Prm1* mRNA translation, this conclusion requires validation by quantifying the levels of polysomal transgenic mRNAs

containing the wild-type *Prm1* 3' UTR, the TCE mutation, and the *hGH* 3' UTR in step 7–9 spermatids. The factor that binds the TCE has not been identified.

The second is a YRS, UCCAUCA, that binds Y-box proteins YBX2/MSY2 and YBX3/MSY4. In this review, the Y-box protein known as MSY4, CSDA and MSY3L (Davies *et al.* 2000) is referred to as YBX3, the name approved by the Mouse Genome Informatics committee. However, it is important to note that both *Ybx3* and *Msy4* fail to acknowledge that the *Ybx3* gene is expressed as two alternatively spliced mRNAs (Mastrangelo & Kleene 2000) encoding isoforms of different sizes and similar abundance, detected by western blots of testicular proteins (Davies *et al.* 2000, Giorgini *et al.* 2002), which are referred to as YBX3L (long) and YBX3S (short). Both isoforms are conserved in multiple mammalian species, implying that each isoform has at least partially distinct functions.

Fajardo *et al.* (1997) studied an *hGH-Prm1* mRNA in which 95 nts were deleted from the middle of the *Prm1* 3' UTR, thereby moving the YRS from its natural position, 109 nts upstream of the poly(A) signal, to an abnormal position, 15 nts upstream of the poly(A) signal. This 3' UTR, with the YRS in an abnormal position, delays hGH expression in transgenic mice, but the initial repression is incomplete because low levels of hGH are detected in round spermatids (Fajardo *et al.* 1997). Giorgini *et al.* (2001) subsequently showed that a mutation in this YRS, which abrogates the binding of Y-box proteins YBX3L/MSY4 and YBX2/MSY2 *in vitro*, results in translational repression in early spermatids. However, a transgene in which the TCEs are mutated in the full-length *Prm1* 3' UTR abrogates translational repression in early

spermatids, suggesting that the YRS in its natural position is insufficient to repress translation (Zhong *et al.* 2001).

The literature to date has revealed that *Prm1* mRNA translation in round spermatids is repressed by an unidentified factor that binds the TCE and that YBX2/MSY2 and YBX3/MSY4 do not repress translation by binding the YRS in its natural position (Zhong *et al.* 2001). The *Prm1* mRNA is also activated by pathways involving the DEAD-box RNA helicase, DDX25/GRTH, and the protein kinase/double-stranded RBP, TARBP2 (Zhong *et al.* 1999, Tsai-Morris *et al.* 2012). However, none of the pathways, direct or indirect, that repress and activate *Prm1* mRNA translation are known.

uORFs in the *Smcp* and *c-Mos* 5' UTRs

The 5' UTRs of 30–50% of human mRNAs contain uORFs, characterized as reading frames that begin with an AUG codon in the 5' UTRs and are discontinuous with the authentic coding sequence (Hinnebusch 2005). The translation of uORFs promotes the release of the ternary complex (eIF2–GTP–tRNA^{iMet}) from the 40S ribosomal subunit, which prevents ribosomes from re-initiating translation at the authentic ORF downstream. Since uORFs are translated, repression by uORFs produces small polysomes instead of free mRNPs.

The repression of mRNA translation by uORFs in various human mRNAs varies from ~30% (fine-tuning) to ~100% (total repression) and is often accompanied by small decreases in mRNA levels (Calvo *et al.* 2009). The strength of repression is determined by many parameters such as secondary structure, rare codons, context of the uORF initiation codon, sequence of the encoded peptide, length and position of the uORF, and *trans*-acting factors (Sachs & Geballe 2006, Medenbach *et al.* 2011). In general, parameters that prolong the translation of uORFs promote the release of the ternary complex and decrease the translation of the authentic ORF. Conversely, parameters that prolong the period between termination of the uORF and initiation at the authentic initiation codon promote rebinding of the ternary complex to the scanning 40S subunit and increase the translation of the authentic ORF. Although the parameters that determine the strength of repression by uORFs are too complex to predict by examining the primary sequence of 5' UTRs, the functions of uORFs in translational repression can easily be demonstrated by mutating the uORF AUG initiation codon to a non-start codon. The uORFs of two mRNAs that are expressed in spermatids have been demonstrated to developmentally regulate gene expression in transgenic mice.

The SMCP protein is first detected in step 11 spermatids 6 days after the mRNA is detected in step 3 spermatids, see Fig. 1 (Shih & Kleene 1992, Cataldo *et al.* 1996). Translational control of the *Smcp* mRNA has been studied in transgenic mice using the *Smcp* promoter, the *Gfp*-coding region, and the permutations of wild-type

and mutant *Smcp* and *Gfp* UTRs. The translational repression of the *Smcp* mRNA in early spermatids is primarily mediated by the 3' UTR, which produces free mRNPs and delays GFP expression by ~4 days. By comparison, the 5' UTR produces a one- to two-day delay and small polysomes (Hawthorne *et al.* 2006, Bagarova *et al.* 2010). The initial repression by the *Smcp* 5' UTR is mediated by two short uORFs, because mutating the uORF AUG codons to AGG eliminates the delay in GFP expression (Bagarova *et al.* 2010). The uORFs are also involved in the developmental control of *Smcp* mRNA translation, because the *Smcp* 3' UTR exerts a positive control that overcomes the small polysomes mediated by the uORFs in late spermatids (Bagarova *et al.* 2010, TA Chordhury and KC Kleene, unpublished observations).

Steel *et al.* (1996) studied the spermatid-specific 5' UTR of the *Mos/c-Mos* mRNA using the β -galactosidase reporter under the control of the constitutive cytomegalovirus IE promoter. A transgene containing the wild-type *Mos* 5' UTR, which contains four short uORFs, restricts β -galactosidase expression to spermatids, whereas a transgene in which the upstream AUG initiation codons are mutated to AAG restricts β -galactosidase expression to the periphery of seminiferous tubules, presumably spermatogonia and spermatocytes. It is unclear whether the striking effect of the *Mos* uORFs on the stage of β -galactosidase expression is mediated by differences in mRNA translation or stability.

Students of translational control in spermatogenic cells have paid little attention to uORFs even though uORFs are a common, versatile, and easily recognized TCE. The *Akap4*, *Gapdhs*, *Spata18*, *Smcp*, *Spata19*, and primate *HILS1* 5' UTRs contain uORFs that are conserved in diverse mammalian species (Chowdhury & Kleene 2012, KC Kleene, unpublished observations). Interestingly, the uORFs of five of these mRNAs have features that correlate with strong translational repression: the termination codons of the *Gapdhs* and *Spata19* uORFs are just upstream or downstream of the authentic AUG codon, the *Akap4* uORFs are especially long, and the *Spata18* and *Gapdhs* uORFs are predicted to bind Y-box proteins and translin respectively (Yang *et al.* 2003, Chowdhury & Kleene 2012). Mutations have not been studied to determine whether the uORFs have large or small effects on the developmental expression of these mRNAs in spermatids.

DAZL, the deleted in azoospermia-like RBP

The deleted in azoospermia, DAZ, family of proteins is named for a cluster of four genes on the Y-chromosome that has been implicated in 10–15% of cases of human male infertility (reviewed in Reynolds & Cooke (2005) and VanGompel & Xu (2011)). DAZ genes are only present in primates, including old-world monkeys and humans, but other vertebrates have two autosomal paralogs, deleted in azoospermia-like (*DAZL*)

and *BOLL/BOULE*, and all invertebrates likely have *BOLL* orthologs.

On a mixed background, the homozygous *Dazl* knockout produces a final block to the development of spermatocytes marked by the failure of chromosome synapsis during the leptotene–zygotene transition (Saunders *et al.* 2003). The mRNAs encoding a major protein in the lateral element of the synaptonemal complex, SYCP3, and the DEAD-box RNA helicase, mouse Vasa homolog (DDX4/MVH), have been identified as targets of DAZL (Venables *et al.* 2001, Reynolds *et al.* 2005). The *Sycp3* and *Ddx4/Mvh* 3' UTRs contain U-rich DAZL-binding elements (discussed below). The *Sycp3* and *Ddx4/Mvh* knockouts also produce blocks at or near the leptotene–zygotene transition respectively similar to the stage of final arrest in *Dazl*-null mice (Tanaka *et al.* 2000, Yuan *et al.* 2000).

DAZL activates *Sycp3* and *Ddx4/Mvh* mRNA translation, because the *Dazl* knockout reduces DDX4/MVH and SYCP3 protein expression in spermatocytes, without altering the levels of either mRNA (Reynolds *et al.* 2005, 2007). In addition, injection of *Xenopus laevis* oocytes with mouse *Dazl* mRNA enhances the translation of luciferase reporter mRNAs containing the *Ddx4/Mvh* and *Sycp3* 3' UTRs, and the stimulation of translation is abrogated by mutation of the DAZL-binding sites (Reynolds *et al.* 2005, 2007).

Collier *et al.* (2005) analyzed the mechanism by which DAZL stimulates mRNA translation with tethering assays in *Xenopus* oocytes. In these tethering assays, synthetic mRNAs encoding fusions of five DAZ family proteins with the bacteriophage MS2 coat protein were injected into *Xenopus* oocytes followed by injections of luciferase reporter mRNA containing 3' UTR MS2 binding sites. Tethering simplifies the assay of translational activity of reporter mRNAs by circumventing the process by which DAZL normally binds its target mRNAs. *Xenopus* DAZL, mouse DAZL, human DAZL, human DAZ, and human Boule stimulate luciferase activity without significant effects on mRNA stability. Collier *et al.* (2005) further demonstrated that all five DAZ family proteins bind the C-terminal domain of *Xenopus* poly(A)-binding protein, PABP (the ortholog of the mammalian cytoplasmic PABP known as PABPC1), and mapped the segment of DAZL that binds the PABP C-terminal domain with deletion mutations using pull-down and yeast two-hybrid assays. The analysis of DAZL deletions in tethering assays demonstrated that binding of DAZL to PABP is necessary for the stimulation of translation. Thus, DAZL activates translation by binding a 3' UTR element and recruiting initiation factor PABP independent of the poly(A) tail.

As summarized in Table 2, DAZL binds a loosely defined set of elements in various mRNAs that are rich in U and contain the trinucleotide GUU. Indeed, the identification of the DAZL-binding elements is the most convincing of those surveyed here because it is based on consistent findings with a variety of assays including

in vivo u.v. cross-linking and mutations that reduce binding of the DAZL protein and stimulation of translation in tethering assays (Venables *et al.* 2001, Reynolds *et al.* 2005). Surprisingly, quantitative binding assays demonstrate that the affinity of DAZL for GUU is independent of the surrounding bases (Jenkins *et al.* 2011). The brevity of this element complicates identifying the targets of DAZL, because GUU appears in many mRNAs, most of which presumably are not strongly upregulated by DAZL. This conundrum might be resolved by evidence that multiple DAZL-binding elements in a single 3' UTR augment translational stimulation (Collier *et al.* 2005). Alternatively, the stimulation by DAZL may be modulated by additional factors.

CPEB1: the cytoplasmic polyadenylation element-binding protein

The cytoplasmic polyadenylation element (CPE), usually U_{4–5}A_{1–2}U, and the CPE-binding protein (CPEB1) form the core of mechanisms that repress and activate mRNA translation in *Xenopus* oocytes (reviewed in Radford *et al.* (2008)). Translationally regulated mRNAs are synthesized in the nucleus with long poly(A) tails. Following export to the cytoplasm, the poly(A) tails are initially shortened to ~20 As, resulting in translational repression, and later lengthened to ~150 As, resulting in translational activation. CPEB1, CPE, and proximity of the CPE and AAUAAA polyadenylation signal are required for both the poly(A) shortening that represses translation by limiting PABP recruitment and the poly(A) lengthening that activates translation by recruiting PABP.

The *Cpeb1* mRNA is expressed in pachytene spermatocytes and round spermatids, but the developmental expression of the protein has not been analyzed (Tay & Richter 2001). The homozygous *Cpeb1* knockout reduces male fertility about 16-fold, accompanied by increased apoptosis and chromatin fragmentation and large decreases in the number of post-meiotic cells and levels of two major synaptonemal complex proteins, SYCP1 and SYCP3. The final stage at which spermatocytes develop normally in *Cpeb1*-null mice has not been determined.

The *Sycp3* mRNA appears to be regulated directly by CPEB1 in wild-type oocytes because its CPE, UUUUAAU, efficiently u.v.-cross-links to *Xenopus* CPEB1, and translational activation is accompanied by poly(A) lengthening. The evidence that CPEB1 activates the *Sycp1* mRNA in oocytes is much weaker because its CPE, UUUUAUUUUU, is a poor match for the CPE consensus sequence and u.v.-cross-links inefficiently to *Xenopus* CPEB1. More importantly, there is little difference in the length of the *Sycp1* poly(A)s in the ovaries of wild-type and *Cpeb1* knockout mice.

Unfortunately, the poly(A) tails on the *Sycp1* and *Sycp3* mRNAs were not measured in *Cpeb1*-null and

wild-type mouse testes. This is an important gap because the factors that regulate meiosis often differ in oocytes and spermatocytes (Cohen *et al.* 2006) and *Sycp3* mRNA translation is also thought to be stimulated by DAZL (Brook *et al.* 2009).

DAZL, CPEB1, and DDX4/MVH probably control the initiation of meiosis by different mechanisms

Gene knockouts reveal that a trio of well-established activators of mRNA translation, CPEB1, DAZL, and DDX4/MVH, are necessary to initiate meiosis in male mice. The *Dazl* knockout blocks the transition from leptotene to zygotene (Saunders *et al.* 2003). The *Ddx4* and *Cpeb1* knockouts also block early meiosis, but the morphology of the chromosomes that distinguish the normal and abnormal stages has not been defined (Tanaka *et al.* 2000, Tay & Richter 2001).

DAZL and CPEB1 are thought to stimulate translation by different mechanisms. DAZL stimulates translation by recruiting PABPC to the body of the 3' UTR, while CPEB1 stimulates translation by poly(A) lengthening recruiting PABPC (Radford *et al.* 2008, Brook *et al.* 2009). The mouse *Sycp3* mRNA contains CPE- and DAZL-RNA-binding elements. Interestingly, the *Ddx4* 3' UTRs in many mammalian species including mouse contain consensus CPEs ~21–24 nt upstream of the poly(A) signal (Kleene 2013 unpublished). Thus, translation of the *Sycp3* and *Ddx4* mRNAs is potentially stimulated by CPEB1 as well as DAZL.

Assigning a mechanism to DDX4 is difficult because DDX4 has multiple functions. DDX4 belongs to the family of DEAD-box RNA helicases that melts secondary structure and often stimulates the translation of large groups of mRNAs (reviewed in Jankowsky (2011) and Parsyan *et al.* (2011)). In addition, DDX4 functions in the assembly of germ-cell RNP granules and blocks the proliferation of retrotransposons, processes with potential indirect effects on post-transcriptional gene expression (Kirino *et al.* 2010, Kuramochi-Miyagawa *et al.* 2010).

Interestingly, the *Drosophila* ortholog of DDX4, Vasa, is an atypical RNA helicase because it stimulates the translation of the *mei-P26* mRNA by binding a U-rich 3' UTR element and recruiting the initiation factor Eif5b (Liu *et al.* 2009, Parsyan *et al.* 2011). The sequence-specific binding of Vasa is thought to be mediated by an RNA-binding motif, RGG, which appears eight times in the ~200-amino acid-long N-terminal segments of Vasa, outside of the RNA helicase domain (Liu *et al.* 2009). Four RGG motifs are present in the N-terminal segment of mouse and human DDX4, and the IUPred algorithm (<http://iupred.enzim.hu>) predicts that the N-terminal segments of Vasa and DDX4 are unstructured (Kleene 2013 unpublished), a feature shared by many RBPs (Castello *et al.* 2012). Thus, DDX4 and Vasa could activate mRNA translation by two mechanisms, by

binding specific sequences in mRNAs and by melting the secondary structure. A related question is whether the two isoforms of DDX4 and Vasa with RGG motifs in different contexts differ functionally.

In adult testis, 858 potential mRNA targets of DDX4 have been identified by immunoprecipitation of native mRNPs with DDX4 antibody and microarray analysis (Nagamori *et al.* 2011), a technique known as RIP-chip. The identification of direct RNA targets of DEAD-box helicases is considered difficult (Jankowsky 2011) and may be complicated by the fact that DDX4 interacts with many RBPs. Most of these putative targets are expressed in pachytene spermatocytes and spermatids. Since the *Ddx4* knockout blocks early meiosis (Tanaka *et al.* 2000), it cannot be used to determine whether DDX4 is necessary for translational regulation in pachytene spermatocytes and spermatids. mRNAs encoding proteins that initiate meiosis have not been identified, probably because they are present at low levels in adult testis. Perhaps, early meiotic targets of DDX4 could be identified in pure populations of early spermatocytes using *in vivo* u.v. crosslinking and immunoprecipitation discussed below, which appears to be able to capture mRNAs that are in direct contact with RNA helicases *in vivo* (Baltz *et al.* 2012, Castello *et al.* 2012).

Defining the roles of DDX4, CPEB1, and DAZL in the expression of the other two factors and meiotic proteins such as SYCP1 and SYCP3 would greatly increase our understanding of the mechanisms that control the initiation of meiosis in spermatocytes. The function of CPEB1 in activating the translation of the *Sycp1*, *Sycp3*, and *Ddx4* mRNAs in early meiosis could easily be determined by measuring poly(A) lengths with ePAT (Janicke *et al.* 2012) and polysomal loading in leptotene and zygotene spermatocytes in wild-type and *Cpeb1*-null mice. However, determining the magnitude of stimulation by each factor will probably require *in vivo* genetic studies of mutations in elements specific for DDX4, CPEB1, and DAZL.

The testis–brain RBP, translin, and TSNAX/TRAX

The testis–brain RBP (TB-RBP) was initially proposed to repress *Prm2* mRNA translation in round spermatids by specifically binding two elements named Y and H in a 40-nt segment of the *Prm2* 3' UTR referred to as transcript c: 5' CUGAGCCCUGAGCUGCCAA-GGAGCCACGAGAUUCUGAGUAC (Kwon & Hecht 1991, 1993). The Y- and H-elements are single and double underlined respectively. Subsequent work has demonstrated that TB-RBP is the DNA- and RBP translin, TSN (Wu *et al.* 1997), and that TSN is usually associated with a partner, translin-associated factor X (TSNAX/TRAX), which lacks nucleic acid-binding activity but possesses RNase activity (reviewed in Jaendling & McFarlane (2010)). TSN has been implicated in several forms of post-transcriptional gene expression of many mRNAs in

spermatogenic cells: translational repression in round spermatids, export of mRNA to the cytoplasm in spermatocytes, and sharing of post-meiotic transcripts by transport of mRNA through cytoplasmic bridges. The functions of TSN, TSNAX, and the Y- and H-elements are confused by controversy and the absence of *in vivo* genetic evidence.

One controversy concerns the proportions of TSN homomer and TSN·TSNAX heteromer in spermatogenic cells. Finkenshtadt *et al.* (2002) reported that virtually all TSN is associated with TSNAX in testis and cerebellum, while Cho *et al.* (2004) reported that the proportion of TSN homomer progressively increases fourfold to fivefold in pachytene spermatocytes, round spermatids, and elongated spermatids. This disparity is important because the affinity of the TSN homomer for transcript c is much greater and exhibits greater sequence specificity than those of the TSN·TSNAX heteromer (Chennathukuzhi *et al.* 2001, Lluís *et al.* 2010).

Another controversy concerns the sequence of bases to which TSN binds. Li & Baraban (2004) demonstrated that the recombinant and natural TSN·TSNAX heteromer in cerebellum extracts binds interspersed G residues instead of the specific bases of the Y- and H-elements. At present, none of the 68 Y- and H- elements in 42 mRNAs predicted computationally by similarity to the *Prm2* Y- and H-elements are confirmed by RNA binding studies of natural and mutated elements (Han *et al.* 1995).

The critical question here is whether TSN represses mRNA translation in round spermatids. Evidence that partially purified TSN from testis extracts and recombinant TSN homomer repress translation in the RRL (Kwon & Hecht 1993, Yang *et al.* 2003) is subject to artifacts of heterologous systems and disagreement about the proportions of TSN homomer and TSN·TSNAX heteromer *in vivo* discussed above.

A gene trap mutation that abolishes the binding of TSN to mRNA has been created (Chennathukuzhi *et al.* 2003) and could be used to analyze the functions of TSN. However, the phenotype of the *Tsn* knockout is in disagreement with the expected consequences of depleting the putative repressor of many mRNAs in round spermatids. Premature translation of the *Prm1* and *Tnp2* mRNAs in transgenic mice produces deformed sperm and reduced male fertility (Lee *et al.* 1995, Tseden *et al.* 2007). The *Tsn* knockout might, therefore, be expected to have drastic effects on spermatid differentiation, because TSN is postulated to repress the translation of the *Gapdhs*, *Prm1*, *Prm2*, *Tnp1*, *Tnp2*, and many other mRNAs. In contrast, the *Tsn* knockout reduces sperm number and litter size by approximately twofold, increases apoptosis in spermatocytes and results in the degradation of TSNAX, but does not increase the frequency of apoptotic round spermatids and abnormal sperm (Chennathukuzhi *et al.* 2003). The phenotype of the *Celf3/Brunol1* knockout is very similar to that of the *Tsn* knockout, and it was interpreted

as evidence that CELF3/BRUNOL1 'might be non-essential for spermatogenesis' (Dev *et al.* 2007). The possibility that TSN does not repress translation *in vivo* cannot be dismissed because there are no reports of determinations whether the *Tsn* knockout results in a precocious activation of repressed mRNAs by the analysis of polysomal loading with sucrose gradients or protein expression in the testes of prepubertal mice.

Recent work has shown that the TSN·TSNAX heteromer, sometimes referred to as C3PO, functions in the RNA interference pathway by increasing the rate of degradation of the passenger RNA by the RNA interference silencing complex (Tian *et al.* 2011, Ye *et al.* 2011). This is probably a minor or redundant function because knockouts of the *Tsn* and/or *Tsnax* genes in fission yeast, flies, and mammals are viable and fertile (Jaendling & McFarlane 2010).

Approaches for connecting elements and factors

The challenges in connecting mRNA targets with regulatory factors originate in the number of different factors and the variety of parameters that influence the binding of RBPs to RNA. RBPs in eukaryotic cells are classified by RBDs, which exhibit different modes of interaction with RNA and functions (reviewed in Kishore *et al.* (2010), Morris *et al.* (2010) and Elliott & Ladomery (2011)). Table 1 lists a diverse assortment of RBPs that have been implicated in translational regulation in spermatogenic cells. These include RBDs that bind specific sequences of bases in a single-stranded RNA (ssRNA), DAZL and CPEB1, or a specific sequence of bases in all mRNAs, PABPC1. Other RBPs, such as TARBP2 and ADAR1, contain the dsRBD that binds the ribose sugars and phosphates of the backbones of double-stranded RNA segments instead of the bases. Some dsRBPs bind dsRNA segments in mRNAs non-specifically, while others bind mRNA-specific configurations of multiple dsRNA segments. Argonaute proteins form complexes with miRNAs that are guided to specific mRNAs by complementary base-pairing (Bartel 2009, Fabian & Sonenberg 2012). Two factors in Table 1, EIF4G3 and PAIP2a, are not known to bind mRNA directly, but instead interact with initiation factors that bind mRNA. The versatility of interactions of RBPs with mRNA is documented in the general reviews cited above as well as specialized reviews of RBP families (Skabkin *et al.* 2006, Brook *et al.* 2009, Jankowsky 2011).

The binding of sncRNAs and RBPs to mRNAs is also influenced by the accessibility to mRNAs (Kishore *et al.* 2010, Morris *et al.* 2010). Access is determined by mRNA secondary structure, RBPs, and sncRNAs, which promote or block binding to specific elements, and RNP remodeling proteins such as Y-box proteins and DEAD-box RNA helicases, which dissociate and rearrange dsRNA segments and RBPs (Skabkin *et al.* 2006, Jankowsky 2011).

The specificity of interactions of RBPs with RNAs can also be defined by intracellular localization. Certain DEAD-box RNA helicases interact non-specifically with RNA *in vitro*, but remodel specific RNAs *in vivo* because the helicases function in a multiprotein complex containing hundreds of proteins and sncRNAs that regulates specific classes of RNAs and RNPs (Jankowsky 2011). This may be a property of the two DEAD-box helicases, DDX4/MVH and DDX25/GRTH, which are major constituents of the cytoplasmic germ-cell RNP granules known as the intermitochondrial cement in spermatocytes and the chromatoid body in spermatids (Meikar *et al.* 2011).

The elements and mRNAs to which factors bind can be identified by biochemical studies of molecular interactions, computational searches for elements shared by co-regulated RNAs, or a combination of both, e.g. co-immunoprecipitation and microarray analysis or deep sequencing (reviewed in Kishore *et al.* (2010), Morris *et al.* (2010) and Ascano *et al.* (2012)). Each approach is distinguished by tradeoffs in simplicity, cost, and reliability in distinguishing *bona fide* targets and false positives, and every approach is capable of producing misleading information. Not surprisingly, the sequences of elements obtained using different techniques differ markedly (Table 2). Some of these differences arise because techniques that define elements precisely reveal that RBPs bind sequences that are often shorter, 3–6 nts, and more degenerate than were thought previously (Ray *et al.* 2009, Jenkins *et al.* 2011, Ascano *et al.* 2012).

The complexity of the parameters that determine the interactions of RBPs with RNA provides a compelling argument for identifying RBP–mRNA interactions in the natural environment of living cells. This has been achieved by variations of a procedure known as CLIP, based on the principle that u.v. irradiation at 254 nm cross-links bases and amino acids that are in direct contact in living cells and does not cross-link proteins (Ule *et al.* 2005). After cross-linking, mRNP fragments containing short segments of mRNA surrounding an element are purified with limited RNase digestion, immunoprecipitation to selectively concentrate the RNA–protein complexes, and SDS–PAGE to separate complexes from non-cross-linked RNAs and proteins.

A recent modification of CLIP, photoactivatable ribonucleoside-enhanced crosslinking and immunoprecipitation (PAR-CLIP), utilizes the incorporation of photoactivatable nucleosides and u.v. cross-linking at 365 nm *in vivo* (Hafner *et al.* 2010, Ascano *et al.* 2012). PAR ribonucleosides, such as 4-thiouridine and 6-thioguanosine, are incorporated into RNA by cells and cross-linked to protein at a wavelength, 365 nm, that minimizes damage to the RNA. Opinions differ whether cross-linking of PAR bases at 365 nm is 1.3- or >100-fold more efficient than that of natural bases at 254 nm (Kishore *et al.* 2011, Ascano *et al.* 2012). Importantly, cross-linking of amino acids to 4-thiouracil and

6-thioguanine in mRNA produces T to C and G to A transitions in cDNA copies respectively, providing criteria to identify the precise binding site and distinguish cross-linked targets from non-cross-linked contaminants. High-throughput sequencing of CLIP cDNAs, HITS-CLIP, has also been shown to detect single-nucleotide deletions at bases that are cross-linked to amino acids (Zhang & Darnell 2011). Thus, deep sequencing of mRNA fragments purified with PAR-CLIP and CLIP is capable of defining elements by the affinity of the RBP for all permutations of bases within and surrounding the element and the context of the element in relation to the local secondary structure and binding of other RBPs and sncRNAs.

The invention of PAR-CLIP has stimulated transcriptome-wide analyses of RBP–mRNA interactions. PAR-CLIP and CLIP, in combination with oligo-dT purification of mRNPs and quantitative proteomics, identify ~800–860 proteins that bind polyadenylated mRNA in cultured mammalian cells (Baltz *et al.* 2012, Castello *et al.* 2012). Using deep sequencing, Baltz *et al.* (2012) also created a transcriptome-wide inventory of nucleotide sites that are bound by RBPs. PAR-CLIP identifies ~5000–7500 mRNAs bound by ELAVL1/HuR using 4-thiouridine, a judicious PAR base because this RBP binds U-rich elements (Kishore *et al.* 2011, Lebedeva *et al.* 2011, Mukherjee *et al.* 2011).

These studies have also shed light on the advantages and disadvantages of various methods for identifying RBP–mRNA interactions that are important in post-transcriptional regulation. The identification of RBP–RNA interactions is influenced by many technical factors including the quality of the antibody, the specificity and levels of the RNase used to purify short RNA–RBP complexes, and the depth of sequencing (Morris *et al.* 2010, Kishore *et al.* 2011, Lebedeva *et al.* 2011, Mukherjee *et al.* 2011).

It is clear that a single procedure does not identify all RBP–mRNA interactions. Irradiation at 254 nm preferentially cross-links pyrimidines with cysteine, lysine, phenylalanine, tyrosine, and tryptophan (Ule *et al.* 2005), and the potential for bias with PAR-CLIP is greater because cross-linking is restricted to elements containing the PAR base. Nevertheless, most RBPs are cross-linked to polyadenylated mRNA with both CLIP and PAR-CLIP, although 10–20% of the RBPs are identified with either procedure, but not both (Castello *et al.* 2012).

PAR-CLIP identifies about fourfold more ELAV1/HuR targets than does immunoprecipitation of native mRNPs without cross-linking, RIP-chip (Mukherjee *et al.* 2011). The greater number of mRNAs identified with PAR-CLIP was explained by the idea that PAR-CLIP detects both stable and transient mRNA–RBP interactions, while RIP-chip only detects stable interactions. This hypothesis is consistent with the observation that mRNA targets identified with RIP-chip exhibit slightly greater decreases

in mRNA levels in the *ELAV1/HUR* siRNA knockdown than those identified with PAR-CLIP. Many targets identified with RIP-chip were not confirmed with PAR-CLIP (Mukherjee *et al.* 2011). The cause of this difference is unclear, but a potential explanation is that *ELAV1/HUR* binds some mRNAs indirectly through complexes with other RBPs.

The identification of mRNA targets with CLIP, PAR-CLIP, or RIP-chip does not indicate that an interaction has major regulatory effects. Indeed, siRNA knockdown of *ELAV1/HuR* reveals that many putative targets undergo very small decreases in mRNA levels and that the most responsive mRNAs have multiple 3' UTR *ELAV1/HuR*-binding sites or binding sites in both introns and 3' UTRs (Lebedeva *et al.* 2011, Mukherjee *et al.* 2011).

The combination of CLIP, proteomics, and deep sequencing has the potential to enormously increase our understanding of the interactions of mRNAs and RBPs in spermatogenic cells. Fortunately, the costs of deep sequencing are decreasing rapidly, and PAR bases can probably be incorporated easily in seminiferous tubule culture (Kleene 1993). Importantly, the ability of CLIP and PAR-CLIP to detect transient RNA–RBP interactions enables the identification of mRNA targets of RBPs that usually do not recognize specific sequences of bases in mRNA, e.g. argonautes, dsRBPs, and RNA helicases (Kishore *et al.* 2011, Zhang & Darnell 2011, Baltz *et al.* 2012, Castello *et al.* 2012). This will facilitate the identification of the direct targets of miRNAs and RBPs in spermatogenic cells that are presently unknown, such as ADAR, TARBP2, DDX4/MVH, and DDX25/GRTH. Despite the successes of CLIP and PAR-CLIP, Castello *et al.* (2012) caution that individual RBP–mRNA interactions should be validated before inferring mechanisms and consequences of post-transcriptional regulation. In fact, CLIP was one of the several techniques that were used to identify the elements to which DAZL binds, and these elements were later validated by studies of mutations in tethering assays (Reynolds *et al.* 2005, 2007).

miRNAs and endo-siRNAs

The role of sncRNAs in spermatogenesis has been studied with knockouts of the *Dicer1* gene encoding a Class III double-stranded RNA endonuclease that functions in the biosynthesis of two classes of sncRNAs, miRNAs and endogenous small interfering RNAs, endo-siRNAs (Song *et al.* 2011). Mature miRNAs and endo-siRNAs bind argonaute proteins and guide complexes to mRNA targets by base-pairing (Bartel 2009, Fabian & Sonenberg 2012). Endo-siRNAs appear to promote mRNA degradation and regulate transcription (Song *et al.* 2011), while miRNAs usually repress mRNA translation, decrease stability, or both (Bartel 2009, Fabian & Sonenberg 2012). A prolonged controversy

concerning the mechanism by which miRNAs repress translation in somatic cells has been resolved recently with evidence that miRNAs function in a pathway that inhibits binding of the small ribosomal subunit to the 5' cap, usually followed by deadenylation (reviewed in Fabian & Sonenberg (2012) and Izaurralde (2012)).

The function of *DICER1* has been studied using conditional *CRE–Lox* knockouts at specific stages of spermatogenesis, because the germ-line deletion of *Dicer1* results in embryonic lethality (Bernstein *et al.* 2003). *Dicer1* knockouts in pre-natal male germ cells and post-natal spermatogonia produce male infertility accompanied by abnormalities in round and elongated spermatids and severe decreases in sperm number (Korhonen *et al.* 2011, Romero *et al.* 2011). However, the pre-natal germ-cell *Dicer1* knockout produces defects in early spermatocytes that are not observed with the spermatogonia knockout.

The phenotypes of male conditional *Dicer1* knockouts will be particularly difficult to connect with specific mRNAs for reasons that have been cogently discussed by Bartel (2009). One reason is that *DICER1* functions in the biosynthesis of hundreds of miRNAs and endo-siRNAs. The large number of sncRNAs that are depleted by *Dicer1* knockouts greatly magnifies the problems in identifying direct targets: each miRNA interacts with tens to thousands of mRNAs, and some of these mRNAs presumably encode RBPs that regulate the expression of other genes at the transcriptional and post-transcriptional levels (Bartel 2009). Another reason is that knockouts of individual miRNAs often fine-tune the expression of hundreds of mRNAs, which cumulatively produce pronounced phenotypes (Bartel 2009). However, subtle changes in the expression of individual mRNAs are difficult to detect experimentally.

An important challenge of future work will be to determine whether individual miRNAs and endo-siRNAs have large or small effects in regulating the translation and stability of specific mRNAs in spermatocytes and spermatids.

Identification of *cis*-elements and factors that coordinate the translation of mRNAs

Deep sequencing identification of the *cis*-elements bound by RBPs will probably lead to insights into the factors that co-regulate mRNAs with similar patterns of developmental regulation (Kishore *et al.* 2010, Morris *et al.* 2010). However, current knowledge of the developmental changes in mRNA translation in meiotic and haploid spermatogenic cells is arguably too imprecise to identify co-regulated mRNAs. The intricate differentiation of spermatids involves chromatin remodeling and the ordered construction of the acrosome, outer dense fibers, fibrous sheath, mitochondrial sheath, and other subcellular structures unique to the spermatozoon

(Russell *et al.* 1990, Meistrich *et al.* 2003). Clustering co-regulated mRNAs according to the stages of translational repression and activation and association with specific organelles is likely to be more informative than the usual Ingenuity-based classifications based on functions in somatic cells. The temporal translational activity of relatively few mRNAs has been precisely described and is often confused by disagreements in reports from different laboratories and multiple mRNA variants encoded by a single gene. For example, the *Odf2* mRNA, which is studied as a typical mRNA that is transcribed in round spermatids and translated in elongated spermatids, is expressed as two variants with different 3' UTRs, 103- and 1085-nt long. The effect of the alternative *Odf2* 3' UTRs on post-transcriptional regulation is unknown.

Iguchi *et al.* (2006) used microarrays and sucrose gradients in a transcriptome-wide analysis of polysomal loading of prepubertal testes in which the most advanced cells were pachytene spermatocytes (17 days), round spermatids (22 days), and elongated spermatids (adult). Unfortunately, the inferences of developmental changes in polysomal loading are confused by high levels of polysomal mRNA at stages at which the corresponding protein cannot be detected and presentation of the findings in a form (absent or present) that does not indicate the proportions of mRNA in the free mRNP and polysomal fractions (discussed in Kleene *et al.* (2010)). Arava *et al.* (2003) provided a state-of-the-art prototype for this kind of study.

Iguchi *et al.* (2006) also proposed that the *Pgk2* and *Dbil5* mRNAs exemplify a novel mode of developmental translational control in which mRNAs are stored in free mRNPs in pachytene spermatocytes and translationally activated in round spermatids. Unfortunately, this idea is not supported by either their data or the literature. RT-PCR and northern blots concur that the *Pgk2* mRNA is expressed from preleptotene spermatocytes to elongated spermatids (McCarrey *et al.* 1992, Yoshioka *et al.* 2007), while immunocytochemistry in four laboratories reveals that the PGK2 protein is first detected in step 9 (Vandeberg *et al.* 1981) or step 12 spermatids (Kramer 1981, Bluthmann *et al.* 1982, Danshina *et al.* 2010). Similarly, *in situ* hybridization and immunohistochemistry demonstrate that the *Dbil5* mRNA and protein are respectively first detected in step 6–8 and step 16 spermatids in rats (Kolmer *et al.* 1997, Pusch *et al.* 2000). The developmental expression of the *Dbil5* and *Pgk2* mRNAs and proteins is diagrammed in Fig. 1. The microarray and RT-qPCR data of Iguchi *et al.* (2006) also show that the polysomal loading of the *Pgk2* and *Dbil5* mRNAs increases sharply between round spermatids and elongated spermatids. Thus, all of these findings concur with the generally accepted view that the developmental regulation of *Pgk2* and *Dbil5* mRNA translation in spermatids circumvents the transcriptional inactivity

imposed by chromatin remodeling in elongating and elongated spermatids.

I know of no mRNAs that have been demonstrated by *in situ* hybridization, immunohistochemistry, and sucrose gradients to undergo significant increases in translational activity in round spermatids. Further work will be required to determine whether RBPs that have been implicated in the development of round spermatids are required to activate the translation of individual mRNAs at specific stages or to sustain a constant rate of mRNA translation (Paronetto *et al.* 2009, VanGompel & Xu 2010).

The task of deducing the functions of factors and elements identified in high-throughput studies of mRNA translation would be facilitated by a carefully annotated compilation of the developmental expression of mRNA variants and proteins in spermatogenic cells. At present, this list does not exist and would require a painstaking analysis of the literature to assess the quality of the evidence and resolve disagreements between reports.

Validating *cis*-elements with studies of mutations *in vivo*

Studies of gene expression in spermatogenic cells at every level, from transcription to translation and stability, have been thwarted by the absence of a suitable technique for validating *cis*-elements with mutations. *In vivo* genetic analyses of mutations in elements have been limited to transgenic mice, which are considered too costly and risky by many. However, it is debatable whether the inconvenience of transgenic mice is worse than the question of whether information based on studies of mutations in heterologous systems might be invalid. Such studies would also benefit from quantification to determine whether a specific element has a major or minor effect on translational activity (Kleene *et al.* 2010). This impasse can potentially be solved by combining several of the strategies described below.

The risk of squandering resources on mutations in *cis*-elements that produce no phenotypes can be reduced by using comparative genomics to identify conserved sequences or RNA binding assays to identify elements. Both approaches have been used successfully by molecular biologists, with the caveat that elements often have negligible regulatory effects. Predicting binding sites of RBPs in spermatogenic cells that are likely to produce strong phenotypes will become easier as information becomes available from transcriptome-wide studies in somatic and spermatogenic cells. The studies cited above indicate that the critical parameters vary among RBPs and elements and include binding affinity, number of binding sites, binding sites in both introns and 3' UTRs, positions within 3' UTRs and 5' UTRs, and distance from the binding sites of miRNAs and other RBPs.

Comparative genomics is appealing because it is fast and cheap. The public databases contain more-or-less

complete sequences of >30 mammalian genomes. In fortuitous situations, such as the *Ddx4/Mvh* CPE noted above, orthologous UTRs in >20 mammals can be identified in an afternoon, leading to insights that force the revision of working hypotheses (Chowdhury & Kleene 2012).

The importance of developing a culture system for analyzing large numbers of mutations in spermatogenic cells is widely recognized. Many types of culture systems have been explored to identify genes and endocrine and paracrine factors that support spermatogenesis and to generate spermatozoa to produce genetically modified animals and remedy male infertility (reviewed in Lo & Domes (2011) and Hunter *et al.* (2012)). It should be noted that culture systems that support abnormal development or yield small numbers of spermatozoa are adequate for many purposes. However, a culture system for studies of the effects of mutations on gene expression must meet particular standards: it must be possible to transfect the cells with DNA, the development of the spermatogenic cells must be sufficiently normal and reproducible to detect subtle differences in gene expression, the system should produce enough cells for biochemical studies, and development should be rapid to enable studies of many mutations. No system fulfills all of these requirements, and the advantages and disadvantages of two options are described below.

Single-cell populations of round spermatids can be maintained for 72 h in 2D culture and develop to step 8, a stage characterized by flagella and displacement of the nucleus and cytoplasm to opposite sides of the cell (Dehugara *et al.* 2012). Similarly, populations of single-cell pachytene spermatocytes can be cultured in 2D culture for 24–36 h and induced with okadaic acid to progress to an abnormal meiotic metaphase I (La Salle *et al.* 2009, Paronetto *et al.* 2009). These culture systems might be useful in studies of the factors and elements that promote the completion of meiosis and round spermatid stage. The shortcomings are that both culture systems require laborious cell separations to prepare enriched cells from large numbers of mice, and it has not been demonstrated that either cell population can be transfected with DNA, although the round spermatids could be transfected with protein.

Another approach is based on recent studies of spermatogonial stem cells (reviewed in Kanatsu-Shinohara & Shinohara (2010) and Ivics *et al.* (2011)). Spermatogonial stem cells proliferate in culture for more than 2 years and transfect readily with DNA. The transfected cells can be purified by cloning in selective media, preserved cryogenically, and used to generate transgenic and knockout mice and rats by implantation into the testes of infertile recipients. The stem cells repopulate the seminiferous tubules and differentiate into spermatozoa, half of which contain the transgene. This is the theoretical maximum because spermatogonia are diploid. Since spermatozoa, derived from genetically

modified spermatogonial stem cells, are able to fertilize eggs through copulation or microinjection of sperm and produce normal offspring, the cultured stem cells appear to be free of epigenetic and genetic mutations (Kanatsu-Shinohara *et al.* 2005). Creating transgenic and knockout mice by transfection of spermatogonial stem cells and repopulation of the testes of sterile recipients is technically easier and more efficient than by conventional approaches using pronuclear injection and blastocyst chimeras. Thus, the use of spermatogonial stem cells has been advanced as a cost-effective method for creating knockout and transgenic mice, and the approach may become widely available in core facilities. Again there is a downside: the repopulation of seminiferous tubules is a prolonged process requiring 2–3 months for the injected stem cells to differentiate into spermatozoa. It is unclear whether this disadvantage offsets the advantages in decreased screening and time to generate transgenic lines.

The analysis of mutations in transgenic mice can also be expedited by studying the GFP reporter in seminiferous tubule squashes in transgenic founders. Each founder can be analyzed in several hours, eliminating delays and costs in establishing lines, and the levels of GFP fluorescence can be quantified with programs such as ImageJ (Ventelä *et al.* 2000a, Bagarova *et al.* 2010). One testis from an adult male can be analyzed biochemically and the other with seminiferous tubule squashes. Phase-contrast images of tubule squashes display the morphological details that distinguish every stage of meiotic and haploid spermatogenic cells more clearly than do photographs of paraffin sections (Kotaja *et al.* 2004). A reporter in which GFP is fused to acrosin localizes GFP in the acrosome, the critical organelle in staging round spermatids, and may prevent the release of soluble GFP from lysed cells (Ventelä *et al.* 2000b). Importantly, the stage at which proteins encoded by the *Smcp-Gfp* and *Prm1-hGH* transgenes are first detected is independent of variations in mRNA levels originating in differences in the number and position of transgene insertions (Braun *et al.* 1989, Fajardo *et al.* 1997, Giorgini *et al.* 2001, Bagarova *et al.* 2010).

A study analyzing approximately ten mutations by whatever means would initiate a new era in studies of gene regulation in spermatogenic cells.

Conclusions and important questions to be answered

Table 1 lists factors that have been implicated in translational regulation in spermatogenic cells. The knockouts of most of these factors produce male infertility or subfertility and abnormal sperm, demonstrating the importance of post-transcriptional regulation in normal development of spermatozoa. However, the knockouts of the *Tsn*, *Celf3*/Brunol1, and *Ybx3* genes do not produce abnormal sperm and therefore appear to have minor functions. Unfortunately, none of

the *cis*-elements mentioned in this review has been demonstrated to interact with a specific factor and regulate translation in its natural position in spermatogenic cells. The failure to connect factors and target mRNAs with translational control limits our understanding of the mechanisms, which in turn hinders formulating hypotheses that are likely to lead to important insights into the mechanisms by which abnormal post-transcriptional regulation decreases male fertility.

The combination of CLIP, deep sequencing, and transgenic mice provides approaches to rigorously identify the direct targets of specific RBPs and miRNAs and to determine their precise function in the regulation of specific mRNAs. These techniques should make it possible to identify individual miRNAs, endo-siRNAs, and mRNA targets that are major regulators of spermatogenesis.

Mutations in regulatory elements in transgenic mice also provide a direct approach to resolving controversies concerning the factors and elements that regulate the translation of the *Prm2*, *Sycp3*, and *Hspa2* mRNAs. The translation of the *Prm2* mRNA has been proposed to be repressed by the Y- and H-elements, a 3' UTR sequence similar to the *Prm1* TCE and two coding region elements that bind the atypically long 26-nt miRNA₄₆₉ (Kwon & Hecht 1993, Zhong *et al.* 2001, Dai *et al.* 2011), while the *Sycp3* and *Ddx4/Mvh* mRNAs have been proposed to be activated by the CPE- and DAZL-binding sites (Tay & Richter 2001, Brook *et al.* 2009, see above). If the activation of *Hspa2* mRNA translation by EIF4G3 and ELAVL1/HuR can be connected with specific elements, it will be possible to determine which pathway is more important for the exit from meiosis (Sun *et al.* 2010, Chi *et al.* 2011). It should also be possible to identify the unknown factor that binds the *Prm1* TCE with CLIP and immunoprecipitation with the appropriate antibody, regardless of whether the factor is an sncRNA or an RBP. Bridging these gaps will increase our understanding of the mechanisms that regulate three important transitions in spermatogenesis: the initiation of meiosis, the completion of meiosis and the differentiation of late spermatids.

Declaration of interest

The author declares that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

Acknowledgements

I am grateful to the many scientists whose careful work provided insights into the complicated and fascinating domains of eukaryotic post-transcriptional gene regulation and spermatogenesis.

References

- Arava Y, Wang Y, Storey JD, Liu CL, Brown PO & Herschlag D 2003 Genome-wide analysis of mRNA translation profiles in *Saccharomyces cerevisiae*. *PNAS* **100** 3889–3894. (doi:10.1073/pnas.0635171100)
- Ascano M, Hafner M, Cekan P, Gerstberger S & Tuschl T 2012 Identification of RNA–protein interaction networks using PAR-CLIP. *Wiley Interdisciplinary Reviews. RNA* **3** 159–177. (doi:10.1002/wrna.1103)
- Bagarova J, Chowdhury TA, Kimura M & Kleene KC 2010 Identification of elements in the Smcp 5' and 3' UTR that repress translation and promote the formation of heavy inactive mRNPs in spermatids by analysis of mutations in transgenic mice. *Reproduction* **140** 853–864. (doi:10.1530/REP-10-0323)
- Baltz AG, Munschauer M, Schwanhäusser B, Vasilie A, Murakawa Y, Schueler M, Youngs N, Penfold-Brown D, Drew K, Milek M *et al.* 2012 The mRNA-bound proteome and its global occupancy profile on protein-coding transcripts. *Molecular Cell* **46** 674–690. (doi:10.1016/j.molcel.2012.05.021)
- Bartel DP 2009 MicroRNAs: target recognition and regulatory functions. *Cell* **136** 215–233. (doi:10.1016/j.cell.2009.01.002)
- Bernstein E, Kim SY, Carmell MA, Murchison EP, Alcorn H, Li MZ, Mills AA, Elledge SJ, Anderson KV & Hannon GJ 2003 Dicer is essential for mouse development. *Nature Genetics* **35** 215–217. (doi:10.1038/ng1253)
- Bluthmann H, Cicurel L, Kuntz GW, Haedenkamp G & Illmensee K 1982 Immunohistochemical localization of mouse testis-specific phosphoglycerate kinase (PGK-2) by monoclonal antibodies. *EMBO Journal* **1** 479–484.
- Braun RE, Peschon J, Behringer RR, Brinster RL & Palmiter RD 1989 Protamine 3'-untranslated sequences regulate temporal translational control and subcellular localization of growth hormone in spermatids of transgenic mice. *Genes and Development* **3** 793–802. (doi:10.1101/gad.3.6.793)
- Brook M, Smith JW & Gray NK 2009 The DAZL and PABP families: RNA-binding proteins with interrelated roles in translational control in oocytes. *Reproduction* **137** 595–617. (doi:10.1530/REP-08-0524)
- Calvo SE, Pagliarini DJ & Mootha VK 2009 Upstream open reading frames cause widespread reduction of protein expression and are polymorphic among humans. *PNAS* **106** 7507–7512. (doi:10.1073/pnas.0810916106)
- Castello A, Fischer B, Eichelbaum K, Horos R, Beckmann BM, Strein C, Davey NE, Humphreys DT, Preiss T, Steinmetz LM *et al.* 2012 Insights into RNA biology from an atlas of mammalian mRNA-binding proteins. *Cell* **149** 1393–1406. (doi:10.1016/j.cell.2012.04.031)
- Cataldo L, Baig K, Oko R, Mastrangelo MA & Kleene KC 1996 Developmental expression, intracellular localization, and selenium content of the cysteine-rich protein associated with the mitochondrial capsules of mouse sperm. *Molecular Reproduction and Development* **45** 320–331. (doi:10.1002/(SICI)1098-2795(199611)45:3<320::AID-MRD9>3.0.CO;2-U)
- Chennathukuzhi VM, Kurihara Y, Bray JD & Hecht NB 2001 Trax (translin-associated factor X), a primarily cytoplasmic protein, inhibits the binding of TB-RBP (translin) to RNA. *Journal of Biological Chemistry* **276** 13256–13263. (doi:10.1074/jbc.M009707200)
- Chennathukuzhi V, Stein JM, Abel T, Donlon S, Yang S, Miller JP, Allman DM, Simmons RA & Hecht NB 2003 Mice deficient for testis-brain RNA binding protein exhibit coordinate loss of TRAX, reduced fertility, altered gene expression in the brain and behavior changes. *Molecular and Cellular Biology* **23** 6419–6434. (doi:10.1128/MCB.23.18.6419-6434.2003)
- Chi MN, Auriol J, Jégou B, Kontoyiannis DL, Turner JM, de Rooij DG & Morello D 2011 The RNA-binding protein ELAVL1/HuR is essential for mouse spermatogenesis, acting both at meiotic and postmeiotic stages. *Molecular Biology of the Cell* **22** 2875–2885. (doi:10.1091/mbc.E11-03-0212)
- Cho YS, Chennathukuzhi VM, Handel MA, Eppig J & Hecht NB 2004 The relative levels of translin-associated factor X (TRAX) and testis brain RNA-binding protein determine their nucleocytoplasmic distribution in male germ cells. *Journal of Biological Chemistry* **279** 31514–31523. (doi:10.1074/jbc.M401442200)
- Chowdhury TA & Kleene KC 2012 Identification of potential regulatory elements in the 5' and 3' UTRs of 12 translationally regulated mRNAs in mammalian spermatids by comparative genomics. *Journal of Andrology* **33** 244–256. (doi:10.2164/jandrol.110.012492)

- Cohen PE, Pollack SE & Pollard JW 2006 Genetic analysis of chromosome pairing, recombination, and cell cycle control during first meiotic prophase in mammals. *Endocrine Reviews* **27** 398–426. (doi:10.1210/er.2005-0017)
- Collier B, Gorgoni B, Loveridge C, Cooke HJ & Gray NK 2005 The DAZL family proteins are PABP-binding proteins that regulate translation in germ cells. *EMBO Journal* **24** 2656–2666. (doi:10.1038/sj.emboj.7600738)
- Connolly CM, Dearth AT & Braun RE 2005 Disruption of murine Tenr results in teratospermia and male infertility. *Developmental Biology* **278** 13–21. (doi:10.1016/j.ydbio.2004.10.009)
- Dai L, Tsai-Morris CH, Sato H, Villar J, Kang JH, Zhang J & Dufau ML 2011 Testis-specific miRNA-469 up-regulated in gonadotropin-regulated testicular RNA helicase (GRTH/DDX25)-null mice silences transition protein 2 and protamine 2 messages at sites within coding region, implication for its role in germ cell development. *Journal of Biological Chemistry* **286** 44306–44318. (doi:10.1074/jbc.M111.282756)
- Danshina PV, Geyer CB, Dai Q, Goulding EH, Willis WD, Kitto GB, McCarrey JR, Eddy EM & O'Brien DA 2010 Phosphoglycerate kinase 2 (PGK2) is essential for sperm function and male fertility in mice. *Biology of Reproduction* **82** 136–145. (doi:10.1095/biolreprod.109.079699)
- Dass B, Tardif S, Park JY, Tian B, Weitlauf HM, Hess RA, Carnes K, Griswold MD, Small CL & Macdonald CC 2007 Loss of polyadenylation protein tauCstF-64 causes spermatogenic defects and male infertility. *PNAS* **104** 20374–20379. (doi:10.1073/pnas.0707589104)
- Davies HG, Giorgini F, Fajardo MA & Braun RE 2000 A sequence-specific RNA binding complex expressed in murine germ cells contains MSY2 and MSY4. *Developmental Biology* **221** 87–100. (doi:10.1006/dbio.2000.9658)
- Dehnugara T, Dhar S & Rao MR 2012 An *in vitro*, short-term culture method for mammalian haploid round spermatids amenable for molecular manipulation. *Molecular Reproduction and Development* **79** 19–30. (doi:10.1002/mrd.21396)
- Dev A, Nayernia K, Meins M, Adham I, Lacone F & Engel W 2007 Mice deficient for RNA-binding protein bruno1 show reduction of spermatogenesis but are fertile. *Molecular Reproduction and Development* **74** 1456–1464. (doi:10.1002/mrd.20742)
- Elliott D & Ladomery M 2011 *Molecular Biology of RNA*. Oxford: Oxford University Press.
- Fabian MR & Sonenberg N 2012 The mechanics of miRNA-mediated gene silencing: a look under the hood of miRISC. *Nature Structural & Molecular Biology* **19** 586–593. (doi:10.1038/nsmb.2296)
- Fajardo MA, Haugen HS, Clegg CH & Braun RE 1997 Separate elements in the 3' untranslated region of the mouse protamine 1 mRNA regulate translational repression and activation during murine spermatogenesis. *Developmental Biology* **191** 42–52. (doi:10.1006/dbio.1997.8705)
- Finkenstadt PM, Jeon M & Baraban JM 2002 Trax is a component of the Translin-containing RNA-binding complex. *Journal of Neurochemistry* **83** 202–210. (doi:10.1046/j.1471-4159.2002.01158.x)
- Giorgini F, Davies HG & Braun RE 2001 MSY2 and MSY4 bind a conserved sequence in the 3' untranslated region of protamine 1 mRNA *in vitro* and *in vivo*. *Molecular and Cellular Biology* **21** 7010–7019. (doi:10.1128/MCB.21.20.7010-7019.2001)
- Giorgini F, Davies HG & Braun RE 2002 Translational repression by MSY4 inhibits spermatid differentiation in mice. *Development* **129** 3669–3679.
- Hafner M, Landthaler M, Burger L, Khorshid M, Hausser J, Berninger P, Rothballer A, Ascano M Jr, Jungkamp AC, Munschauer M *et al.* 2010 Transcriptome-wide identification of RNA-binding protein and micro-RNA target sites by PAR-CLIP. *Cell* **141** 129–141. (doi:10.1016/j.cell.2010.03.009)
- Han JR, Gu W & Hecht NB 1995 Testis-brain RNA-binding protein, a testicular translational regulatory RNA-binding protein, is present in the brain and binds to the 3' untranslated regions of transported brain mRNAs. *Biology of Reproduction* **53** 707–717. (doi:10.1095/biol-reprod53.3.707)
- Hawthorne SK, Busanelli RR & Kleene KC 2006 The 5'UTR and 3'UTR of the sperm mitochondria-associated cysteine-rich protein mRNA regulate translation in spermatids by multiple mechanisms in transgenic mice. *Developmental Biology* **297** 118–126. (doi:10.1016/j.ydbio.2006.04.468)
- Hinnebusch AG 2005 Translational regulation of GCN4 and the general amino acid control of yeast. *Annual Review of Microbiology* **59** 407–450. (doi:10.1146/annurev.micro.59.031805.133833)
- Hunter D, Anand-Ivell R, Danner S & Ivell R 2012 Models of *in vitro* spermatogenesis. *Spermatogenesis* **2** 32–43. (doi:10.4161/spmg.19383)
- Idler RK & Yan W 2012 Control of messenger RNA fate by RNA binding proteins, an emphasis on mammalian spermatogenesis. *Journal of Andrology* **33** 309–337. (doi:10.2164/jandrol.111.014167)
- Iguchi N, Tobias JW & Hecht NB 2006 Expression profiling reveals meiotic male germ cell mRNAs that are translationally up- and down-regulated. *PNAS* **103** 7712–7717. (doi:10.1073/pnas.0510999103)
- Ivics Z, Izsvák Z, Medrano G, Chapman KM & Hamra FK 2011 Sleeping beauty transposon mutagenesis in rat spermatogonial stem cells. *Nature Protocols* **6** 1521–1535. (doi:10.1038/nprot.2011.378)
- Izaurrealde E 2012 Elucidating the temporal order of silencing. *EMBO Reports* **13** 662–663. (doi:10.1038/embor.2012.91)
- Jackson RJ, Hellen CU & Pestova TV 2010 The mechanism of eukaryotic translation initiation and principles of its regulation. *Nature Reviews. Molecular Cell Biology* **11** 113–127. (doi:10.1038/nrm2838)
- Jaendling A & McFarlane RJ 2010 Biological roles of translin and translin-associated factor-X, RNA metabolism comes to the fore. *Biochemical Journal* **429** 225–234. (doi:10.1042/BJ20100273)
- Janicke A, Vancuylenberg J, Boag PR, Traven A & Beilharz TH 2012 ePAT: a simple method to tag adenylated RNA to measure poly(A)-tail length and other 3' RACE applications. *RNA* **18** 1289–1295. (doi:10.1261/rna.031898.111)
- Jankowsky E 2011 RNA helicases at work: binding and rearranging. *Trends in Biochemical Sciences* **36** 19–29. (doi:10.1016/j.tibs.2010.07.008)
- Jenkins HT, Malkova B & Edwards TA 2011 Kinked β -strands mediate high-affinity recognition of mRNA targets by the germ-cell regulator DAZL. *PNAS* **108** 18266–18271. (doi:10.1073/pnas.1105211108)
- Kanatsu-Shinohara M & Shinohara T 2010 Germline modification using mouse spermatogonial stem cells. *Methods in Enzymology* **477** 17–36. (doi:10.1016/S0076-6879(10)77002-6)
- Kanatsu-Shinohara M, Ogonuki N, Iwano T, Lee J, Kazuki Y, Inoue K, Miki H, Takehashi M, Toyokuni S, Shinkai Y *et al.* 2005 Genetic and epigenetic properties of mouse male germline stem cells during long-term culture. *Development* **132** 4155–4163. (doi:10.1242/dev.02004)
- Kashiwabara S, Noguchi J, Zhuang T, Ohmura K, Honda A, Sugiura S, Miyamoto K, Takahashi S, Inoue K, Ogura A *et al.* 2002 Regulation of spermatogenesis by testis-specific, cytoplasmic poly(A) polymerase TPAP. *Science* **298** 1999–2002. (doi:10.1126/science.1074632)
- Kierszenbaum AL & Tres LL 1975 Structural and transcriptional features of the mouse spermatid genome. *Journal of Cell Biology* **65** 258–270. (doi:10.1083/jcb.65.2.258)
- Kirino Y, Vourekas A, Kim N, de Lima Alves F, Rappsilber J, Klein PS, Jongens TA & Mourelatos Z 2010 Arginine methylation of vasa protein is conserved across phyla. *Journal of Biological Chemistry* **285** 8148–8854. (doi:10.1074/jbc.M109.089821)
- Kishore S, Luber S & Zavolan M 2010 Deciphering the role of RNA-binding proteins in post-transcriptional control of gene expression. *Briefings in Functional Genomics* **9** 391–404. (doi:10.1093/bfpg/elq028)
- Kishore S, Jaskiewicz L, Burger L, Hausser J, Khorshid M & Zavolan M 2011 A quantitative analysis of CLIP methods for identifying binding sites of RNA-binding proteins. *Nature Methods* **8** 559–564. (doi:10.1038/nmeth.1608)
- Kleene KC 1989 Poly(A) shortening accompanies the activation of translation of five mRNAs during spermiogenesis in the mouse. *Development* **106** 367–373.
- Kleene KC 1993 Multiple controls over the efficiency of translation of the mRNAs encoding transition proteins, protamines, and the mitochondrial capsule selenoprotein in late spermatids in mice. *Developmental Biology* **159** 720–731. (doi:10.1006/dbio.1993.1277)
- Kleene KC 2003 Patterns, mechanisms and functions of translational regulation in mammalian spermatogenic cells. *Cytogenetic and Genome Research* **103** 217–224. (doi:10.1159/000076807)
- Kleene KC & Smith J 1994 Translational activity of mouse protamine 1 messenger ribonucleoprotein particles in the reticulocyte and wheat germ cell-free translation systems. *Molecular Reproduction and Development* **37** 12–20. (doi:10.1002/mrd.1080370103)

- Kleene KC, Bagarova J, Hawthorne SK & Catado LM 2010 Quantitative analysis of mRNA translation in mammalian spermatogenic cells with sucrose and Nycodenz gradients. *Reproductive Biology and Endocrinology* **8** 155. (doi:10.1186/1477-7827-8-155)
- Kolmer M, Peltö-Huikko M, Parvinen M, Höög C & Alho H 1997 The transcriptional and translational control of diazepam binding inhibitor expression in rat male germ-line cells. *DNA and Cell Biology* **16** 59–72. (doi:10.1089/dna.1997.16.59)
- Korhonen HM, Meikar O, Yadav RP, Papaioannou MD, Romero Y, Da Ros M, Herrera PL, Toppari J, Nef S & Kotaja N 2011 Dicer1 is required for haploid male germ cell differentiation in mice. *PLoS ONE* **6** e24821. (doi:10.1371/journal.pone.0024821)
- Kotaja N, Kimmins S, Brancorsini S, Hentsch D, Vonesch JL, Davidson I, Parvinen M & Sassone-Corsi P 2004 Preparation, isolation and characterization of stage-specific spermatogenic cells for cellular and molecular analysis. *Nature Methods* **1** 249–254. (doi:10.1038/nmeth1204-249)
- Kramer JM 1981 Immunofluorescent localization of PGK-1 and PGK-2 isozymes within specific cells of the mouse testis. *Developmental Biology* **87** 30–36. (doi:10.1016/0012-1606(81)90057-9)
- Kress C, Gautier-Courteille C, Osborne HB, Babinet C & Paillard L 2007 Inactivation of CUG-BP1/CELF1 causes growth, viability, and spermatogenesis defects in mice. *Molecular and Cellular Biology* **27** 1146–1157. (doi:10.1128/MCB.01009-06)
- Kuramochi-Miyagawa S, Watanabe T, Gotoh K, Takamatsu K, Chuma S, Kojima-Kita K, Shiromoto Y, Asada N, Toyoda A, Fujiyama A *et al.* 2010 MVH in piRNA processing and gene silencing of retrotransposons. *Genes and Development* **24** 887–892. (doi:10.1101/gad.1902110)
- Kwon YK & Hecht NB 1991 Cytoplasmic protein binding to highly conserved sequences in the 3' untranslated region of mouse protamine 2 mRNA, a translationally regulated transcript of male germ cells. *PNAS* **88** 3584–3588. (doi:10.1073/pnas.88.9.3584)
- Kwon YK & Hecht NB 1993 Binding of a phosphoprotein to the 3' untranslated region of the mouse protamine 2 mRNA temporally represses its translation. *Molecular and Cellular Biology* **13** 6547–6557. (doi:10.1128/MCB.13.10.65470)
- Kwon YK, Murray MT & Hecht NB 1993 Proteins homologous to the *Xenopus* germ cell-specific RNA-binding proteins p54/p56 are temporally expressed in mouse male germ cells. *Developmental Biology* **158** 99–100. (doi:10.1006/dbio.1993.1170)
- La Salle S, Sun F & Handel MA 2009 Isolation and short-term culture of mouse spermatocytes for analysis of meiosis. *Methods in Molecular Biology* **558** 279–297. (doi:10.1007/978-1-60761-103-5_17)
- Lebedeva S, Jens M, Theil K, Schwanhäusser B, Selbach M, Landthaler M & Rajewsky N 2011 Transcriptome-wide analysis of regulatory interactions of the RNA-binding protein HuR. *Molecular Cell* **43** 340–352. (doi:10.1016/j.molcel.2011.06.008)
- Lee K, Haugen HS, Clegg CH & Braun RE 1995 Premature translation of protamine 1 mRNA causes precocious nuclear condensation and arrests spermatid differentiation in mice. *PNAS* **92** 12451–12455. (doi:10.1073/pnas.92.26.12451)
- Lee K, Fajardo MA & Braun RE 1996 A testis cytoplasmic RNA-binding protein that has the properties of a translational repressor. *Molecular and Cellular Biology* **16** 3023–3034.
- Levy S, Avni D, Hariharan N, Perry RP & Meyuhas O 1991 Oligopyrimidine tract at the 5' end of mammalian ribosomal protein mRNAs is required for their translational control. *PNAS* **88** 3319–3323. (doi:10.1073/pnas.88.8.3319)
- Li Z & Baraban JM 2004 High affinity binding of the Translin/Trax complex to RNA does not require the presence of Yor H elements. *Brain Research. Molecular Brain Research* **120** 123–129. (doi:10.1016/j.molbrainres.2003.10.010)
- Ling J, Morley SJ, Pain VM, Marzluff WF & Gallie DR 2002 The histone 3'-terminal stem-loop-binding protein enhances translation through a functional and physical interaction with eukaryotic initiation factor 4G (eIF4G) and eIF3. *Molecular and Cellular Biology* **22** 7853–7867. (doi:10.1128/MCB.22.22.7853-7867.2002)
- Liu N, Han H & Lasko P 2009 Vasa promotes *Drosophila* germline stem cell differentiation by activating mei-P26 translation by directly interacting with a (U)-rich motif in its 3' UTR. *Genes and Development* **23** 2742–2752. (doi:10.1101/gad.1820709)
- Lluis M, Hoe W, Schleit J & Robertus J 2010 Analysis of nucleic acid binding by a recombinant translin-trax complex. *Biochemical and Biophysical Research Communications* **396** 709–713. (doi:10.1016/j.bbrc.2010.04.166)
- Lo KC & Domes T 2011 Can we grow sperm? A translational perspective on the current animal and human spermatogenesis models *Asian Journal of Andrology* **13** 677–682. (doi:10.1038/aja.2011.88)
- Lu ZH, Books JT & Ley TJ 2006 Cold shock domain family members YB-1 and MSY4 share essential functions during murine embryogenesis. *Molecular and Cellular Biology* **26** 8410–8417. (doi:10.1128/MCB.01196-06)
- Lyabin DN, Eliseeva IA, Skabkina OV & Ovchinnikov LP 2011 Interplay between Y-box-binding protein 1 (YB-1) and poly(A) binding protein (PABP) in specific regulation of YB-1 mRNA translation. *RNA Biology* **8** 883–892. (doi:10.4161/rna.8.5.16022)
- Mali P, Kaipia A, Kangasniemi M, Toppari J, Sandberg M, Hecht NB & Parvinen M 1989 Stage-specific expression of nucleoprotein mRNAs during rat and mouse spermiogenesis. *Reproduction, Fertility, and Development* **1** 369–382. (doi:10.1071/RD9890369)
- Mastrangelo MA & Kleene KC 2000 Developmental expression of Y-box protein 1 mRNA and alternatively spliced Y-box protein 3 mRNAs in spermatogenic cells in mice. *Molecular Human Reproduction* **6** 779–788. (doi:10.1093/molehr/6.9.779)
- Mathews MB, Sonenberg N & Hershey JWB 2007 Origins and principles of translational control. In *Translational Control in Biology and Medicine*, pp 1–40. Eds MB Mathews, N Sonenberg & JWB Hershey. Cold Spring Harbor, NY: Cold Spring Harbor Press.
- McCarrey JR, Berg WM, Paragioudakis SJ, Zhang PL, Dilworth DD, Arnold BL & Rossi JJ 1992 Differential transcription of Pdg genes during spermatogenesis in the mouse. *Developmental Biology* **154** 160–168. (doi:10.1016/0012-1606(92)90056-M)
- McGrew LL, Dworkin-Rastl E, Dworkin MB & Richter JD 1989 Poly(A) elongation during *Xenopus* oocyte maturation is required for translational recruitment and is mediated by a short sequence element. *Genes and Development* **3** 803–815. (doi:10.1101/gad.3.6.803)
- Medenbach J, Seiler M & Hentze MW 2011 Translational control via protein-regulated upstream open reading frames. *Cell* **145** 902–913. (doi:10.1016/j.cell.2011.05.005)
- Mehta A, Trotta CR & Peltz SW 2006 Derepression of the Her-2 uORF is mediated by a novel post-transcriptional control mechanism in cancer cells. *Genes and Development* **20** 939–953. (doi:10.1101/gad.1388706)
- Meikar O, Da Ros M, Korhonen H & Kotaja N 2011 Chromatoid body and small RNAs in male germ cells. *Reproduction* **142** 195–209. (doi:10.1530/REP-11-0057)
- Meistrich ML, Mohapatra B, Shirley CR & Zhao M 2003 Roles of transition nuclear proteins in spermiogenesis. *Chromosoma* **111** 483–488. (doi:10.1007/s00412-002-0227-z)
- Messina V, Di Sauro A, Pedrotti S, Adesso L, Latina A, Geremia R, Rossi P & Sette C 2010 Differential contribution of the MTOR and MNK pathways to the regulation of mRNA translation in meiotic and postmeiotic mouse male germ cells. *Biology of Reproduction* **83** 607–615. (doi:10.1095/biolreprod.110.085050)
- Mestdagh P, Boström AK, Impens F, Fredlund E, Van Peer G, De Antonellis P, von Stedingk K, Ghesquière B, Schulte S, Dews M *et al.* 2010 The miR-17-92 microRNA cluster regulates multiple components of the TGF- β pathway in neuroblastoma. *Molecular Cell* **40** 762–773. (doi:10.1016/j.molcel.2010.11.038)
- Morris AR, Mukherjee N & Keene JD 2010 Systematic analysis of posttranscriptional gene expression. *Wiley Interdisciplinary Reviews. Systems Biology and Medicine* **2** 162–180. (doi:10.1002/wsbm.54)
- Mueller PP & Hinnebusch AG 1986 Multiple upstream AUG codons mediate translational control of GCN4. *Cell* **45** 201–207. (doi:10.1016/0092-8674(86)90384-3)
- Mukherjee N, Corcoran DL, Nusbaum JD, Reid DW, Georgiev S, Hafner M, Ascano M Jr, Tuschl T, Ohler U & Keene JD 2011 Integrative regulatory mapping indicates that the RNA-binding protein HuR couples pre-mRNA processing and mRNA stability. *Molecular Cell* **43** 327–339. (doi:10.1016/j.molcel.2011.06.007)
- Nagamori I, Cruickshank VA & Sassone-Corsi P 2011 Regulation of an RNA granule during spermatogenesis: acetylation of MVH in the chromatoid body of germ cells. *Journal of Cell Science* **124** 4346–4355. (doi:10.1242/jcs.096461)

- Nguyen-Chi M & Morello D 2011 RNA-binding proteins, RNA granules, and gametes: is unity strength? *Reproduction* **142** 803–817. (doi:10.1530/REP-11-0257)
- Nguyen Chi M, Chalmel F, Agius E, Vanzo N, Khabar KS, Jégou B & Morello D 2009 Temporally regulated traffic of HuR and its associated ARE-containing mRNAs from the chromatoid body to polysomes during mouse spermatogenesis. *PLoS ONE* **4** e4900. (doi:10.1371/journal.pone.0004900)
- Oko R, Korley R, Murray MT, Hecht NB & Hermo L 1996 Germ cell-specific DNA and RNA binding proteins p48/52 are expressed at specific stages of male germ cell development and are present in the chromatoid body. *Molecular Reproduction and Development* **44** 1–13. (doi:10.1002/(SICI)1098-2795(199605)44:1<1::AID-MRD1>3.0.CO;2-S)
- Onohara Y & Yokota S 2012 Expression of DDX25 in nuage components of mammalian spermatogenic cells: immunofluorescence and immunoelectron microscopic study. *Histochemistry and Cell Biology* **137** 37–51. (doi:10.1007/s00418-011-0875-2)
- Onohara Y, Fujiwara T, Yasukochi T, Himeno M & Yokota S 2010 Localization of mouse vasa homolog protein in chromatoid body and related nuage structures of mammalian spermatogenic cells during spermatogenesis. *Histochemistry and Cell Biology* **133** 627–639. (doi:10.1007/s00418-010-0699-5)
- Ostareck DH, Ostareck-Lederer A, Shatsky IN & Hentze MW 2001 Lipoygenase mRNA silencing in erythroid differentiation. The 3'UTR regulatory complex controls 60S ribosomal subunit joining. *Cell* **104** 281–290. (doi:10.1016/S0092-8674(01)00212-4)
- Paronetto MP & Sette C 2010 Role of RNA-binding proteins in mammalian spermatogenesis. *International Journal of Andrology* **33** 2–12. (doi:10.1111/j.1365-2605.2009.00959.x)
- Paronetto MP, Messina V, Bianchi E, Barchi M, Vogel G, Moretti C, Palombi F, Stefanini M, Geremia R, Richard S *et al.* 2009 Sam68 regulates translation of target mRNAs in male germ cells, necessary for mouse spermatogenesis. *Journal of Cell Biology* **185** 235–249. (doi:10.1083/jcb.200811138)
- Parsyan A, Svitkin Y, Shahbazian D, Gkogkas C, Lasko P, Merrick WC & Sonenberg N 2011 mRNA helicases: the tacticians of translational control. *Nature Reviews. Molecular Cell Biology* **12** 235–245. (doi:10.1038/nrm3083)
- Pires-daSilva A, Nayernia K, Engel W, Torres M, Stoykova A, Chowdhury K & Gruss P 2001 Mice deficient for spermatid perinuclear RNA-binding protein show neurologic, spermatogenic, and sperm morphological abnormalities. *Developmental Biology* **233** 319–328. (doi:10.1006/dbio.2001.0169)
- Pusch W, Balvers M, Weinbauer GF & Ivell R 2000 The rat endozepine-like peptide gene is highly expressed in late haploid stages of male germ cell development. *Biology of Reproduction* **63** 763–768. (doi:10.1095/biolreprod63.3.763)
- Radford HE, Meijer HA & de Moor CH 2008 Translational control by cytoplasmic polyadenylation in *Xenopus* oocytes. *Biochimica et Biophysica Acta* **1779** 217–229. (doi:10.1016/j.bbagr.2008.02.002)
- Ray D, Kazan H, Chan ET, Peña Castillo L, Chaudhry S, Talukder S, Blencowe BJ, Morris Q & Hughes TR 2009 Rapid and systematic analysis of the RNA recognition specificities of RNA-binding proteins. *Nature Biotechnology* **27** 667–670. (doi:10.1038/nbt.1550)
- Reynolds N & Cooke HJ 2005 Role of the DAZ genes in male fertility. *Reproductive BioMedicine Online* **10** 72–80. (doi:10.1016/S1472-6483(10)60806-1)
- Reynolds N, Collier B, Maratou K, Bingham V, Speed RM, Taggart M, Semple CA, Gray NK & Cooke HJ 2005 Dazl binds *in vivo* to specific transcripts and can regulate the pre-meiotic translation of Mvh in germ cells. *Human Molecular Genetics* **14** 3899–3909. (doi:10.1093/hmg/ddi414)
- Reynolds N, Collier B, Bingham V, Gray NK & Cooke HJ 2007 Translation of the synaptonemal complex component Sycp3 is enhanced *in vivo* by the germ cell specific regulator Dazl. *RNA* **13** 974–981. (doi:10.1261/ra.465507)
- Romero Y, Meikar O, Papaioannou MD, Conne B, Grey C, Weier M, Pralong F, De Massy B, Kaessmann H, Vassalli JD *et al.* 2011 Dicer11 depletion in male germ cells leads to infertility due to cumulative meiotic and spermiogenic defects. *PLoS ONE* **6** e25241. (doi:10.1371/journal.pone.0025241)
- Russell LD, Ettlin RA, Sinha Hikim AP & Clegg ED 1990 *Histological and Histopathological Evaluation of the Testis*. Clearwater, FL: Cache River Press.
- Sachs MS & Geballe AP 2006 Downstream control of upstream open reading frames. *Genes and Development* **20** 915–921. (doi:10.1101/gad.1427006)
- Saunders PT, Turner JM, Ruggiu M, Taggart M, Burgoyne PS, Elliott D & Cooke HJ 2003 Absence of mDazl produces a final block on germ cell development at meiosis. *Reproduction* **126** 589–597. (doi:10.1530/rep.0.1260589)
- Schmidt EE, Hanson ES & Capecchi MR 1999 Sequence-independent assembly of spermatid mRNAs into messenger ribonucleoprotein particles. *Molecular and Cellular Biology* **19** 3904–3915.
- Shih DM & Kleene KC 1992 A study by *in situ* hybridization of the stage of appearance and disappearance of the transition protein 2 and the mitochondrial capsule seleno-protein mRNAs during spermatogenesis in the mouse. *Molecular Reproduction and Development* **33** 222–227. (doi:10.1002/mrd.1080330216)
- Shyu AB, Greenberg ME & Belasco JG 1989 The c-fos transcript is targeted for rapid decay by two distinct mRNA degradation pathways. *Genes and Development* **3** 60–72. (doi:10.1101/gad.3.1.60)
- Skabkin MA, Liabin DN & Ovchinnikov LP 2006 Nonspecific and specific interaction of Y-box binding protein 1 (YB-1) with mRNA and posttranscriptional regulation of protein synthesis in animal cells. *Molecular Biology* **40** 620–633. (doi:10.1134/S0026893306040078)
- Song R, Hennig GW, Wu Q, Jose C, Zheng H & Yan W 2011 Male germ cells express abundant endogenous siRNAs. *PNAS* **108** 13159–13164. (doi:10.1073/pnas.1108567108)
- Steel LF, Telly DL, Leonard J, Rice BA, Monks B & Sawicki JA 1996 Elements in the murine c-mos messenger RNA 5'-untranslated region repress translation of downstream coding sequences. *Cell Growth and Differentiation* **7** 1415–1424.
- Sun F, Palmer K & Handel MA 2010 Mutation of Eif4g3, encoding a eukaryotic translation initiation factor, causes male infertility and meiotic arrest of mouse spermatocytes. *Development* **137** 1699–1707. (doi:10.1242/dev.043125)
- Svitkin YV, Evdokimova VM, Brasey A, Pestova TV, Fantus D, Yanagiya A, Imataka H, Skabkin MA, Ovchinnikov LP, Merrick WC *et al.* 2009 General RNA-binding proteins have a function in poly(A)-binding protein-dependent translation. *EMBO Journal* **28** 58–68. (doi:10.1038/emboj.2008.259)
- Tanaka SS, Toyooka Y, Akasu R, Katoh-Fukui Y, Nakahara Y, Suzuki R, Yokoyama M & Noce T 2000 The mouse homolog of *Drosophila* Vasa is required for the development of male germ cells. *Genes and Development* **14** 841–853. (doi:10.1101/gad.14.7.841)
- Tay J & Richter JD 2001 Germ cell differentiation and synaptonemal complex formation are disrupted in CPEB1 knockout mice. *Developmental Cell* **1** 201–213. (doi:10.1016/S1534-5807(01)00025-9)
- Tian Y, Simanshu DK, Ascano M, Diaz-Avalos R, Park AY, Juranek SA, Rice WJ, Yin Q, Robinson CV, Tuschl T *et al.* 2011 Multimeric assembly and biochemical characterization of the Trax-translin endonuclease complex. *Nature Structural & Molecular Biology* **18** 658–664. (doi:10.1038/nsmb.2069)
- Tsai-Morris CH, Sheng Y, Lee E, Lei KJ & Dufau ML 2004 Gonadotropin-regulated testicular RNA helicase (GRTH/Ddx25) is essential for spermatid development and completion of spermatogenesis. *PNAS* **101** 6373–6378. (doi:10.1073/pnas.0401855101)
- Tsai-Morris CH, Sato H, Guttli R & Dufau ML 2012 Role of gonadotropin regulated testicular RNA helicase (GRTH/Ddx25) on polysomal associated mRNAs in mouse testis. *PLoS ONE* **7** e32470. (doi:10.1371/journal.pone.0032470)
- Tseden K, Topaloglu O, Meinhardt A, Dev A, Adham I, Müller C, Wolf S, Böhm D, Schlüter G, Engel W *et al.* 2007 Premature translation of transition protein 2 mRNA causes sperm abnormalities and male infertility. *Molecular Reproduction and Development* **74** 273–279. (doi:10.1002/mrd.20570)
- Ule J, Jensen K, Mele A & Darnell RB 2005 CLIP: a method for identifying protein–RNA interaction sites in living cells. *Methods* **37** 376–386. (doi:10.1016/j.ymeth.2005.07.018)

- Vandeberg JL, Lee CY & Goldberg E 1981 Immunohistochemical localization of phosphoglycerate kinase isozymes in mouse testes. *Journal of Experimental Zoology* **217** 435–441. (doi:10.1002/jez.1402170315)
- VanGompel MJ & Xu EY 2010 A novel requirement in mammalian spermatid differentiation for the DAZ-family protein Boule. *Human Molecular Genetics* **19** 2360–2369. (doi:10.1093/hmg/ddq109)
- VanGompel MJ & Xu EY 2011 The roles of the DAZ family in spermatogenesis: more than just translation? *Spermatogenesis* **1** 36–46. (doi:10.4161/spmg.1.1.14659)
- Venables JP, Ruggiu M & Cooke HJ 2001 The RNA-binding specificity of the mouse Dazl protein. *Nucleic Acids Research* **29** 2479–2483. (doi:10.1093/nar/29.12.2479)
- Ventelä S, Okabe M, Tanaka H, Nishimune Y, Toppari J & Parvinen M 2000a Expression of green fluorescent protein under β -actin promoter in living spermatogenic cells of the mouse: stage-specific regulation by FSH. *International Journal of Andrology* **23** 236–242. (doi:10.1046/j.1365-2605.2000.00237.x)
- Ventelä S, Mulari M, Okabe M, Tanaka H, Nishimune Y, Toppari J & Parvinen M 2000b Regulation of acrosome formation in mice expressing green fluorescent protein as a marker. *Tissue & Cell* **32** 501–507. (doi:10.1016/S0040-8166(00)80006-3)
- Wu XQ, Gu W, Meng X & Hecht NB 1997 The RNA-binding protein, TB-RBP, is the mouse homologue of translin, a recombination protein associated with chromosomal translocations. *PNAS* **94** 5640–5645. (doi:10.1073/pnas.94.11.5640)
- Yanagiya A, Delbes G, Svitkin YV, Robaire B & Sonenberg N 2010 The poly(A)-binding protein partner Paip2a controls translation during late spermiogenesis in mice. *Journal of Clinical Investigation* **120** 3389–3400. (doi:10.1172/JCI43350)
- Yang J, Chennathukuzhi V, Miki K, O'Brien DA & Hecht NB 2003 Mouse testis brain RNA-binding protein/translin selectively binds to the messenger RNA of the fibrous sheath protein glyceraldehyde 3-phosphate dehydrogenase-S and suppresses its translation *in vitro*. *Biology of Reproduction* **68** 853–859. (doi:10.1095/biolreprod.102.008631)
- Yang J, Morales CR, Medvedev S, Schultz RM & Hecht NB 2007 In the absence of the mouse DNA/RNA-binding protein MSY2, messenger RNA instability leads to spermatogenic arrest. *Biology of Reproduction* **76** 48–54. (doi:10.1095/biolreprod.106.055095)
- Ye X, Huang N, Liu Y, Paroo Z, Huerta C, Li P, Chen S, Liu Q & Zhang H 2011 Structure of C3PO and mechanism of human RISC activation. *Nature Structural & Molecular Biology* **18** 650–657. (doi:10.1038/nsmb.2032)
- Yoshioka H, Geyer CB, Hornecker JL, Patel KT & McCarrey JR 2007 *In vivo* analysis of developmentally and evolutionarily dynamic protein–DNA interactions regulating transcription of the Pdgfra gene during mammalian spermatogenesis. *Molecular and Cellular Biology* **27** 7871–7885. (doi:10.1128/MCB.00990-07)
- Yuan L, Liu JG, Zhao J, Brundell E, Daneholt B & Höög C 2000 The murine SCP3 gene is required for synaptonemal complex assembly, chromosome synapsis, and male fertility. *Molecular Cell* **5** 73–83. (doi:10.1016/S1097-2765(00)80404-9)
- Zhang C & Darnell RB 2011 Mapping *in vivo* protein–RNA interactions at single-nucleotide resolution from HITS-CLIP data. *Nature Biotechnology* **29** 607–614. (doi:10.1038/nbt.1873)
- Zhong J, Peters AH, Lee K & Braun RE 1999 A double-stranded RNA binding protein required for activation of repressed messages in mammalian germ cells. *Nature Genetics* **22** 171–174. (doi:10.1038/9684)
- Zhong J, Peters AH, Kafer K & Braun RE 2001 A highly conserved sequence essential for translation repression of the protamine 1 messenger RNA in murine spermatids. *Biology of Reproduction* **64** 1784–1789. (doi:10.1095/biolreprod.64.6.1784)

Received 17 September 2012

First decision 29 October 2012

Revised manuscript received 25 February 2013

Accepted 10 April 2013