Oviductal secretion and gamete interaction

Sergio Ghersevich, Estefanía Massa and Carlos Zumoffen

Laboratory of Reproductive Studies, Area of Clinical Biochemistry, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531, 2000 Rosario, Argentina

Correspondence should be addressed to S Ghersevich; Email: sghersev@fbioyf.unr.edu.ar

Abstract

Experimental evidence from the last 30 years supports the fact that the oviduct is involved in the modulation of the reproductive process in eutherian mammals. Oviductal secretion contains molecules that contribute to regulation of gamete function, gamete interaction, and the early stages of embryo development. The oviductal environment would act as a sperm reservoir, maintaining sperm viability, and modulating the subpopulation of spermatozoa that initiates the capacitation process. It could also contribute to prevent the premature acrosome reaction and to reduce polyspermy. Many studies have reported the beneficial effects of the oviductal environment on fertilization and on the first stages of embryo development. Some oviductal factors have been identified in different mammalian species. The effects of oviductal secretion on the reproductive process could be thought to result from the dynamic combined action (inhibitory or stimulatory) of multiple factors present in the oviductal lumen at different stages of the ovulatory cycle and in the presence of gametes or embryos. It could be hypothesized that the absence of a given molecule would not affect fertility as its action could be compensated by another factor with similar functions. However, any alteration in this balance could affect certain events of the reproductive process and could perhaps impair fertility. Thus, the complexity of the reproductive process warrants a continuous research effort to unveil the mechanisms and factors behind its regulation in the oviductal microenvironment.

Reproduction (2015) **149** R1–R14

The oviduct

The oviduct of eutherian mammals is the organ where the fertilization process takes place. It is a seromuscular tubular organ, whose distal part surrounds the ovary and whose proximal portion is attached to the uterus. Based on morphological and anatomical differences, the oviduct can be divided into four segments (composed of similar cell types, but in different proportions): the infundibulum (with the fimbriae surrounding the ovary), the ampulla, the isthmus, and the uterotubal junction (Fig. 1; Hess *et al.* 2006, Suarez 2006).

The ampulla is the site of oocyte fertilization. The mucosa of the ampulla shows a very complex pattern of folds, which are projected toward the oviductal lumen (Hess *et al.* 2006, Suarez 2006). The isthmic mucosa contains fewer folds than the ampulla.

The oviductal epithelium is mainly formed by columnar ciliated cells and secretory cells (showing surface microvilli); lymphoid cells can be observed close to the basement membrane. The secretory nonciliated columnar cells present a typical structure of cells that actively synthesize proteins (Suarez 2006). Marked differences in the morphological characteristics of the secretory granules from oviductal cells were reported in different species analyzed. In cattle, during the follicular

phase of the estrous cycle, nonciliated cells of the ampulla and fimbriae contain large amount of secretory granules, while in the isthmus the number of cytoplasm granules was smaller and show different structural characteristics (Killian 2011).

Primate oviductal epithelial cells possess estrogen and progesterone receptors and undergo cyclical changes related to the menstrual cycle (Brenner *et al.* 1990, Hess *et al.* 2006). In the presence of progesterone, after estradiol levels decrease, loss of *ciliated* epithelium occurs and the secretory cells tend to lose their biosynthetic structures. Estrogens stimulate the secretion of the oviductal epithelium and this secretion is highest in the proliferative phase (Lippes *et al.* 1981, Suarez 2006, Killian 2011). The oviductal fluid contains amino acids, proteins, simple, and complex carbohydrates, ions, lipids, and phospholipids. Some of these components are metabolic substrates, such as lactate, pyruvic acid, amino acids, and glucose, whose levels differ from those present in the uterine fluid and the serum (Leese 1988, Leese *et al.* 2008, Hugentobler *et al.* 2010). Experimental evidence indicated that ion concentrations in oviductal fluid also differ from those of serum, suggesting that the oviductal epithelium modulates ion levels (Leese 1988, Leese *et al.* 2008, Hugentobler *et al.* 2010).

Figure 1 Schematic representation of the oviduct and its suggested involvement in the reproductive process.

The volume and some protein components of the oviductal fluid change throughout the cycle. Part of the complex mixture of proteins that are present in the oviductal fluid come from serum transudate, but there are also specific proteins synthesized and secreted by the oviductal epithelium, and some of them could be regulated by cyclic hormonal changes, with increased biosynthesis at the periovulatory period (Buhi *et al.* 2000, Buhi 2002).

Gamete transport within the oviductal lumen would be a highly controlled process (Suarez 2006, Kölle *et al.* 2009). There is evidence showing that the ciliary activity of the oviductal epithelial cells plays a key role in gamete and embryo transportation. It has been determined that the frequency of ciliary movement of the human oviductal epithelium increases after ovulation (Suarez 2006, Kölle *et al.* 2009).

The following pages will present a review of experimental data that support the central role of the oviduct and its secretion in the reproductive process (Fig. 2).

Sperm–oviduct epithelial cell interaction

Once spermatozoa reach the oviduct, they could follow two pathways. Some of them quickly migrate to the ampulla region, and usually they are not able to fertilize the oocyte, while most of spermatozoa are retained in the isthmus region forming a sperm reservoir, in the presence of the oviductal fluid (Figs 1 and 2; Croxatto 2002, Suarez 2008a). Some spermatozoa from the reservoir will retain their viability and their fertilizing ability until ovulation takes place (Suarez 2008a).

In different mammalian species, it has been shown that spermatozoa sequestered in the isthmus region could attach to epithelial cells, delaying sperm capacitation until ovulation-associated signals induce their release, allowing their transit to the ampulla. Such interaction would involve the sperm acrosomal region and the apical region of the oviduct epithelial cells, mainly of the ciliated cells (Petrunkina *et al.* 2001,

Croxatto 2002, Suarez 2008a, Coy *et al.* 2012a). In some mammals, such as canine, cattle, and horses, the interaction between spermatozoa and oviduct epithelial cells appears to be associated with sperm survival and capacitation state (Kawakami *et al.* 2001, Suarez 2006, 2008b). The molecular and biological mechanisms behind the sperm–epithelium interaction have not been clarified yet.

Different studies have investigated the effects of oviductal molecules that could be involved in the sperm–oviduct interaction. Members of the annexin family of proteins detected in the oviductal epithelium were identified as potential receptors for sperm proteins in bovine and porcine species (Ignatz *et al.* 2007, Suarez *et al.* 2008a, Teijeiro *et al.* 2009). Results from a recent study on pigs suggested that sperm binding to oviduct requires the presence of proteins with 6-sialylated biantennary glycans in the membrane of epithelial cells (Kadirvel *et al.* 2012).

Participation of molecules from oviductal secretion in sperm–oviduct interaction

Heparin and other sulfated glycoconjugates, which could be detected in oviductal fluid, were shown to induce the synchronous release of sperm adhering to bovine oviduct epithelium *in vitro*. Heparin-released sperm showed significantly higher intracellular Ca^{++} levels and increased levels of tyrosine-phosphorylated proteins compared with adhering spermatozoa (Gualtieri *et al.* 2005). Another study reported that uncapacitated bovine spermatozoa adhered to the oviduct, and their release was associated with capacitation (Talevi *et al.* 2007). The authors suggested that thiol-reducing agents in the oviductal fluid, such as sulfated glycoconjugates and disulfide reductants, may modulate the redox status of sperm surface proteins, leading to the release of

Figure 2 Reported effects of the oviductal secretions on gamete function and gamete interaction in different experimental models. The symbols '+' and '-' indicate stimulatory or inhibitory effects respectively.

spermatozoa from the oviductal epithelium through the reduction of sperm surface protein disulfides to sulfhydryls (Talevi *et al.* 2007, Gualtieri *et al.* 2010).

An endocannabinoid known as anandamide is synthesized in the oviductal epithelia and detected in the oviductal fluid, and binds to cannabinoid receptors. These receptors were detected in mammalian reproductive tissues and male gametes from different species, such as human, porcine, and bovine species (Schuel *et al.* 2002a,b, Maccarrone *et al.* 2005, Rossato *et al.* 2005, Gervasi *et al.* 2009, Osycka-Salut *et al.* 2012). Gervasi *et al.* (2013) reported that anandamide levels in bovine oviductal fluid varied during estrous cycle, with the highest values detected during the periovulatory period, suggesting that this endocannabinoid levels may be regulated by ovarian hormones. It was reported that anandamide could modulate the bovine sperm–oviductal epithelium interaction, by inhibiting sperm binding and inducing sperm release from epithelial cells (Gervasi *et al.* 2009, Osycka-Salut *et al.* 2012). However, other authors observed that anandamide had no effects on sperm–oviduct binding and sperm release, but might contribute to the oviduct sperm-reservoir function, decreasing motility and capacitation, and prolonging sperm fertile life (Talevi *et al.* 2010).

Oviductal secretion and sperm function

Different experimental approaches have been used in order to assess the effects of the oviductal fluid on gamete function, fertilization, and the initial embryo development.

In some studies, the oviductal fluid was collected by placing a cannula in the oviduct (Leese 1988, Kavanaugh *et al.* 1992, Grippo *et al.* 1995, Way *et al.* 1997, Kumaresan *et al.* 2012).

In addition, the conditioned media from cultures of oviductal cells or tissue explants from different species were used to study the effects of the oviductal secretion on gamete function (Verhage *et al.* 1988, Yeung *et al.* 1994, Yao *et al.* 2000, Quintero *et al.* 2005, Munuce *et al.* 2009, Zumoffen *et al.* 2010).

Numerous studies in mammals have shown that the oviductal secretion or some of its components were able to modulate sperm function, sperm–zona interaction, and the process of fertilization (Fig. 2; Rodríguez-Martínez 2007, Mugnier *et al.* 2009, Killian 2011).

Co-incubation of spermatozoa with oviduct epithelial cells or their conditioned media maintained sperm viability and motility (Zhu *et al.* 1994, Kervancioglu *et al.* 2000, Quintero *et al.* 2005, Munuce *et al.* 2009, Zumoffen *et al.* 2010). However, the mechanisms by which oviductal cells and their secretory products favor the survival of spermatozoa remain unknown.

It has been reported that spermatozoa incubated in the presence of oviductal fluid or co-cultured with oviduct epithelial cells showed a pattern of hyperactivated

motility related to the capacitation state, and it would be a preliminary step to undergo the acrosome reaction (Suarez, 2006, 2008b). It has also been observed that co-culture of sperm with oviductal cells or incubation in the presence of oviductal fluid showed beneficial effects on the human sperm membrane stabilization (Zhu *et al.* 1994, Yao *et al.* 1999). The decrease in human sperm response to acrosome reaction inducers in the presence of the oviductal cells or their conditioned media suggests that oviductal secretion could exert a stabilizing effect on sperm, contributing to avoid a premature acrosome reaction in the absence of the female gamete (Morales *et al.* 1996, Yao *et al.* 1999). Results from our laboratory indicated a significant decrease in the ionophore-induced acrosome reaction in human sperm incubated previously in the presence of a conditioned medium from human oviductal tissue cultures (Quintero *et al.* 2005). Another study showed that the exposure of boar sperm to porcine oviductal fluid collected in the follicular phase of the estrous cycle promoted boar sperm viability and acrosomal integrity (Coy *et al.* 2010). A recent study has reported that incubation of boar sperm in the presence of pre-ovulatory oviductal fluid caused a significant increase in sperm protein tyrosine phosphorylation compared with incubation with post-ovulatory oviductal fluid or the control medium (Kumaresan *et al.* 2012). In addition, the observed phosphorylation patterns appeared to be individual dependent in porcine species.

Numerous research studies have investigated the oviductal factors that could be involved in the reported effects. The presence of adequate levels of bicarbonate and calcium seems to be essential in regulating sperm capacitation, motility, and acrosome reaction (Rodríguez-Martínez 2007, Leese *et al.* 2008, Abou-haila & Tulsiani 2009, Lishko *et al.* 2012). Hyaluronan and sulfated glycosaminoglycans, such as heparin, have been detected in oviductal fluid from different mammalian species (Kawakami *et al.* 2000, Tienthai *et al.* 2000, 2004, Bergqvist & Rodríguez-Martínez 2006). Glycosaminoglycans were involved in modulating sperm viability and capacitation in some species (Kawakami *et al.* 2000, Tienthai *et al.* 2000, Bergqvist & Rodríguez-Martínez 2006).

Transuded serum proteins, such as albumin and HDLs, are found in the oviductal fluid and would be involved in the cholesterol efflux from sperm membrane during capacitation (Travis & Kopf 2002, Leese *et al.* 2008, Abou-haila & Tulsiani 2009).

Involvement of oviductal proteins in sperm function

The oviduct synthesizes and secretes proteins whose functions have not been clarified with certainty (Leese *et al.* 2008, Killian 2011). Notorious differences in the proteomic profiles of the oviduct between follicular and luteal phases of the reproductive cycle in pig were

reported (Seytanoglu *et al.* 2008). In addition, the presence of gametes appears to alter the pattern of protein synthesis and secretion of the oviductal cells. Supporting this idea, the binding of equine spermatozoa to homologous oviductal cells could change the pattern of protein secretion of the epithelial cells (Thomas *et al.* 1995a). The authors suggested that, during this interaction, spermatozoa would be exposed to oviductal protein factors that would maintain their viability and motility and would also facilitate the elimination of deleterious metabolic products (Thomas *et al.* 1995b). Georgiou *et al.* (2005, 2007) also showed that the presence of spermatozoa in the pig oviduct could alter the expression and secretion of specific oviductal proteins.

Recently, we have shown that proteins from conditioned media of oviduct tissue culture decreased the follicular fluid-induced acrosome reaction and the phosphorylation in tyrosine residues of human sperm proteins in a dose-dependent manner (Zumoffen *et al.* 2010).

In the last 30 years, several investigations have been directed to identify proteins from the oviductal fluid to analyze their potential effects on the reproductive process. Some of these oviductal proteins and their reported actions will be described as follows.

Oviductins

The production and secretion of high-molecular-weight (MW 70–130 kDa) glycoproteins from the oviductal epithelium seem to be associated with hormonal changes during the ovulatory cycle in different species (Buhi *et al.* 2000, Buhi 2002, Leese *et al.* 2008, Avilés *et al.* 2010). Oviductins, also known as oviduct-specific glycoproteins, have been found in the oviducts of every mammalian species studied to date, and the cDNA sequences for these glycoproteins indicated that they show a high homology among species (Donnelly *et al.* 1991, Arias *et al.* 1994, Sendai *et al.* 1995, Buhi *et al.* 1996). Bovine oviductin, referred to as an estrus-associated protein, was isolated from the oviductal fluid and was able to bind to the head and middle piece of spermatozoa (King & Killian 1994). Bovine oviductin has been shown to promote *in vitro* sperm capacitation and maintain both the viability and motility of spermatozoa compared with the control medium without any added protein and these effects were dose dependent (King *et al.* 1994, Abe *et al.* 1995). Mouse oviductin has been shown to bind to the equatorial and acrosomal regions of mouse sperm heads (Lyng & Shur 2009). Other studies indicated that hamster oviductin bound to the anterior acrosomal region of the sperm and enhanced sperm capacitation (Kimura *et al.* 1994, Saccary *et al.* 2013). In addition, it was reported that the sperm-binding sites of hamster oviductin were related to the sperm capacitation status and the acrosome reaction (Kan & Esperanzate 2006).

Osteopontin

This phosphoprotein, which contains repetitive amino acid sequences of arginine–glycine–aspartic acid (RGD), was detected in the bovine oviductal secretion and shown to have a variable expression throughout the estrous cycle (Gabler *et al.* 2003). Glycoproteins containing the RGD recognition sequence, which would be ligands of integrins, could be present on the extracellular cover of bovine oocytes and sperm (Ikawa *et al.* 2010). A study by Souza *et al.* (2008) revealed that distribution of sperm-binding sites of osteopontin changed after incubation in the bovine oviductal fluid, and the authors suggested that the protein could participate in sperm–oocyte interaction. Another study has reported that osteopontin increased bovine sperm capacitation (Monaco *et al.* 2009).

Glycodelins

These glycoproteins have been detected in the human oviduct at least in four isoforms, namely glycodelin S, glycodelin A, glycodelin F, and glycodelin C, based on the differences in glycosylation (Yeung *et al.* 2006, Chiu *et al.* 2007a, b). Glycodelins are highly homologous to beta-lactoglobulins, which were detected in the female reproductive tract from several species (Huhtala *et al.* 1987). Glycodelin A is produced and secreted by the oviductal epithelium. Recombinant glycodelin A was shown to inhibit capacitation of human and hamster spermatozoa (Dutta *et al.* 2001). Glycodelin F is expressed in the human oviductal epithelium and in granulosa luteal cells and was shown to bind to the acrosomal region of human sperm head and suppress the progesterone-induced AR (Yeung *et al.* 2006). Thus, it was suggested that glycodelin F could help to prevent a premature acrosome reaction, before the human spermatozoa contact the zona pellucida (ZP; Yeung *et al.* 2007). Glycodelin C, generated from glycodelins A and F in oocyte cumulus cells, has been shown to bind to sperm head, mainly in the equatorial region (Chiu *et al.* 2007b).

Atrial natriuretic peptide

This peptide is expressed in the pig oviductal epithelium and is present in the oviductal fluid. Its receptor was detected in spermatozoa (Zhang *et al.* 2006). The pre-incubation of boar spermatozoa with atrial natriuretic peptide has been shown to induce the acrosome reaction (Zhang *et al.* 2006).

Sperm adhesion molecule 1 (SPAM1)

This protein is a hyaluronidase also known as PH-20 present in sperm membrane. It has been found to be secreted in the female reproductive tract (including the oviduct) and it can bind to sperm (Griffiths *et al.* 2008). It was shown that hyaluronic acid interacts with the

PH-20 protein anchored on the macaque and human sperm membrane increasing the basal levels of intracellular calcium and promoting the induced acrosome reaction (Sabeur *et al.* 1998, Cherr *et al.* 1999).

Lactoferrin

This glycoprotein was isolated in our laboratory from the conditioned medium of human oviductal tissue culture based on its binding ability to sperm membrane and was further identified as human lactoferrin (Zumoffen *et al.* 2013). It was detected in tubal fluid and appeared to be estrogen regulated in human oviduct epithelial cells (Zumoffen *et al.* 2013). Results of our study indicated that lactoferrin presents different binding patterns to sperm related to the capacitation status and the acrosome reaction, suggesting that the protein could participate in sperm–oocyte interaction (Fig. 3; Zumoffen *et al.* 2013). In addition, lactoferrin was able to modulate some sperm functions related to capacitation (C Zumoffen and S Ghersevich unpublished observations).

Figure 3 Fluorescence micrographs showing the acrosomal staining (detected with *Pisum sativum*-rhodamine) and the binding of FITC-conjugated lactoferrin in human spermatozoa incubated under capacitating conditions. (A) Intact acrosome sperm. (B) Acrosome-reacted sperm. (C) Lactoferrin localized to the sperm head in the same cell as (A). (D) Lactoferrin was detected mainly in the equatorial segment in the same spermatozoon as (B).

Involvement of oviductal secretion in gamete interaction

Numerous studies have investigated the effects of the oviductal secretion on gamete interaction. Experiments conducted on cows have shown that the presence of nonluteal oviductal fluid or fluid collected from the isthmic oviductal region increased more sperm–zona binding than the fluid collected in the luteal stage of the estrous cycle or the fluid from the ampulla respectively (Grippio *et al.* 1995, Way *et al.* 1997). The authors suggested that the effects of bovine oviductal fluid on gamete function and interaction depended on the oviductal region and the stage of the estrous cycle (Grippio *et al.* 1995, Way *et al.* 1997). Taitzoglou *et al.* (2007) reported changes in exposed saccharide residues of bovine sperm during capacitation in the presence of oviductal fluid and they suggested that these modifications could influence sperm–zona binding, zona penetration, and interaction with the oolemma.

It has been reported that the exposure of hamster oocytes to the oviductal environment increased sperm–zona binding, zona penetration and fertilization (Boatman *et al.* 1994). In accordance with the previous results, incubation of hamster oocytes in the presence of oviductal fluid improved both sperm–zona binding, zona penetration and fertilization (Kito & Bavister 1996).

A study conducted on dogs reported that incubation in the presence of oviductal fluid increased the sperm–ZP binding (Kawakami *et al.* 1998). Another study revealed that co-culture of equine gametes with homologous and heterologous (porcine) oviductal cells increased equine IVF rates (Mugnier *et al.* 2009). Coy *et al.* (2010) reported that exposure of boar spermatozoa to pre-ovulatory porcine oviductal fluid increased zona binding and polyspermy during IVF.

In the case of human species, co-incubation with oviductal cells or their conditioned media was found to reduce the human sperm binding to ZP (Morales *et al.* 1996, Yao *et al.* 1999).

It has been shown that exposure of ZP to oviductal fluid increased ZP resistance to proteolysis. This effect was associated with reduced polyspermy by decreasing sperm–ZP binding in porcine and bovine species (Coy *et al.* 2008). The increased ZP resistance to proteolysis by oviductal fluid seems not to be species specific, at least among ruminants and rabbits (Katska *et al.* 1989, 1999). Supporting this suggestion partially, the presence of human oviductal fluid increased ZP resistance to proteolysis in rabbit, sheep, pig, and bovine species, but not in human species, suggesting that this effect could depend on the species (Mondéjar *et al.* 2013a).

Involvement of oviductal proteins in gamete interaction

Experimental evidence supports that proteins play a central role in modulating gamete interaction. Once the

spermatozoon detects the oocyte, it must cross two barriers, the cumulus cells and the ZP, before it could contact the female gamete plasma membrane (Ikawa *et al.* 2010, Coy *et al.* 2012a; Fig. 2). The ZP is a glycoprotein cover that surrounds the oocyte in mammals and would be responsible for the species specificity of gamete interaction (Wassarman *et al.* 1999).

It has been suggested that species-specific zona adhesion is not mediated by a single receptor. Instead, sperm–zona binding would involve a multiplicity of receptor–ligand interactions and would require the coordinated action of different proteins (Nixon *et al.* 2005, Gahlay *et al.* 2010, Clark *et al.* 2011, Avella *et al.* 2013). The initial mouse sperm–zona binding could also involve components of the ZP acquired in the oviduct after ovulation (Lyng & Shur 2009).

Experimental evidence suggested that there is a dynamic aggregation of zona proteins believed to be important in sperm–ZP recognition to the regions of sperm that mediate this binding event (Gahlay *et al.* 2010). In addition, a recent study has demonstrated that mannose, fucose, and β -N-acetylglucosamine were terminal carbohydrates on the mouse oocyte ZP involved in cross-linking or aggregation with receptors on the sperm membrane (Wu & Sampson 2014). It has been suggested that mouse sperm receptors for zona proteins interact with both the glycans and the protein backbone of the ZP (Clark *et al.* 2011).

It has been proposed that the interaction between integrin-like proteins on the oolemma and disintegrins of transmembrane proteins that contains a disintegrin and metalloprotease domain (ADAM), found on the sperm membrane would be involved in the gamete fusion process (Ikawa *et al.* 2010, Inoue *et al.* 2011). However, this idea is mainly supported by results obtained from studies on the fusion of gamete membranes carried out in mouse models.

Different experimental approaches have demonstrated that oviductal proteins could interact with gametes and affect gamete interaction (Fig. 2). A previous study has reported that six proteins from the bovine oviductal fluid were attached to the homologous ZP (Staros & Killian 1998). In a study carried out in our laboratory, the presence of proteins from the conditioned media of human oviductal tissue culture resulted in a dose-dependent inhibition of the sperm–ZP binding and decreased the detection of sperm D-mannose-binding sites, which are associated with gamete interaction (Munuce *et al.* 2009).

Some of the oviductal proteins suggested to be involved in modulating gamete interaction have been identified and their reported effects are described below.

Oviductins

These glycoproteins secreted by the oviductal epithelium were shown to interact with the oocyte ZP in different species. O'Day-Bowman *et al.* (1996) reported that

incubation of human or baboon spermatozoa in the presence of the oviductin increased hemizona binding and penetration to hamster oocytes. The presence of oviductin also increased the *in vitro* sperm fertilizing ability in bovine species (King *et al.* 1994, Martus *et al.* 1998). Oviductins were suggested to participate in the initial sperm–zona adhesion in mice and were found to be associated with both the ZP and the perivitelline space of mouse oocytes (Ensslin *et al.* 2007, Lyng & Shur 2009). Despite the results mentioned above, a study using *Ovgp1*^{−/−} mice indicated that fertility of *Ovgp1*^{−/−} females was within the normal limits, suggesting that, at least in mice, the protein was not essential for the process of *in vivo* fertilization (Araki *et al.* 2003).

An early study demonstrated that hamster oviductin was bound to the ZP during transit of the oocyte in the oviduct (St-Jacques *et al.* 1992). However, the exposure to oviductin was shown to decrease the sperm–zona interaction and inhibit IVF of cumulus-free oocytes in hamsters (Kimura *et al.* 1994, Saccary *et al.* 2013). It has been reported that exposure to porcine oviductin before and during IVF decreased the incidence of polyspermy in pig oocytes and reduced the number of bound sperm (Kouba *et al.* 2000, McCauley *et al.* 2003). Oviductin and heparin-like glycosaminoglycans have been implicated in the pre-fertilization ZP hardening in cows and pigs, which could affect sperm binding and would contribute to prevent polyspermy (Coy *et al.* 2008). In addition to oviductin, proteins from the heat shock protein family and the protein disulfide isomerase A4 appear to be involved in ZP hardening in bovine species (Mondéjar *et al.* 2013b).

Osteopontin

A study on bovine species demonstrated the binding of osteopontin (which is secreted by the oviductal epithelium) to the ZP (Gonçalves *et al.* 2008). It has also been reported that the pre-incubation of spermatozoa or oocytes with oviductal fluid pre-treated with antibodies against osteopontin reduced sperm–ZP binding and IVF (Gonçalves *et al.* 2007, 2008). Another study has demonstrated that the exposure to osteopontin during IVF decreased polyspermy in pigs (Hao *et al.* 2006, 2008). In addition, it has also been demonstrated that exposure to osteopontin during IVF also increased the fertilization efficiency in pigs and slightly increased the IVF rates in equine species (Hao *et al.* 2006, 2008, Mugnier *et al.* 2009).

Glycodelins

Exposure to glycodelin A has been shown to decrease human gamete interaction *in vitro* (Oehninger *et al.* 1995). This effect would result from blocking the binding of the sperm fucosyltransferase 5 (suggested to be the sperm glycodelin A receptor) to the ZP (Chiu *et al.* 2007a). The presence of glycodelin F has also been reported to reduce gamete interaction (Chiu *et al.* 2003).

Glycodelin C would remove the previously attached glycodelins A and F from spermatozoa, enhancing the zona-binding capacity of sperm passing through the cumulus oophorus (Chiu *et al.* 2007b).

HSPA5

This protein also known as GRP8 was first isolated from the plasma membrane of oviduct epithelial cells and its expression in these cells changed throughout the estrous cycle and would be under hormonal control (Bauersachs *et al.* 2004, Boilard *et al.* 2004). The presence of this protein was detected in human oviductal fluid and the conditioned medium from tubal tissue cultures (Marín-Briggiler *et al.* 2010). Recombinant GRP78 was found to bind to spermatozoa and was able to decrease sperm–zona interaction in a dose-dependent manner (Marín-Briggiler *et al.* 2010).

Atrial natriuretic peptide

It has been shown that pre-incubation of spermatozoa with this peptide increased the oocyte penetration rate and decreased the polyspermy rate and the average number of sperm per penetrated oocyte in pigs (Zhang *et al.* 2006).

S100A11

The expression of this protein, which belongs to the S100 family of proteins, was detected in the oviductal epithelium, mainly in the ampullary region of the mouse oviduct and at the estrous stage of the estrous cycle (Hanaue *et al.* 2011). S100A11 was also detected bound to the plasma membrane of cumulus cells surrounding the oocytes. Pre-treatment of the cumulus cell–oocyte complex with recombinant S100A11 significantly reduced the efficiency of IVF in mice. The authors suggested that the effect could be mediated through the binding of S100A11 to the plasma membrane of the cumulus cells (Miwa *et al.* 2010, Hanaue *et al.* 2011).

Deleted in malignant brain tumors 1 (DMBT1)

This protein was shown to be expressed by the oviductal epithelium and was localized to the ZP and the cytoplasm of oocytes in equine and porcine species (Ambruosi *et al.* 2013). Pre-incubation of oocytes with recombinant deleted in malignant brain tumors 1 (DMBT1) increased the monospermic IVF rate in pigs (Ambruosi *et al.* 2013). The reported effect was also observed when oviductal fluids were used instead of the recombinant protein, but there was no effect when an antibody against DMBT1 was previously added to the oviductal fluid (Ambruosi *et al.* 2013).

Glycosidase enzymes

These enzymes have been detected in hamster, porcine, and bovine oviductal secretion (Tulsiani *et al.* 1996,

Carrasco *et al.* 2008a, b). It has been suggested that these enzymes could affect the protein and carbohydrate distribution on the sperm and the ZP surface and could modulate sperm–oocyte binding and gamete–oviductal epithelium interaction, both suggested to be carbohydrate-mediated events occurring in the presence of the oviductal fluid (Carrasco *et al.* 2008a, b).

Proteases

Plasminogen activators were detected in porcine and bovine oviductal flushing and their relative concentrations were found to change during the ovulatory cycle (Roldán-Olarte *et al.* 2005). Plasminogen is present in the oviductal fluid through the estrous cycle (Mondéjar *et al.* 2012). Plasminogen activators and their main substrate, plasminogen, were also found in the cumulus cell extracellular matrix and oocyte ZP (Roldán-Olarte *et al.* 2005, Mondéjar *et al.* 2012). A recent study has demonstrated that the presence of plasminogen in the IVF medium decreased sperm penetration of oocytes in porcine and bovine species (Mondéjar *et al.* 2012). It has been suggested that sperm binding to oocytes triggers the releasing of plasminogen activators from the oocyte and the generated plasmin causes sperm detachment from the ZP (Coy *et al.* 2012b). Supporting this idea, it has been reported that, after fertilization, plasminogen and plasminogen activator immunolabeling decreases in the oocyte, suggesting its conversion into plasmin (Mondéjar *et al.* 2012).

Sperm adhesion molecule 1

This hyaluronidase, together with other enzymes, has been implicated in assisting the sperm penetration through the cumulus cell layer surrounding the ZP (Lin *et al.* 1994). It has also been involved in the binding of acrosome-reacted sperm to the ZP (Myles & Primakoff 1997, Redgrove *et al.* 2013).

Lactoferrin

In a recent study, human lactoferrin was shown to bind to ZP. In addition, the presence of the protein caused a dose-dependent decrease in the human sperm–zona interaction (Zumoffen *et al.* 2013). Lactoferrin was also shown to reduce the availability of sperm D-mannose receptors (C Zumoffen and S Ghersevich, unpublished observations). The latter effect could partially explain the inhibition of sperm–zona binding in the presence of the protein.

Involvement of oviductal secretion in embryo development

Some studies have suggested that co-culture of gametes or embryos with oviduct epithelial cells benefits IVF rate and embryo development respectively (Liu *et al.* 1995, Wiemer *et al.* 1995, Romar *et al.* 2001, Kattal *et al.* 2008,

Mugnier *et al.* 2009). However, these results were not supported by studies on other mammalian species (Izquierdo *et al.* 2002, Kidson *et al.* 2003, Shirazi & Motaghi 2013). The difference in the mentioned results may reflect the fact that every study was performed under very different conditions. However, potential species-specific effects of co-culture with oviductal cells on embryo development could not be ruled out, but they remain to be demonstrated yet.

It has been proposed that co-culture with oviduct epithelial cells would reduce the undesirable factors in the culture medium and this would benefit embryo development. It is possible that epithelial cells, through their metabolic capacity, reduce the atmospheric oxygen pressure and the levels of substances such as glucose, which act as inhibitors of embryonic development *in vitro*. Another possible mechanism would involve the production of embryotropic factors, such as certain growth factors, whose presence in the oviduct has been clearly documented. Growth factors in the oviduct were suggested to be involved in preimplantation embryo development (Dalton *et al.* 1994, Einspanier *et al.* 1999, Pushpakumara *et al.* 2002, Wijayagunawardane *et al.* 2005, Itoh *et al.* 2006, Sun *et al.* 2006, Kawamura *et al.* 2007, Weng *et al.* 2009, Swangchan-Uthai *et al.* 2011).

It has been reported that the condition of the oviduct where bovine embryos were placed until reaching the blastocyst stage influenced their gene expression patterns, especially for those genes that regulate metabolic activity (Gad *et al.* 2011). Thus, factors from the oviductal secretion could affect the embryo gene expression.

In addition, Lee *et al.* (2005, 2006) have analyzed the interaction between embryos and oviduct in mice and their results suggested that the presence of the embryo could affect the protein expression of the mouse oviductal epithelium, as shown for the phospholipid transfer protein (PLTP) and the demilune cell and parotid protein.

Phospholipid transfer protein

In the presence of an embryo, the mouse oviductal epithelium secretes PLTP, which showed a higher expression in the embryo-containing oviduct than in the control oviduct (Lee *et al.* 2005). As *Pltp* mRNA increased in the oviductal epithelia of pregnant mice, the authors suggested that it could be involved in *in vivo* preimplantation embryo development (Lee *et al.* 2005).

Demilune cell and parotid protein

This protein was also highly expressed in mouse oviductal lumen in the presence of embryos (Lee *et al.* 2006). Demilune cell and parotid protein was shown to stimulate the growth of preimplantation embryos, suggesting that it may participate in embryo–maternal dialog (Lee *et al.* 2006). Other proteins present in the oviductal secretion suggested to be involved in modulating embryo development are mentioned below.

Oviductins

They were localized to the perivitelline space and the membrane of embryos from different species before the implantation. Their densely glycosylated mucin-type domains would act as a protective shield around the oocyte and the early embryo and would be important for early stages of embryo development (Malette *et al.* 1995, Boatman 1997, Velasquez *et al.* 2001, Buhi 2002, Wolf *et al.* 2003, Gonçalves *et al.* 2008). Porcine oviductin increased post-cleavage embryo development to blastocyst (Kouba *et al.* 2000, McCauley *et al.* 2003). In a study on goats, Pradeep *et al.* also reported that oviductin increased the cleavage rate, and morula and blastocyst formation (Pradeep *et al.* 2011).

Osteopontin

A study on bovine species reported that pre-incubation of oocytes with oviductal fluid pre-treated with antibodies against osteopontin reduced the *in vitro* embryo development when compared with the oviductal fluid alone (Gonçalves *et al.* 2008). Supporting the mentioned results, the presence of osteopontin improved the efficiency of *in vitro* embryo production in bovine species (Monaco *et al.* 2009). In addition, the exposure to osteopontin during IVF has also been shown to improve *in vitro* development of porcine embryos (Hao *et al.* 2006, 2008).

Human oviduct-derived embryotropic factor 3

This factor contains complement protein 3 (C3) and its derivatives C3b and inactivated complement 3b (iC3b) (Lee *et al.* 2004). C3 is produced and secreted by human and mouse oviductal cells. Both derivatives, but not C3, were embryotropic, while iC3b was most efficient in enhancing the mouse blastocyst development, with larger size and higher hatching rate (Lee *et al.* 2004). Oviductal cells possess C3 convertase activity converting C3 to C3b (Tse *et al.* 2008). It has been shown that the mouse preimplantation embryos may cooperate with oviductal cells to produce embryotropic iC3b (Tse *et al.* 2008). The levels of C3 and iC3b in mouse oviductal fluid were highest on day 3 of pregnancy, when they could enhance the blastocyst development and result in larger size and higher embryo hatching rate *in vitro* (Lee *et al.* 2009). Based on these data, it has been suggested that the oviduct produced C3/C3b, which could be converted into iC3b in the presence of the embryo stimulating its development.

Concluding remarks

Until a few decades ago, the oviduct was considered as a simple passive conduit that provided an optimal microenvironment in terms of temperature, pH, osmotic pressure, nutrients, and oxygen pressure, enabling both

the fertilization process and the early stages of embryo development (Pauerstein & Eddy 1979). The accumulated experimental evidence reviewed in the present work supports that the oviduct is actively involved in the reproductive process, considering that its secretion contains molecules capable of modulating gamete functions and interaction. It would also contribute to regulate the early stages of embryo development.

Based on the reported data in different mammalian species, it has been considered that spermatozoa in the oviduct interact with factors that would help select subpopulations of male gametes that remain viable while developing an optimal fertilizing ability. The oviductal environment would also contribute to decrease the number of sperm that could interact with the oocytes.

Numerous studies have identified specific molecules, most of which are proteins, in the oviductal environment that could be involved in modulating different stages of the reproductive process. The expression of certain protein components of the oviductal secretion would be subject to cyclic changes of sex steroids. In addition, the presence of gametes or embryos could affect the protein expression from oviductal epithelial cells. It could be thought that the resulting effect of the oviductal secretion, either inhibitory or stimulatory, on the reproductive process would result from the dynamic cooperative action of multiple factors present in the oviduct at different stages of the ovulatory cycle (Fig. 4). The combined action of these factors, either inducers

or repressors, could contribute to the regulation of the complex mechanism of reproduction in the oviduct. The balance between the stimulatory and inhibitory effects would result from the regulation of the expression of these oviductal factors, which could be dependent on the ovulatory cycle and the oviductal region, induced by the presence of gametes or embryos, constitutive, or even influenced by individual characteristics or by certain disorders. Thus, the deficiency of a given molecule might not impair fertility capacity because its action could be compensated by another factor with similar functions. However, any alteration in this balance could affect some stages of the reproductive process and could perhaps impair fertility.

Therefore, the complexity of the reproductive process warrants a continuous research effort to unveil the mechanisms and oviductal factors behind its regulation.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the review.

Funding

This review did not receive any specific grant from any funding agency in the public, commercial, or not-for-profit sector.

Acknowledgements

The authors thank M Robson, M de Sanctis, and G Raimundo for their assistance in manuscript proofreading.

References

- Abe H, Sendai Y, Satoh T & Hoshi H 1995 Bovine oviduct-specific glycoprotein: a potent factor for maintenance of viability and motility of bovine spermatozoa *in vitro*. *Molecular Reproduction and Development* **42** 226–232. (doi:10.1002/mrd.1080420212)
- Abou-haila A & Tulsiani DR 2009 Signal transduction pathways that regulate sperm capacitation and the acrosome reaction. *Archives of Biochemistry and Biophysics* **485** 72–81. (doi:10.1016/j.abb.2009.02.003)
- Ambruosi B, Accogli G, Douet C, Canepa S, Pascal G, Monget P, Moros Nicolás C, Holmskov U, Mollenhauer J, Robbe-Masselot C *et al.* 2013 Deleted in malignant brain tumor 1 is secreted in the oviduct and involved in the mechanism of fertilization in equine and porcine species. *Reproduction* **146** 119–133. (doi:10.1530/REP-13-0007)
- Araki Y, Nohara M, Yoshida-Komiya H, Kuramochi T, Ito M, Hoshi H, Shinkai Y & Sendai Y 2003 Effect of a null mutation of the oviduct-specific glycoprotein gene on mouse fertilization. *Biochemical Journal* **374** 551–557. (doi:10.1042/BJ20030466)
- Arias EB, Verhage HG & Jaffe RC 1994 Complementary deoxyribonucleic acid cloning and molecular characterization of an estrogen-dependent human oviductal glycoprotein. *Biology of Reproduction* **51** 685–694. (doi:10.1095/biolreprod51.4.685)
- Avella MA, Xiong B & Dean J 2013 The molecular basis of gamete recognition in mice and humans. *Molecular Human Reproduction* **19** 279–289. (doi:10.1093/molehr/gat004)

Figure 4 Drawing showing the suggested actions of the oviductal secretion in different stages of the reproductive process. The effect of the oviductal secretion would result from the combined action of several factors, either stimulatory or inhibitory. The balance between the stimulatory and inhibitory effects would result from the regulation of the expression of these oviductal factors, which could be dependent on the ovulatory cycle and the oviductal region, induced by the presence of gametes, constitutive, or even associated with a given disorder. Among oviductal proteins from different species with potential inhibitory actions, the following ones could be mentioned: glycodeilins A and F, GRP78, S100 A11, and lactoferrin. On the other hand, proteins such as oviductins, osteopontin, atrial natriuretic peptide, glycodeilin C, and sperm adhesion molecule 1 were reported to have some stimulatory effects *in vitro*. The symbols '+' and '-' indicate stimulatory or inhibitory effects respectively.

- Avilés M, Gutiérrez-Adán A & Coy P 2010 Oviductal secretions: will they be key factors for the future ARTs? *Molecular Human Reproduction* **16** 896–906. (doi:10.1093/molehr/gaq056)
- Bauersachs S, Rehfeld S, Ulbrich SE, Mallok S, Prelle K, Wenigerkind H, Einspanier R, Blum H & Wolf E 2004 Monitoring gene expression changes in bovine oviduct epithelial cells during the oestrous cycle. *Journal of Molecular Endocrinology* **32** 449–466. (doi:10.1677/jme.0.0320449)
- Bergqvist AS & Rodríguez-Martínez H 2006 Sulphated glycosaminoglycans (S-GAGs) and syndecans in the bovine oviduct. *Animal Reproduction Science* **93** 46–60. (doi:10.1016/j.anireprosci.2005.06.029)
- Boatman DE 1997 Responses of gametes to the oviductal environment. *Human Reproduction* **12** 133–149.
- Boatman DE, Felson SE & Kimura J 1994 Changes in morphology, sperm penetration and fertilization of ovulated hamster eggs induced by oviductal exposure. *Human Reproduction* **9** 519–526.
- Boilard M, Reyes-Moreno C, Lachance C, Massicotte L, Bailey JL, Sirard MA & Leclerc P 2004 Localization of the chaperone proteins GRP78 and HSP60 on the luminal surface of bovine oviduct epithelial cells and their association with spermatozoa. *Biology of Reproduction* **71** 1879–1889. (doi:10.1095/biolreprod.103.026849)
- Brenner RM, West NB & McClellan MC 1990 Estrogen and progesterone receptors in the reproductive tract of male and female primates. *Biology of Reproduction* **42** 11–19. (doi:10.1095/biolreprod42.1.11)
- Buhi WC 2002 Characterization and biological roles of oviduct-specific, oestrogen-dependent glycoprotein. *Reproduction* **123** 355–362. (doi:10.1530/rep.0.1230355)
- Buhi WC, Alvarez IM, Choi I, Cleaver BD & Simmen FA 1996 Molecular cloning and characterization of an estrogen-dependent porcine oviductal secretory glycoprotein. *Biology of Reproduction* **55** 1305–1314. (doi:10.1095/biolreprod55.6.1305)
- Buhi WC, Alvarez IM & Koube AJ 2000 Secreted proteins of the oviduct. *Cells, Tissues, Organs* **166** 165–179. (doi:10.1159/000016731)
- Carrasco LC, Romar R, Aviles M, Gadea J & Coy P 2008a Glycosidase determination in bovine oviductal fluid at the follicular and luteal phases of the oestrous cycle. *Reproduction, Fertility, and Development* **20** 808–817. (doi:10.1071/RD08113)
- Carrasco LC, Romar R, Aviles M, Gadea J & Coy P 2008b Determination of glycosidase activity in porcine oviductal fluid at different phases of the estrous cycle. *Reproduction* **136** 833–842. (doi:10.1530/REP-08-0221)
- Cherr GN, Yudin AI, Li MW, Vines CA & Overstreet JW 1999 Hyaluronic acid and the cumulus extracellular matrix induce increases in intracellular calcium in macaque sperm via the plasma membrane protein PH-20. *Zygote* **7** 211–222. (doi:10.1017/S0967199499000593)
- Chiu PC, Koistinen R, Koistinen H, Seppala M, Lee KF & Yeung WS 2003 Zona-binding inhibitory factor-1 from human follicular fluid is an isoform of glycodelin. *Biology of Reproduction* **69** 365–372. (doi:10.1095/biolreprod.102.012658)
- Chiu PC, Chung MK, Koistinen R, Koistinen H, Seppala M, Ho PC, Ng EH, Lee KF & Yeung WS 2007a Glycodelin-A interacts with fucosyltransferase on human sperm plasma membrane to inhibit spermatozoa–zona pellucida binding. *Journal of Cell Science* **120** 33–44. (doi:10.1242/jcs.03258)
- Chiu PC, Chung MK, Koistinen R, Koistinen H, Seppala M, Ho PC, Ng EH, Lee KF & Yeung WS 2007b Cumulus oophorus-associated glycodelin-C displaces sperm-bound glycodelin-A and -F and stimulates spermatozoa–zona pellucida binding. *Journal of Biological Chemistry* **282** 5378–5388. (doi:10.1074/jbc.M607482200)
- Clark GF 2011 Molecular models for mouse sperm–oocyte binding. *Glycobiology* **21** 3–5. (doi:10.1093/glycob/cwq159)
- Coy P, Canovas S, Mondéjar I, Saavedra MD, Romar R, Grullón L, Matas C & Aviles M 2008 Oviduct-specific glycoprotein and heparin modulate sperm–zona pellucida interaction during fertilization and contribute to the control of polyspermy. *PNAS* **105** 15809–15814. (doi:10.1073/pnas.0804422105)
- Coy P, Lloyd R, Romar R, Satake N, Matas C, Gadea J & Holt WV 2010 Effects of porcine pre-ovulatory oviductal fluid on boar sperm function. *Theriogenology* **74** 632–642. (doi:10.1016/j.theriogenology.2010.03.005)
- Coy P, García-Vázquez FA, Visconti PE & Avilés M 2012a Roles of the oviduct in mammalian fertilization. *Reproduction* **144** 649–660. (doi:10.1530/REP-12-0279)
- Coy P, Jiménez-Movilla M, García-Vázquez FA, Mondéjar I, Grullón L & Romar R 2012b Oocytes use the plasminogen-plasmin system to remove supernumerary spermatozoa. *Human Reproduction* **27** 1985–1993. (doi:10.1093/humrep/des146)
- Croxatto HB 2002 Physiology of gamete and embryo transport through the Fallopian tube. *Reproductive Biomedicine Online* **4** 160–169. (doi:10.1016/S1472-6483(10)61935-9)
- Dalton T, Kover K, Dey SK & Andrews GK 1994 Analysis of the expression of growth factor, interleukin-1, and lactoferrin genes and the distribution of inflammatory leukocytes in the preimplantation mouse oviduct. *Biology of Reproduction* **51** 597–606. (doi:10.1095/biolreprod51.4.597)
- Donnelly KM, Fazleabas AT, Verhage HG, Mavrogianis PA & Jaffe RC 1991 Cloning of a recombinant complementary DNA to a baboon (*Papio anubis*) estradiol-dependent oviduct-specific glycoprotein. *Molecular Endocrinology* **5** 356–364. (doi:10.1210/mend-5-3-356)
- Dutta B, Ain R, Seshagiri PB & Karande AK 2001 Differential influence of recombinant non-glycosylated and glycosylated glycodelin on human sperm function: comparative studies with hamster spermatozoa. *Reproduction, Fertility, and Development* **13** 111–118. (doi:10.1071/RD00119)
- Einspanier R, Gabler C, Bieser B, Einspanier A, Berisha B, Kosmann M, Wollenhaupt K & Schams D 1999 Growth factors and extracellular matrix proteins in interactions of cumulus–oocyte complex, spermatozoa and oviduct. *Journal of Reproduction and Fertility Supplement* **54** 359–365.
- Ensslin MA, Lyng R, Raymond A, Copland S & Shur BD 2007 Novel gamete receptors that facilitate sperm adhesion to the egg coat. *Journal of Reproduction & Fertility Supplement* **63** 367–383.
- Gabler C, Chapman DA & Killian GJ 2003 Expression and localization of osteopontin and integrins in the bovine oviduct during the estrous cycle. *Reproduction* **126** 721–729. (doi:10.1530/rep.0.1260721)
- Gad A, Besenfelder U, Rings F, Ghanem N, Salilew-Wondim D, Hossain MM, Tesfaye D, Lonergan P, Becker A, Cinar U *et al.* 2011 Effect of reproductive tract environment following controlled ovarian hyperstimulation treatment on embryo development and global transcriptome profile of blastocysts: implications for animal breeding and human assisted reproduction. *Human Reproduction* **26** 1693–1707. (doi:10.1093/humrep/der110)
- Gahlay G, Gauthier L, Baibakov B, Epifano O & Dean J 2010 Gamete recognition in mice depends on the cleavage status of an egg's zona pellucida protein. *Science* **329** 216–219. (doi:10.1126/science.1188178)
- Georgiou S, Sostaric E, Wong CH, Snijders APL, Wright PC & Moore HD 2005 Gametes alter the oviductal secretory proteome. *Molecular & Cellular Proteomics* **4** 1785–1796. (doi:10.1074/mcp.M500119-MCP200)
- Georgiou AS, Snijders AP, Sostaric E, Aflatoonian R, Vazquez JL, Vazquez JM, Roca J, Martinez EA, Wright PC & Fazeli A 2007 Modulation of the oviductal environment by gametes. *Journal of Proteome Research* **6** 4656–4666. (doi:10.1021/pr070349m)
- Gervasi MG, Rapanelli M, Ribeiro ML, Farina M, Billi S, Franchi AM & Perez-Martinez S 2009 The endocannabinoid system in bull sperm and bovine oviductal epithelium: role of anandamide in sperm–oviduct interaction. *Reproduction* **137** 403–414. (doi:10.1530/REP-08-0204)
- Gervasi MG, Marczylo TH, Lam PM, Rana S, Franchi AM, Konje JC & Perez-Martinez S 2013 Anandamide levels fluctuate in the bovine oviduct during the oestrous cycle. *PLoS ONE* **8** e72521. (doi:10.1371/journal.pone.0072521)
- Gonçalves RF, Wolinetz CD & Killian GJ 2007 Influence of arginine–glycine–aspartic acid (RGD), integrins (α V and α 5) and osteopontin on bovine sperm–egg binding, and fertilization *in vitro*. *Theriogenology* **67** 468–474. (doi:10.1016/j.theriogenology.2006.08.013)
- Gonçalves RF, Staros AL & Killian GJ 2008 Oviductal fluid proteins associated with the bovine zona pellucida and the effect on *in vitro* sperm–egg binding, fertilization and embryo development. *Reproduction in Domestic Animals* **43** 720–729. (doi:10.1111/j.1439-0531.2007.00978.x)
- Griffiths GS, Miller KA, Galileo DS & Martin-DeLeon PA 2008 Murine SPAM1 is secreted by the estrous uterus and oviduct in a form that can bind to sperm during capacitation: acquisition enhances hyaluronic acid-binding ability and cumulus dispersal efficiency. *Reproduction* **135** 293–301. (doi:10.1530/REP-07-0340)

- Grippe AA, Way AL & Killian GJ 1995 Effect of bovine ampullary and isthmus oviductal fluid on motility, acrosome reaction and fertility of bull spermatozoa. *Journal of Reproduction & Fertility* **105** 57–64. (doi:10.1530/jrf.0.1050057)
- Gualtieri R, Boni R, Tosti E, Zagami M & Talevi R 2005 Intracellular calcium and protein tyrosine phosphorylation during the release of bovine sperm adhering to the fallopian tube epithelium *in vitro*. *Reproduction* **129** 51–60. (doi:10.1530/rep.1.00374)
- Gualtieri R, Mollo V, Barbato V & Talevi R 2010 Ability of sulfated glycoconjugates and disulfide-reductants to release bovine epididymal sperm bound to the oviductal epithelium *in vitro*. *Theriogenology* **73** 1037–1043. (doi:10.1016/j.theriogenology.2009.11.030)
- Hanaue M, Miwa N, Uebi T, Fukuda Y, Katagiri Y & Takamatsu K 2011 Characterization of S100A11, a suppressive factor of fertilization, in the mouse female reproductive tract. *Molecular Reproduction and Development* **78** 91–103. (doi:10.1002/mrd.21273)
- Hao Y, Mathialagan N, Walters E, Mao J, Lai L, Becker D, Li W, Critser J & Prather RS 2006 Osteopontin reduces polyspermy during *in vitro* fertilization of porcine oocytes. *Biology of Reproduction* **75** 726–733. (doi:10.1095/biolreprod.106.052589)
- Hao Y, Murphy CN, Spate L, Wax D, Zhong Z, Samuel M, Mathialagan N, Schatten H & Prather RS 2008 Osteopontin improves *in vitro* development of porcine embryos and decreases apoptosis. *Molecular Reproduction and Development* **75** 291–298. (doi:10.1002/mrd.20794)
- Hess AP, Nayak NR & Giudice LC 2006 Oviduct and endometrium: cyclic changes in the primate oviduct and endometrium. In *The Physiology of Reproduction*, 3 edn, pp 337–382. Eds E Knobil & JD Neill. New York, NY: Raven Press.
- Hugentobler S, Sreenan J, Humpherson P, Leese H, Diskin M & Morris D 2010 Effects of changes in the concentration of systemic progesterone on ions, amino acids and energy substrates in cattle oviduct and uterine fluid and blood. *Reproduction, Fertility, and Development* **22** 684–694. (doi:10.1071/RD09129)
- Huhtala ML, Seppälä M, Närvänen A, Palomäki P, Julkunen M & Bohn H 1987 Amino acid sequence homology between human placental protein 14 and beta-lactoglobulins from various species. *Endocrinology* **120** 2620–2622. (doi:10.1210/endo-120-6-2620)
- Ignatz GG, Cho MY & Suarez SS 2007 Annexins are candidate oviductal receptors for bovine sperm surface proteins and thus may serve to hold bovine sperm in the oviductal reservoir. *Biology of Reproduction* **77** 906–913. (doi:10.1095/biolreprod.107.062505)
- Ikawa M, Inoue N, Benham AM & Okabe M 2010 Fertilization: a sperm's journey to and interaction with the oocyte. *Journal of Clinical Investigation* **120** 984–994. (doi:10.1172/JCI41585)
- Inoue N, Ikawa M & Okabe M 2011 The mechanism of sperm–egg interaction and the involvement of IZUMO1 in fusion. *Asian Journal of Andrology* **13** 81–87. (doi:10.1038/aja.2010.70)
- Itoh H, Nasu K, Matsumoto H, Kawano Y, Yoshimatsu J & Narahara H 2006 Hypoxia regulates vascular endothelial growth factor and soluble fms-like tyrosine kinase-1 secretion by human oviductal epithelial cells and stromal fibroblasts. *Fertility and Sterility* **85** (Suppl 1) 1097–1102. (doi:10.1016/j.fertnstert.2005.10.023)
- Izquierdo D, Villamediana P, López-Bejar M & Paramio MT 2002 Effect of *in vitro* and *in vivo* culture on embryo development from prepubertal goat IVM–IVF oocytes. *Theriogenology* **57** 1431–1441. (doi:10.1016/S0093-691X(02)00647-7)
- Kadirvel G, Machado SA, Korneli C, Collins E, Miller P, Bess KN, Aoki K, Tiemeyer M, Bovin N & Miller DJ 2012 Porcine sperm bind to specific 6-sialylated biantennary glycans to form the oviduct reservoir. *Biology of Reproduction* **87** 147–156. (doi:10.1095/biolreprod.112.103879)
- Kan FW & Esperanzate PW 2006 Surface mapping of binding of oviductin to the plasma membrane of golden hamster spermatozoa during *in vitro* capacitation and acrosome reaction. *Molecular Reproduction and Development* **73** 756–766. (doi:10.1002/mrd.20459)
- Katska L, Kauffold P, Smorag Z, Duschinski U, Torner H & Kanitz W 1989 Influence of hardening of the zona pellucida on *in vitro* fertilization of bovine oocytes. *Theriogenology* **32** 767–777. (doi:10.1016/0093-691X(89)90465-2)
- Katska L, Kania G, Smorag Z, Wayda E & Plucienniczak G 1999 Developmental capacity of bovine IVM/IVF oocytes with experimentally induced hardening of the zona pellucida. *Reproduction in Domestic Animals* **34** 255–259. (doi:10.1111/j.1439-0531.1999.tb01248.x)
- Kattal N, Cohen J & Barmat LI 2008 Role of coculture in human *in vitro* fertilization: a meta-analysis. *Fertility and Sterility* **90** 1069–1076. (doi:10.1016/j.fertnstert.2007.07.1349)
- Kavanaugh JF, Grippe AA & Killian GJ 1992 Cannulation of the bovine ampullary and isthmus oviduct. *Journal of Investigative Surgery* **5** 11–17. (doi:10.3109/08941939209031588)
- Kawakami E, Hori T & Tsutsui T 1998 Induction of dog sperm capacitation by oviductal fluid. *Journal of Veterinary Medical Science* **60** 197–202. (doi:10.1292/jvms.60.197)
- Kawakami E, Arai T, Oishi I, Hori T & Tsutsui T 2000 Induction of dog sperm capacitation by glycosaminoglycans and glycosaminoglycan amounts of oviductal and uterine fluids in bitches. *Journal of Veterinary Medical Science* **62** 65–68. (doi:10.1292/jvms.62.65)
- Kawakami E, Kashiwagi C, Hori T & Tsutsui T 2001 Effects of canine oviduct epithelial cells on movement and capacitation of homologous spermatozoa *in vitro*. *Animal Reproduction Science* **68** 121–131. (doi:10.1016/S0378-4320(01)00135-X)
- Kawamura K, Kawamura N, Fukuda J, Kumagai J, Hsueh AJ & Tanaka T 2007 Regulation of preimplantation embryo development by brain-derived neurotrophic factor. *Developmental Biology* **311** 147–158. (doi:10.1016/j.ydbio.2007.08.026)
- Kervancioglu ME, Saridogan E, Aitken RJ & Djahanbakhch O 2000 Importance of sperm-to-epithelial cell contact for capacitation of human spermatozoa in fallopian tube epithelial cell cocultures. *Fertility and Sterility* **74** 780–784. (doi:10.1016/S0015-0282(00)01514-4)
- Kidson A, Schoevers E, Langendijk P, Verheijden J, Colenbrander B & Bevers M 2003 The effect of oviductal epithelial cell co-culture during *in vitro* maturation on sow oocyte morphology, fertilization and embryo development. *Theriogenology* **59** 1889–1903. (doi:10.1016/S0093-691X(02)01291-8)
- Killian G 2011 Physiology and endocrinology symposium: evidence that oviduct secretions influence sperm function: a retrospective view for livestock. *Journal of Animal Science* **89** 1315–1322. (doi:10.2527/jas.2010-3349)
- Kimura H, Matsuda J, Ogura A, Asano T & Naiki M 1994 Affinity binding of hamster oviductin to spermatozoa and its influence on *in vitro* fertilization. *Molecular Reproduction and Development* **39** 322–327. (doi:10.1002/mrd.1080390309)
- King RS & Killian GJ 1994 Purification of bovine estrus-associated protein and localization of binding on sperm. *Biology of Reproduction* **51** 34–42. (doi:10.1095/biolreprod51.1.34)
- King RS, Anderson SH & Killian GJ 1994 Effect of bovine oviductal estrus-associated protein on the ability of sperm to capacitate and fertilize oocytes. *Journal of Andrology* **15** 468–478. (doi:10.1002/j.1939-4640)
- Kito S & Bavister BD 1996 Kinetics of sperm penetration and fertilization *in vitro* in hamster follicular and oviductal ova. *Journal of Experimental Zoology* **274** 373–383. (doi:10.1002/(SICI)1097-010X(19960415)274:6<373::AID-JEZ6>3.0.CO;2-L)
- Kölle S, Dubielzig S, Reese S, Wehrend A, König P & Kummer W 2009 Ciliary transport, gamete interaction, and effects of the early embryo in the oviduct: ex vivo analyses using a new digital videomicroscopic system in the cow. *Biology of Reproduction* **81** 267–274. (doi:10.1095/biolreprod.108.073874)
- Kouba AJ, Abeydeera LR, Alvarez IM, Day BN & Buhi WC 2000 Effects of the porcine oviduct-specific glycoprotein on fertilization, polyspermy, and embryonic development *in vitro*. *Biology of Reproduction* **63** 242–250. (doi:10.1095/biolreprod63.1.242)
- Kumaresan A, Johannisson A, Humblot P & Bergqvist AS 2012 Oviductal fluid modulates the dynamics of tyrosine phosphorylation in cryopreserved boar spermatozoa during capacitation. *Molecular Reproduction and Development* **79** 525–540. (doi:10.1002/mrd.22058)
- Lee YL, Lee KF, Xu JS, He QY, Chiu JF, Lee WM, Luk JM & Yeung WS 2004 The embryotrophic activity of oviductal cell-derived complement C3b and iC3b, a novel function of complement protein in reproduction. *Journal of Biological Chemistry* **279** 12763–12768. (doi:10.1074/jbc.M311160200)
- Lee KF, Kwok KL, Chung MK, Lee YL, Chow JF & Yeung WS 2005 Phospholipid transfer protein (PLTP) mRNA expression is stimulated by developing embryos in the oviduct. *Journal of Cellular Biochemistry* **95** 740–749. (doi:10.1002/jcb.20444)

- Lee KF, Xu JS, Lee YL & Yeung WS 2006 Demilune cell and parotid protein from murine oviductal epithelium stimulates preimplantation embryo development. *Endocrinology* **147** 79–87. (doi:10.1210/en.2005-0596)
- Lee YL, Cheong AW, Chow WN, Lee KF & Yeung WS 2009 Regulation of complement-3 protein expression in human and mouse oviducts. *Molecular Reproduction and Development* **76** 301–308. (doi:10.1002/mrd.20955)
- Leese HJ 1988 The formation and function of oviduct fluid. *Journal of Reproduction & Fertility* **82** 843–856. (doi:10.1530/jrf.0.0820843)
- Leese H, Hugentobler S, Gray S, Morris D, Sturme R, Whitear S & Sreenan J 2008 Female reproductive tract fluids: composition, mechanism of formation and potential role in the developmental origins of health and disease. *Reproduction, Fertility, and Development* **20** 1–8. (doi:10.1071/RD07153)
- Lin Y, Mahan K, Lathrop WF, Myles DG & Primakoff P 1994 A hyaluronidase activity of the sperm plasma-membrane protein Ph-20 enables sperm to penetrate the cumulus layer surrounding the egg. *Journal of Cell Biology* **125** 1157–1163. (doi:10.1083/jcb.125.5.1157)
- Lippes J, Krasner J, Alfonso LA, Dacalos ED & Lucero R 1981 Human oviductal fluid proteins. *Fertility and Sterility* **36** 623–629.
- Lishko PV, Kirichok Y, Ren D, Navarro B, Chung JJ & Clapham DE 2012 The control of male fertility by spermatozoan ion channels. *Annual Review of Physiology* **74** 453–475. (doi:10.1146/annurev-physiol-020911-153258)
- Liu LP, Chan ST, Ho PC & Yeung WS 1995 Human oviductal cells produce high molecular weight factor(s) that improves the development of mouse embryo. *Human Reproduction* **10** 2781–2786. (doi:10.1093/molehr/1.8.395)
- Lyng R & Shur BD 2009 Mouse oviduct-specific glycoprotein is an egg-associated ZP3-independent sperm-adhesion ligand. *Journal of Cell Science* **122** 3894–3906. (doi:10.1242/jcs.058776)
- Maccarrone M, Barboni B, Paradisi A, Bernabò N, Gasperi V, Pistilli MG, Fezza F, Lucidi P & Mattioli M 2005 Characterization of the endocannabinoid system in boar spermatozoa and implications for sperm capacitation and acrosome reaction. *Journal of Cell Science* **118** 4393–4404. (doi:10.1242/jcs.02536)
- Malette B, Paquette Y, Merlen Y & Bleau G 1995 Oviductins possess chitinase- and mucin-like domains: a lead in the search for the biological function of these oviduct-specific ZP-associating glycoproteins. *Molecular Reproduction and Development* **41** 384–397. (doi:10.1002/mrd.1080410315)
- Marín-Briggiler C, González-Echeverría MF, Munuce MJ, Ghersevich S, Caille AM, Corrigan V, Panayi G, Hellman U & Vazquez-Levin MH 2010 Glucose-regulated protein 78 (Grp78/BiP) is secreted by human oviduct epithelial cells and modulates sperm-zona pellucida binding. *Fertility and Sterility* **93** 1574–1584. (doi:10.1016/j.fertnstert.2008.12.132)
- Martus NS, Verhage HG, Mavrogianis PA & Thibodeaux JK 1998 Enhancement of bovine oocyte fertilization *in vitro* with a bovine oviductal specific glycoprotein. *Journal of Reproduction and Fertility* **113** 323–329. (doi:10.1530/jrf.0.1130323)
- McCauley TC, Buhi WC, Wu GM, Mao J, Caamano JN, Didion BA & Day BN 2003 Oviduct-specific glycoprotein modulates sperm-zona binding and improves efficiency of porcine fertilization *in vitro*. *Biology of Reproduction* **69** 828–834. (doi:10.1095/biolreprod.103.016444)
- Miwa N, Ogawa M, Shinmyo Y, Hiraoka Y, Takamatsu K & Kawamura S 2010 Dicalcin inhibits fertilization through its binding to a glycoprotein in the egg envelope in *Xenopus laevis*. *Journal of Biological Chemistry* **285** 15627–15636. (doi:10.1074/jbc.M109.079483)
- Monaco E, Gasparrini B, Boccia L, De Rosa A, Attanasio L, Zicarelli L & Killian G 2009 Effect of osteopontin (OPN) on *in vitro* embryo development in cattle. *Theriogenology* **71** 450–457. (doi:10.1016/j.theriogenology.2008.08.012)
- Mondéjar I, Grullón LA, García-Vázquez FA, Romar R & Coy P 2012 Fertilization outcome could be regulated by binding of oviductal plasminogen to oocytes and by releasing of plasminogen activators during interplay between gametes. *Fertility and Sterility* **97** 453–461. (doi:10.1016/j.fertnstert.2011.11.032)
- Mondéjar I, Avilés M & Coy P 2013a The human is an exception to the evolutionarily-conserved phenomenon of pre-fertilization zona pellucida resistance to proteolysis induced by oviductal fluid. *Human Reproduction* **28** 718–728. (doi:10.1093/humrep/des423)
- Mondéjar I, Martínez-Martínez I, Avilés M & Coy P 2013b Identification of potential oviductal factors responsible for zona pellucida hardening and monospermy during fertilization in mammals. *Biology of Reproduction* **89** 67–75. (doi:10.1095/biolreprod.113.111385)
- Morales P, Palma V, Salgado AM & Villalón M 1996 Sperm interaction with human oviductal cells *in vitro*. *Human Reproduction* **11** 1504–1509. (doi:10.1093/oxfordjournals.humrep.a019426)
- Mugnier S, Kervella M, Douet C, Canepa S, Pascal G, Deleuze S, Duchamp G, Monget P & Goudet G 2009 The secretions of oviduct epithelial cells increase the equine *in vitro* fertilization rate: are osteopontin, atrial natriuretic peptide A and oviductin involved? *Reproductive Biology and Endocrinology* **7** 129–145. (doi:10.1186/1477-7827-7-129)
- Munuce MJ, Serravallo A, Caille A, Zumoffen C, Botti G, Cabada M & Ghersevich S 2009 Human tubal secretion can modify the affinity of human spermatozoa for the zona pellucida. *Fertility and Sterility* **91** 407–413. (doi:10.1016/j.fertnstert.2007.11.077)
- Myles DG & Primakoff P 1997 Why did the sperm cross the cumulus? To get to the oocyte. Functions of the sperm surface proteins PH-20 and fertilin in arriving at, and fusing with, the egg. *Biology of Reproduction* **56** 320–327. (doi:10.1095/biolreprod56.2.320)
- Nixon B, Asquith KL & Aitken R 2005 The role of molecular chaperones in mouse sperm-egg interactions. *Molecular and Cellular Endocrinology* **240** 1–10. (doi:10.1016/j.mce.2005.06.004)
- O'Day-Bowman MB, Mavrogianis PA, Reuter LM, Johnson DE, Fazleabas AT & Verhage HG 1996 Association of oviduct-specific glycoproteins with human and baboon (*Papio anubis*) ovarian oocytes and enhancement of human sperm binding to human hemizonae following *in vitro* incubation. *Biology of Reproduction* **54** 60–69. (doi:10.1095/biolreprod54.1.60)
- Oehninger S, Coddington CC, Hodgen GD & Seppala M 1995 Factors affecting fertilization: endometrial placental protein 14 reduces the capacity of human spermatozoa to bind to the human zona pellucida. *Fertility and Sterility* **63** 377–383. (doi:10.1016/0020-7292(95)97553-E)
- Osycka-Salut C, Gervasi MG, Pereyra E, Cella M, Ribeiro ML, Franchi AM & Perez-Martinez S 2012 Anandamide induces sperm release from oviductal epithelia through nitric oxide pathway in bovines. *PLoS ONE* **7** e30671. (doi:10.1371/journal.pone.0030671)
- Pauerstein CJ & Eddy CA 1979 The role of the oviduct in reproduction; our knowledge and our ignorance. *Journal of Reproduction and Fertility* **55** 223–229. (doi:10.1530/jrf.0.0550223)
- Petrunkina AM, Gehlhaar R, Drommer W, Waberski D & Töpfer-Petersen E 2001 Selective sperm binding to pig oviductal epithelium *in vitro*. *Reproduction* **121** 889–896. (doi:10.1530/rep.0.1210889)
- Pradeep MA, Jagadeesh J, De AK, Kaushik JK, Malakar D, Kumar S, Dang AK, Das SK & Mohanty AK 2011 Purification, sequence characterization and effect of goat oviduct-specific glycoprotein on *in vitro* embryo development. *Theriogenology* **75** 1005–1015. (doi:10.1016/j.theriogenology.2010.11.007)
- Pushpakumara PG, Robinson RS, Demmers KJ, Mann GE, Sinclair KD, Webb R & Wathes DC 2002 Expression of the insulin-like growth factor (IGF) system in the bovine oviduct at oestrus and during early pregnancy. *Reproduction* **123** 859–868. (doi:10.1530/rep.0.1230859)
- Quintero I, Ghersevich S, Caille A, Munuce M, Daniele SM & Morisoli L 2005 Effects of human oviductal *in vitro* secretion on spermatozoa and search of sperm-oviductal proteins interactions. *International Journal of Andrology* **28** 137–143. (doi:10.1111/j.1365-2605.2005.00525.x)
- Redgrove KA, Anderson AL, McLaughlin EA, O'Bryan MK, Aitken RJ & Nixon B 2013 Investigation of the mechanisms by which the molecular chaperone HSPA2 regulates the expression of sperm surface receptors involved in human sperm-oocyte recognition. *Molecular Human Reproduction* **19** 120–135. (doi:10.1093/molehr/gas064)
- Rodríguez-Martínez H 2007 Role of the oviduct in sperm capacitation. *Theriogenology* **68** (Suppl 1) S138–S146. (doi:10.1016/j.theriogenology.2007.03.018)
- Roldán-Olarte M, Jiménez-Díaz M & Miceli DC 2005 Plasminogen detection in oocytes and plasminogen activator activities in the porcine oviduct during the estrous cycle. *Zygote* **13** 115–123. (doi:10.1017/S0967199405003175)
- Romar R, Coy P, Campos I, Gadea J, Matás C & Ruiz S 2001 Effect of co-culture of porcine sperm and oocytes with porcine oviductal epithelial cells on *in vitro* fertilization. *Animal Reproduction Science* **68** 85–98. (doi:10.1016/S0378-4320(01)00133-6)

- Rossato M, Ion Popa F, Ferigo M, Clari G & Foresta C 2005 Human sperm express cannabinoid receptor Cb1, the activation of which inhibits motility, acrosome reaction, and mitochondrial function. *Journal of Clinical Endocrinology and Metabolism* **90** 984–991. (doi:10.1210/jc.2004-1287)
- Sabeur K, Cherr GN, Yudin AI & Overstreet JW 1998 Hyaluronic acid enhances induction of the acrosome reaction of human sperm through interaction with the PH-20 protein. *Zygote* **6** 103–111. (doi:10.1017/S0967199498000021)
- Saccary L, She YM, Oko R & Kan FW 2013 Hamster oviductin regulates tyrosine phosphorylation of sperm proteins during *in vitro* capacitation. *Biology of Reproduction* **89** 38–49. (doi:10.1095/biolreprod.113.109314)
- Schuel H, Burkman LJ, Lippes J, Crickard K, Forester E, Piomelli D & Giuffrida A 2002a *N*-acylethanolamines in human reproductive fluids. *Chemistry and Physics of Lipids* **121** 211–227. (doi:10.1016/S0009-3084(02)00158-5)
- Schuel H, Burkman LJ, Lippes J, Crickard K, Mahony MC, Giuffrida A, Picone RP & Makriyannis A 2002b Evidence that anandamide-signaling regulates human sperm functions required for fertilization. *Molecular Reproduction and Development* **63** 376–387. (doi:10.1002/mrd.90021)
- Sendai Y, Komiya H, Suzuki K, Onuma T, Kikuchi M, Hoshi H & Araki Y 1995 Molecular cloning and characterization of a mouse oviduct-specific glycoprotein. *Biology of Reproduction* **53** 285–294. (doi:10.1095/biolreprod53.2.285)
- Seytanoglu A, Georgiou AS, Sostaric E, Watson PF, Holt WV & Fazeli A 2008 Oviductal cell proteome alterations during the reproductive cycle in pigs. *Journal of Proteome Research* **7** 2825–2833. (doi:10.1021/pr8000095)
- Shirazi A & Motaghi E 2013 The *in vitro* fertilization of ovine oocytes in the presence of oviductal cells and its effect on the expression of zygote arrest 1 (zZar1) and subsequent embryonic development. *Journal of reproduction & infertility* **14** 8–16.
- Souza CE, Moura AA, Monaco E & Killian GJ 2008 Binding patterns of bovine seminal plasma proteins A1/A2, 30 kDa and osteopontin on ejaculated sperm before and after incubation with isthmic and ampullary oviductal fluid. *Animal Reproduction Science* **105** 72–89. (doi:10.1016/j.anireprosci.2007.11.027)
- Staros AL & Killian GJ 1998 *In vitro* association of six oviductal fluid proteins with the bovine zona pellucida. *Journal of Reproduction and Fertility* **112** 131–137. (doi:10.1530/jrf.0.1120131)
- St-Jacques S, Malette B, Chevalier S, Roberts KD & Bleau G 1992 The zona pellucida binds the mature form of an oviductal glycoprotein (oviductin). *Journal of Experimental Zoology* **262** 97–104. (doi:10.1002/jez.1402620113)
- Suarez SS 2006 Gamete and zygote transport. In *The Physiology of Reproduction*, 3rd edn, pp 133–148. Eds E Knobil & JD Neill. New York, NY: Raven Press.
- Suarez SS 2008a Regulation of sperm storage and movement in the mammalian oviduct. *International Journal of Developmental Biology* **52** 455–462. (doi:10.1387/ijdb.072527ss)
- Suarez SS 2008b Control of hyperactivation in sperm. *Human Reproduction Update* **14** 647–657. (doi:10.1093/humupd/dmn029)
- Sun XX, Gemzell-Danielsson K, Li HZ, Ståbi B & Stavreus-Evers A 2006 Expression of heparin-binding epidermal growth factor-like growth factor and its receptors in the human fallopian tube and endometrium after treatment with mifepristone. *Fertility and Sterility* **85** 171–178. (doi:10.1016/j.fertnstert.2005.08.011)
- Swangchan-Uthai T, Walsh SW, Alexander SL, Cheng Z, Crowe MA, Evans AC & Wathes DC 2011 Comparison of mRNA for IGFs and their binding proteins in the oviduct during the peri-oestrous period between dairy heifers and lactating cows. *Reproduction* **142** 457–465. (doi:10.1530/REP-11-0154)
- Taitzoglou IA, Kokoli AN & Killian GJ 2007 Modifications of surface carbohydrates on bovine spermatozoa mediated by oviductal fluid: a flow cytometric study using lectins. *International Journal of Andrology* **30** 108–114. (doi:10.1111/j.1365-2605.2006.00717.x)
- Talevi R, Zagami M, Castaldo M & Gualtieri R 2007 Redox regulation of sperm surface thiols modulates adhesion to the fallopian tube epithelium. *Biology of Reproduction* **76** 728–735. (doi:10.1095/biolreprod.106.056028)
- Talevi R, Barbato V, De Iorio S, Mollo V, Capriglione T, Ricchiari L, Samo A & Gualtieri R 2010 Is there a role for endocannabinoids in sperm–oviduct interaction? *Reproduction* **140** 247–257. (doi:10.1530/REP-10-0095)
- Teijeiro JM, Ignatz GG & Marini PE 2009 Annexin A2 is involved in pig (*Sus scrofa*) sperm–oviduct interaction. *Molecular Reproduction and Development* **76** 334–341. (doi:10.1002/mrd.20958)
- Thomas PG, Ignatz GG, Ball BA, Brinsko SP & Currie WB 1995a Effect of coculture with stallion spermatozoa on *de novo* protein synthesis and secretion by equine oviduct epithelial cells. *American Journal of Veterinary Research* **56** 1657–1662.
- Thomas PG, Ball BA, Brinsko SP 1995b Changes associated with induced capacitation influence the interaction between equine spermatozoa and oviduct epithelial cells monolayers. In *Equine Reproduction VI: Biology of Reproduction Monograph Series I*. eds Sharp DC, Bazer FW (Society for the Study of Reproduction, Inc. Madison, WI:) pp 697–706.
- Tienthai P, Kjellén L, Pertoft H, Suzuki K & Rodriguez-Martinez H 2000 Localization and quantitation of hyaluronan and sulfated glycosaminoglycans in the tissues and intraluminal fluid of the pig oviduct. *Reproduction, Fertility, and Development* **12** 173–182. (doi:10.1071/RD00034)
- Tienthai P, Johannisson A & Rodriguez-Martinez H 2004 Sperm capacitation in the porcine oviduct. *Animal Reproduction Science* **80** 131–146. (doi:10.1016/S0378-4320(03)00134-9)
- Travis AJ & Kopf GS 2002 The role of cholesterol efflux in regulating the fertilization potential of mammalian spermatozoa. *Journal of Clinical Investigation* **110** 731–736. (doi:10.1172/JCI0216392)
- Tse PK, Lee YL, Chow WN, Luk JM, Lee KF & Yeung WS 2008 Preimplantation embryos cooperate with oviductal cells to produce embryotrophic inactivated complement-3b. *Endocrinology* **149** 1268–1276. (doi:10.1210/en.2007-1277)
- Tulsiani DR, Chayko CA, Orgebin-Crist MC & Araki Y 1996 Temporal surge of glycosyltransferase activities in the genital tract of the hamster during the estrous cycle. *Biology of Reproduction* **54** 1032–1037. (doi:10.1095/biolreprod54.5.1032)
- Velasquez LA, Maisey K, Fernandez R, Valdes D, Cardenas H, Imarai M, Delgado J, Aguilera J & Croxatto HB 2001 PAF receptor and PAF acetylhydrolase expression in the endosalpinx of the human fallopian tube: possible role of embryo-derived PAF in the control of embryo transport to the uterus. *Human Reproduction* **16** 1583–1587. (doi:10.1093/humrep/16.8.1583)
- Verhage HG, Fazleabas AT & Donnelly K 1988 The *in vitro* synthesis and release of proteins by the human oviduct. *Endocrinology* **122** 1639–1645. (doi:10.1210/endo-122-4-1639)
- Wassarman PM, Chen J, Cohen N, Litscher E, Liu C, Qi H & Williams Z 1999 Structure and function of the mammalian egg zona pellucida. *Journal of Experimental Zoology* **285** 251–258. (doi:10.1002/(SICI)1097-010X(19991015)285:3<251::AID-JEZ8>3.0.CO;2-1)
- Way AL, Schuler AM & Killian GJ 1997 Influence of bovine ampullary and isthmic oviductal fluid on sperm–egg binding and fertilization *in vitro*. *Journal of Reproduction and Fertility* **109** 95–101. (doi:10.1530/jrf.0.1090095)
- Weng Q, Shi Z, Watanabe G & Taya K 2009 Immunolocalization of NGF and its receptors trkA and p75 in the oviducts of golden hamsters during the estrous cycle. *Experimental Animals* **58** 543–546. (doi:10.1538/expanim.58.543)
- Wiemer K, Dale B, Hu Y, Steuerwald N, Maxso WS & Hoffman DI 1995 Blastocyst development in co-culture: development and morphological aspects. *Human Reproduction* **10** 3226–3232.
- Wijayagunawardane MP, Kodithuwakku SP, Yamamoto D & Miyamoto A 2005 Vascular endothelial growth factor system in the cow oviduct: a possible involvement in the regulation of oviductal motility and embryo transport. *Molecular Reproduction and Development* **72** 511–520. (doi:10.1002/mrd.20379)
- Wolf E, Arnold GJ, Bauersachs S, Beier HM, Blum H, Einspanier R, Frohlich T, Herrler A, Hiendleder S, Kolbe S *et al.* 2003 Embryo-maternal communication in bovine – strategies for deciphering a complex cross-talk. *Reproduction in Domestic Animals* **38** 276–289. (doi:10.1046/j.1439-0531.2003.00435.x)
- Wu L & Sampson NS 2014 Fucose, mannose, and β -N-acetylglucosamine glycopolymers initiate the mouse sperm acrosome reaction through convergent signaling pathways. *ACS Chemical Biology* **9** 468–475. (doi:10.1021/cb400550j)

- Yao Y, Ho P & Yeung WS** 1999 Effects of human oviductal cell coculture on various functional parameters of human spermatozoa. *Fertility and Sterility* **71** 232–239. (doi:10.1016/S0015-0282(98)00430-0)
- Yao Y, Ho P & Yeung WS** 2000 Human oviductal cells produce a factor(s) that maintains the motility of human spermatozoa *in vitro*. *Fertility and Sterility* **73** 479–486. (doi:10.1016/S0015-0282(99)00563-4)
- Yeung WS, Ng VK, Lau EY & Ho PC** 1994 Human oviductal cells and their conditioned medium maintain the motility and hyperactivation of human spermatozoa *in vitro*. *Human Reproduction* **9** 656–660.
- Yeung WS, Lee KF, Koistinen R, Koistinen H, Seppala M, Ho PC & Chiu PC** 2006 Roles of glycodefin in modulating sperm function. *Molecular and Cellular Endocrinology* **250** 149–156. (doi:10.1016/j.mce.2005.12.038)
- Yeung WS, Lee KF, Koistinen R, Koistinen H, Seppala M, Ho PC & Chiu PC** 2007 Glycodefin: a molecule with multi-functions on spermatozoa. *Society of Reproduction and Fertility Supplement* **63** 143–151.
- Zhang M, Hong H, Zhou B, Jin S, Wang C, Fu M, Wang S & Xia G** 2006 The expression of atrial natriuretic peptide in the oviduct and its functions in pig spermatozoa. *Journal of Endocrinology* **189** 493–507. (doi:10.1677/joe.1.06483)
- Zhu J, Barrat CL, Lippes J, Pacey AA & Cooke ID** 1994 The sequential effects of human cervical mucus, oviductal fluid, and follicular fluid on sperm function. *Fertility and Sterility* **61** 1129–1135.
- Zumoffen CM, Caille AM, Munuce MJ, Cabada MO & Ghersevich SA** 2010 Proteins from human oviductal tissue- conditioned medium modulate sperm capacitation. *Human Reproduction* **25** 1504–1512. (doi:10.1093/humrep/deq063)
- Zumoffen CM, Gil R, Caille AM, Morente C, Munuce MJ & Ghersevich SA** 2013 A protein isolated from human oviductal tissue *in vitro* secretion, identified as human lactoferrin, interacts with spermatozoa and oocytes and modulates gamete interaction. *Human Reproduction* **28** 1297–1308. (doi:10.1093/humrep/det016)

Received 12 March 2014

First decision 17 April 2014

Revised manuscript received 26 August 2014

Accepted 4 September 2014