Oocyte ageing and epigenetics

Zhao-Jia Ge^{1,2,3}, Heide Schatten⁴, Cui-Lian Zhang^{1,3} and Qing-Yuan Sun²

¹Reproductive Medicine Center, Henan Provincial People's Hospital, #7 Weiwu Road, Jinshui District, Zhengzhou, Henan Province 450003, People's Republic of China, ²State Key Laboratory of Reproductive Biology, Institute of Zoology, Chinese Academy of Sciences, #1 Beichen West Road, Chaoyang District, Beijing 100101, People's Republic of China, ³Reproductive Medicine Center, People's Hospital of Zhengzhou University, Zhengzhou, Henan Province 450003, People's Republic of China and ⁴Department of Veterinary Pathobiology, University of Missouri, Columbia, Missouri 65211, USA

Correspondence should be addressed to C-L Zhang; Email: zcl6086@gmail.com or to Q-Y Sun; Email: sunqy@ioz.ac.cn

Abstract

It has become a current social trend for women to delay childbearing. However, the quality of oocytes from older females is compromised and the pregnancy rate of older women is lower. With the increased rate of delayed childbearing, it is becoming more and more crucial to understand the mechanisms underlying the compromised quality of oocytes from older women, including mitochondrial dysfunctions, aneuploidy and epigenetic changes. Establishing proper epigenetic modifications during oogenesis and early embryo development is an important aspect in reproduction. The reprogramming process may be influenced by external and internal factors that result in improper epigenetic changes in germ cells. Furthermore, germ cell epigenetic changes might be inherited by the next generations. In this review, we briefly summarise the effects of ageing on oocyte quality. We focus on discussing the relationship between ageing and epigenetic modifications, highlighting the epigenetic changes in oocytes from advanced-age females and in post-ovulatory aged oocytes as well as the possible underlying mechanisms.

Reproduction (2015) **149** R103–R114

Introduction

The average age of women bearing children has increased by ~5 years in the past several decades (te Velde & Pearson 2002). However, advanced maternal age has deleterious effects on oocyte maturation and embryonic development, for which the decreasing oocyte quality with ageing may play a key role (Henderson & Edwards 1968, Tarin *et al.* 1998a, Sher *et al.* 2007, Simsek-Duran *et al.* 2013, Di Emidio *et al.* 2014, Rambags *et al.* 2014). These aspects have been well reviewed in our previous publications (Miao *et al.* 2009, Qiao *et al.* 2014); therefore, we summarise what is known on epigenetic alterations and relate changes in the epigenome with alterations in gene expression, mitochondrial dysfunction, nutrition, and hormonal homeostasis (Fig. 1).

The risk of childlessness, stillbirth and multiple births for daughters born by aged mothers (≥ 40 years) is increased when compared with daughters born to young mothers (24–30 years of age) (Smits *et al.* 2002, Sekhon *et al.* 2014, Selemani *et al.* 2014). Several studies have indicated that maternal age is negatively correlated with the health of the offspring (Rocca *et al.* 1991,

Kemkes-Grottenthaler 2004, Brion *et al.* 2008, Gale 2010). The Barker hypothesis suggests that the foetal development process is the origin of diseases in offspring (Barker 1995), and compromised pre-ovulation oocyte quality, especially related with epigenetic changes, may be crucial as well (Ge *et al.* 2014a). Epigenetics, including DNA methylation, histone modification and non-coding RNA, regulate gene expression by changing the conformation of chromosomes other than changing DNA sequences. However, epigenetic modifications may be affected by external and internal factors and the epigenetic changes may be inherited by daughter cells or the offspring (Flanagan *et al.* 2006, Bird 2007, Goldberg *et al.* 2007). Therefore, establishing proper genomic imprinting is a crucial event during oocyte maturation and early embryonic development in humans and other mammalian species. In the mouse female germline, DNA methylation is re-methylated during oocyte maturation after birth and it is completely established at the germinal vesicle (GV) stage. This process is mainly catalysed by DNA methyltransferase 3s (DNMT3s, reviewed by Tomizawa *et al.* (2012)). In mammalian oocytes, the reprogramming process

Figure 1 Schematics about relationship between epigenetic changes in oocytes and advanced maternal age. Advanced maternal age causes decrease in oocyte quality, including expression of HDACs and DNMTs, mitochondrial dysfunction, abnormal nutrition supply, changed levels of hormones and others. Thus, many pathways in oocytes may be disrupted, which may be involved in the process of establishing proper epigenetic modification in oocytes.

takes place during oogenesis for histone modification and non-coding RNAs, too which is well discussed in previous reviews (Gu *et al.* 2010, Hales *et al.* 2011). Thus, if the internal and/or external factors are changed during oogenesis, the epigenetic reprogramming process may be disturbed. For instance, in female mice DNA methylation of imprinted genes in oocytes is altered by maternal diabetes and obesity (Ge *et al.* 2013a,b, 2014b). Furthermore, if one-carbon metabolism is abnormal, DNA methylation is changed in many tissues and it is correlated with diseases (Steegers-Theunissen *et al.* 2013).

Altered epigenetic modification may be an important factor for the complications seen in children of older mothers (Ge *et al.* 2014a,b). In this paper, we review the evidence for epigenetic alterations that have been detected in oocytes and that could be responsible for the age-related effects and the potential mechanisms.

Ageing and epigenetic modifications in tissues

Damage to the reproductive and other systems increases with ageing in humans and in many other complex organisms, and there is a decrease in the adaptability and energy utilisation of the system. The accumulation of damage is caused by external and internal factors. Genetic factors only explain 20–30% of the variation in the human lifespan for twins and long-lived families, although these factors are crucial for survival to very old ages (Herskind *et al.* 1996, Mitchell *et al.* 2001, Poulsen *et al.* 2007). The other 70–80% of the variation may be caused by stochastic events, the environment and other non-genetic factors. Therefore, epigenetics, which is the link between the environment and genes and which regulates gene expression by mechanisms other than changes in the underlying DNA sequence (Goldberg *et al.* 2007), has been recognised as a possible contribution to the ageing phenotype (Wilson & Jones 1983, Fraga *et al.* 2005, McCauley & Dang 2015).

In mammals, epigenetic modifications are established during foetal development and most of them will be maintained throughout life by DNMT1. But the established epigenetic modifications in organs may be changed after birth if the external and/or internal environment is altered. The genome-wide result shows that global hypomethylation is associated with ageing (Liu *et al.* 2003, 2011). Genome-wide analysis shows that ageing is associated with a decrease in global genomic methylation, including CpG-poor promoters and tissue-specific genes (Heyn *et al.* 2012). Similar results have been obtained by other studies in blood and different tissues (Bjornsson *et al.* 2008, Moore *et al.* 2008, Bollati *et al.* 2009, Liu *et al.* 2009a, Kim *et al.* 2010, Gentilini *et al.* 2013, Wnuk *et al.* 2014). During the lifetime of monozygotic twins, the older monozygous twins exhibit remarkable differences not only in genomic distribution of 5-methylcytosine DNA but also in histone acetylation compared with the early years of life (Fraga *et al.* 2005). In the process of postnatal development and ageing of rhesus, dimethylation of histone 3 lysine 4 (H3K4me2) globally increases at promoters and enhancers (Han *et al.* 2012). Small non-coding RNA known as microRNA (miRNA) negatively modulates gene expression through binding to target mRNAs. The expression of miRNA is regulated by DNA methylation (Anwar & Lehmann 2014). Recently, the expression of miRNAs in ageing organs of mice and humans has also been observed (Pincus *et al.* 2011, Inukai *et al.* 2012, Ye *et al.* 2014). In older (>38 years) women, the miRNA profiling in the follicular fluid is clearly different compared with younger ones (<31 years) (Diez-Fraile *et al.* 2014). Although Anwar & Lehmann (2014) discussed that DNA methylation regulates the expression of miRNAs, the detailed mechanisms remained obscure. Besides DNA methylation, histone modifications and miRNAs, the chromatin structure and transposable elements in organisms are also altered with ageing in *Caenorhabditis elegans*, *Drosophila* and mouse model systems and the tissue culture-based replicative senescence model of cell ageing (Wood & Helfand 2013). Although the mechanism causing epigenetic changes in tissues after birth may be different from that in oocytes, the alterations in epigenetic modifications in different organs may indicate that epigenetic modifications in oocytes may be affected by ageing.

Epigenetic changes in ageing oocytes

Herein, 'ageing oocyte' is defined as oocyte ageing that occurs in the ovaries of females who show a progressive decline in oocyte number and poor oocyte quality in ovaries during reproductive ageing (Tatone *et al.* 2008). The pregnancy rate of older women is lower compared with that of younger women, and the lower pregnancy rate (Menken *et al.* 1986) may mainly be caused by

compromised oocyte quality (Wang *et al.* 2012). The child born to an older mother tends to exhibit the onset of some diseases in adulthood (Rocca *et al.* 1991, Kemkes-Grottenthaler 2004). These observations prompted us to ponder the mechanisms underlying this phenomenon. External and internal environmental alterations may cause changes in epigenetic modifications in oocytes. In animal models, the establishment of epigenetic modifications in oocytes is affected by maternal diets (Steegers-Theunissen *et al.* 2013), non-communicable diseases (Ge *et al.* 2013b) and other factors (Li *et al.* 2014). If the changes in epigenetic modifications occur in the germline, it would affect embryonic development and health of the offspring, and this effect may even extend to further generations (Ge *et al.* 2014b). Therefore, we discuss the association of oocyte quality, highlighting the epigenetic modifications, with maternal ageing.

Ageing and oocyte quality

Many studies in the IVF–ET setting have demonstrated that ageing does not appear to affect the ability of oocytes to become fertilised, but compared with younger women, the implantation rates are lower and the spontaneous abortion rates are higher for older women (Romeu *et al.* 1987, Warburton *et al.* 1987, Lim & Tsakok 1997). Navot *et al.* (1991) reported that the age-related decline in female fertility is attributed to oocyte quality. For young mothers, about 20% of oocytes are aneuploid (Hassold *et al.* 2007), but the percentage increases to 50% or more in the oocytes of older mothers (Fragouli *et al.* 2011, Handyside *et al.* 2012). One reason may be that the level of cohesin, which is the key protein that regulates chromosome separation, falls below the level required to stabilise chiasmata and to hold sister centromeres tightly together in pre-ovulatory ageing oocytes (Lister *et al.* 2010, Jessberger 2012). Cohesion loss may be responsible for age-related meiotic segregation errors in mammalian oocytes (Petronczki *et al.* 2003, Revenkova *et al.* 2004, Tsutsumi *et al.* 2014). In humans, the expression of REC8 and SMC1B, the subunit of cohesin, in oocytes is decreased in 40-year-old women compared with 20-year-old women (Tsutsumi *et al.* 2014). There are other factors that may induce oocyte aneuploidy, which have been well reviewed recently (Jones & Lane 2013). Mitochondrial dysfunctions are observed in pre-ovulatory ageing oocytes (Dorland *et al.* 1998, Bentov *et al.* 2011, Eichenlaub-Ritter *et al.* 2011). As the amount of mitochondria decreases, the mutation of mitochondrial DNA may become increased and mitochondrial function is also affected in the oocytes of advanced-age female bovine, hamsters and mice (Eichenlaub-Ritter *et al.* 2011, Iwata *et al.* 2011, Simsek-Duran *et al.* 2013). Gene expression and chromatin structure (Wood & Helfand 2013) are also found to be affected by ageing. Hamatani *et al.* (2004)

compared the difference in the expression profile at the transcript level of metaphase II (MII) oocytes of 5- to 6-week-old mice with that of 42- to 45-week-old mice. Among ~11 000 genes whose transcripts are detected in oocytes, 5% showed obvious expression changes (Hamatani *et al.* 2004). Similar expression profiles in aged mouse oocytes have been reported in another study (Pan *et al.* 2008). An alteration in the expression profile of human MII oocytes is also associated with female ageing (Grondahl *et al.* 2010, Santonocito *et al.* 2013). In mouse oocytes, the mRNA and protein expression levels are altered with ageing, and dysfunctions of the ageing ovary may be a reason for the altered expression of mRNAs and proteins in pre-ovulatory ageing oocytes (Schwarzer *et al.* 2014, Tatone *et al.* 2014).

Ageing oocytes and epigenetics

Previous discussions indicate that the epigenetic modifications of oocytes may be affected by advanced maternal age because the expression of DNMTs and histone acetyltransferases (*Myst1* (*Kat8*) and *Mrgx* (*Morf412*); Hamatani *et al.* 2004) is altered with ageing. Therefore, the changes in epigenetic modifications may partly explain why the child of an older woman is predisposed to the onset of hypertension, obesity and other diseases in adulthood (Aagesen *et al.* 1984, Malini & Ramachandra 2006).

Ageing oocytes and DNA methylation

If DNMT1, which maintains the DNA methylation patterns in oocytes and embryos (Mertineit *et al.* 1998), is deleted in mouse oocytes, the embryos show a loss of allele-specific expression and methylation at certain imprinted loci, and the fetuses of homozygous females die during the last third of gestation (Howell *et al.* 2001). In 5- to 6-week-old mouse oocytes, the gene expression profile is different from that in 42- to 45-week-old mouse oocytes. DNMT1, 3b and 3l are involved in the differential gene expression (Hamatani *et al.* 2004). Yue *et al.* (2012) found that the changes in genome-wide DNA methylation in oocytes and preimplantation embryos of 35- to 40-week-old mice were associated with decreased expression of DNMTs. The pregnancy rate of older Kunming mice (35–40 weeks old) is lower than that of younger mice, and the stillbirth and foetal malformation rate are higher in the older group compared with the younger group, which may be associated with abnormal DNA methylation in oocytes (Yue *et al.* 2012). In humans, TAP73 expression which is regulated by DNA methylation patterns is lower in the oocytes of women older than 38 years of age compared with the oocytes of women younger than 36 years of age (Guglielmino *et al.* 2011). There are still no direct proofs that the DNA methylation status in human oocytes is affected by ageing.

However, one study (Lopes *et al.* 2009) reports that the increase in resorption sites, morphological abnormalities

and delayed development are related with the age of C57BL/6 mice (43–47 weeks old), but the monoallelic expression of the imprinted genes *H19* and *Snrpn* is normal in the blastocysts of aged female mice and the DNA methylation patterns of the differentially methylated regions (DMRs) of *Snrpn*, *Kcnq1ot1*, *U2af1-rs1* (*Zrsr1*), *Peg1*, *Igf2r* and *H19* are not altered. By Restriction Landmark Genome Scanning, the investigators also did not find significant differences in genome-wide DNA methylation in embryos and placentas from aged female mice (Lopes *et al.* 2009). This is contradictory to previous reports (Hamatani *et al.* 2004). The authors propose that the contradiction may be related to the materials selected and the limitation of the technique utilised in their study (Lopes *et al.* 2009). During zebrafish ageing, two CpG island shores are hypomethylated in oocytes, but they are *de novo* methylated in fertilised eggs (Shimoda *et al.* 2014). This suggests that the loss of methylation might be reset after fertilisation because there is a de-methylation and re-methylation process during early embryonic development.

Although reports about DNA methylation changes in oocytes from advanced-age females are contradictory, the popular viewpoint proposes that the DNA methylation in oocytes may be changed by pre-ovulatory ageing.

Ageing oocytes and histone modifications

Histone modifications, including methylation, acetylation, ubiquitination and other modifications, represent another crucial and well-investigated epigenetic modification. During meiosis, histone is deacetylated globally at the MI and MII stages by histone deacetylase (HDAC) activity in mammalian oocytes as revealed by

immunostaining (Kim *et al.* 2003, Akiyama *et al.* 2004, Reddy & Villeneuve 2004). Akiyama *et al.* (2006) reported that if meiotic histone deacetylation was inhibited, aneuploidy occurred in fertilised mouse oocytes and this resulted in embryonic death in the uterus at an early stage of development. HDAC is downregulated at transcript level in ageing mouse (42- to 45-week-old) oocytes (Hamatani *et al.* 2004) although histone still remains acetylated in the oocytes of 10-month-old female mice (Akiyama *et al.* 2006). This suggests that histone modification in pre-ovulatory ageing oocytes may be affected (Table 1) and during development it may result in embryonic death. Similar results were obtained by Manosalva & Gonzalez (2009) and Suo *et al.* (2010). The expression of *Sirt2* which is related with the acetylation of histone H4K16 in the oocytes of aged mice is lower compared with that in younger mouse oocytes (Zhang *et al.* 2014a).

Another study found that the histone methylation in mouse GV oocytes was affected by advanced maternal age (Manosalva & Gonzalez 2010) (Table 1). Concomitantly, the GV and MII oocytes of older females lack H3K9me3, H3K36me2, H3K79me2 and H4K20me2 compared with the GV and MII oocytes of younger females. Meanwhile, the expression of the histone methylation-related factors (*Cbx1* and *Sirt1*) is changed in ageing GV oocytes. Histone 3 lysine 4 methylation in mouse GV oocytes is also changed by ageing (Shao *et al.* 2015).

In humans, the mRNA expression profile of MII oocytes is related with ageing. The differently expressed genes are involved in many biological processes, such as cell cycle, metabolism, apoptosis, protein modification and others (Grondahl *et al.* 2010). Recently, van den Berg *et al.* have

Table 1 Effects of advanced maternal age on epigenetics in oocytes.

Species	Epigenetic changes	References
Mouse	Genome-wide DNA methylation is lower in 35- to 40-week-old mouse oocytes	Yue <i>et al.</i> (2012)
Mouse	DNMTs (DNMT1, 3a, 3b and 3l) expression is decreased in aged mouse oocytes	Hamatani <i>et al.</i> (2004) and Pan <i>et al.</i> (2008)
Mouse	Histone deacetylase is downregulated and histone remains acetylated in older mouse oocytes. Histone acetylation of H4K12 is affected in aged GV and MII oocytes	Hamatani <i>et al.</i> (2004), Akiyama <i>et al.</i> (2006), Manosalva & Gonzalez (2009) and Suo <i>et al.</i> (2010)
Mouse	<i>Sirt2</i> expression is lower in the oocytes of old mice compared with young mice	Zhang <i>et al.</i> (2014a,b)
Mouse	H3K9me3, H3K36me2, H3K79me2 and H4K20me2 are altered in aged oocytes	Manosalva & Gonzalez (2010)
Mouse	The expression of <i>Cbx1</i> and <i>Sirt1</i> is changed in the oocytes of older mice	Shao <i>et al.</i> (2014)
Mouse	Histone 3 lysine 4 methylation is changed in aged GV oocytes	
Bovine	Non-imprinted genes (<i>SLC2A1</i> , <i>PRDX1</i> , <i>ZAR1</i> and <i>BTS</i>) are hypomethylated in the oocytes of adult cows compared with prepubertal calves	Diederich <i>et al.</i> (2012)
Zebrafish	Two CpG island shores hypomethylated in oocytes with ageing	Shimoda <i>et al.</i> (2014)
Human	The deacetylation of H4K12 in human MII oocytes is affected at an age-dependent manner	van den Berg <i>et al.</i> (2011)
Human	The expression of ubiquitin I, <i>USP2</i> , <i>USP34</i> and <i>USP42</i> is affected in the oocytes of women aged 37–39 years	Grondahl <i>et al.</i> (2010)
Human	MicroRNAs expression profiling of the follicular fluid of younger females is different from that of the follicular fluid of older females	Diez-Fraile <i>et al.</i> (2014)
Equine	Three miRNAs are expressed in significantly higher amounts in exosomes isolated from follicular fluid of old compared to young mares	da Silveira <i>et al.</i> (2012)

shown that the histone acetylation staining of H4K5, H4K8, H4K12 and H4K16 was intensive in GV oocytes; however in MI and MII oocytes, chromatin was deacetylated in variable proportions. They also investigated the relationship between histone acetylation and maternal age. The results indicate that advanced maternal age negatively influences the deacetylation of H4K12 in human MII oocytes (van den Berg *et al.* 2011).

For women at aged 37–39 years, the gene for ubiquitin 1 (a ubiquitin-like protein) is downregulated in the oocytes, but three genes for the ubiquitin-specific peptidases *USP2*, *USP34* and *USP42* are upregulated (Grondahl *et al.* 2010). This suggests that the ubiquitination may be affected by age in human oocytes (Steuerwald *et al.* 2007). This indicates that histone ubiquitination might also be affected in oocytes by ageing, but there is still no solid evidence to confirm it.

Ageing oocytes and miRNA

miRNA is a kind of small non-coding RNA which functions in post-transcriptional regulation of gene expression upon recruitment into effector complexes (miRNA protein complexes or microRNPs; Truesdell *et al.* 2012). The post-transcriptional regulation may be particularly crucial for early mammalian development, from maturation of the germ line to initiation of gastrulation, because the genome is transcriptionally silent from the fully grown oocyte stage until zygotic genome activation (Abe *et al.* 2010). Small RNA is present in mouse oocytes, including miRNA (Tam *et al.* 2008), but Suh *et al.* (2010) suggest that miRNA function is globally suppressed during oocyte maturation and preimplantation development. However, loss of *Dicer*, which is crucial for the generation of endo-siRNA and miRNA, in mouse oocytes results in severe spindle and chromosomal segregation defects (Murchison *et al.* 2007), while loss of *Dgcr8*, which is essential only for miRNA processing, in the mouse has no effects on mRNA expression (Suh *et al.* 2010). Once *Ago2* is knocked out in mouse oocytes, the phenotype is similar to that observed in *Dicer*-knockout mouse oocytes (Kaneda *et al.* 2009). These results suggest that miRNAs function may be suppressed in mouse oocytes. However, miRNA mediates mRNA translation activation by FXR1 in *Xenopus laevis* oocytes (Truesdell *et al.* 2012). In mouse oocytes, miRNA-335-5p could affect oocyte maturation by regulating cytoskeleton dynamics (Cui *et al.* 2013). A similar result that miRNA-27a activation is not suppressed is observed in porcine oocytes (Chen *et al.* 2012). Many studies also demonstrated that miRNAs were essential for follicle development in different species (Abramov *et al.* 2013, Sohel *et al.* 2013, Yang *et al.* 2013, Zhang *et al.* 2014b). miRNAs expressed in oocytes could regulate bovine early embryogenesis (Tripurani *et al.* 2013). This contradiction has been well reviewed in previously published papers (Svoboda & Flemer 2010, Suh & Blelloch 2011).

It was shown that 79 miRNAs and 41 miRNAs existed in the microvesicles and exosomes isolated from equine follicular fluid respectively (Table 1), and three miRNAs are expressed significantly higher in exosomes isolated from follicular fluid of old mares compared with young ones (da Silveira *et al.* 2012). In humans, miRNAs are abundant in MII oocytes and cumulus cells and they may be essential for follicle development (Assou *et al.* 2013). Moreover, the miRNA expression profile in follicular fluid of women with polycystic ovary syndrome or premature ovarian failure is different from that in follicular fluid of unaffected women (Roth *et al.* 2014). In addition, miRNAs expression profiling of the follicular fluid of younger (<31 years) and older (>38 years) individuals was also investigated and the result showed that the expression of four miRNAs is different. These miRNAs are involved in carbohydrate digestion and absorption, p53 signalling and other biological processes that may be related with fertility. Therefore, this set of miRNAs and their respective targets should be evaluated in relationship with reproductive ageing (Diez-Fraile *et al.* 2014). However, the correlation between miRNA expression and oocyte quality during maternal ageing is still unknown.

Post-ovulatory ageing of oocytes and epigenetic modifications

After ovulation, when the arrested MII oocytes are not fertilised during the window of the optimal fertilisation time *in vivo* or *in vitro*, the unfertilised oocytes undergo a time-dependent decline concerning quality and this is called 'post-ovulatory ageing of oocytes' (Liang *et al.* 2012). Studies indicate that post-ovulatory ageing of mouse oocytes decreases the pregnancy rate, litter size and increases the percentage of male offspring compared with control females (Tarin *et al.* 1999, 2002, Kosubek *et al.* 2010, Liang *et al.* 2011). F1 offspring derived from post-ovulatory ageing oocytes are prone to the onset of growth retardation, delayed development of the righting reflex and emotionality (Tarin *et al.* 1999). Otherwise, post-ovulatory ageing of mouse oocytes decreases reproductive fitness and longevity of offspring (Tarin *et al.* 2002). This suggests that the epigenetic modification in post-ovulatory ageing of oocytes may be altered (Table 2). In our laboratory, we have analysed methylation patterns of imprinted genes in mouse oocytes during the post-ovulatory process. We examined methylation patterns of *Snrpn* and *Peg1* in *in vivo* and *in vitro* oocytes at 13, 21 and 29 h of human chorionic gonadotrophin (hCG) injection, respectively, and loss of methylation was observed at 29 h of hCG injection (Liang *et al.* 2008). Imamura *et al.* (2005) also reported *Peg1* lost methylation in oocytes during post-ovulatory ageing. However, only a small number of oocytes showed aberrant methylation in the DMR of *Peg3* in offspring derived from post-ovulatory ageing mouse

Table 2 Effects of post-ovulatory ageing of oocytes on epigenetic modifications.

Species	Epigenetic changes	References
Mouse	DNA methylation patterns of <i>Snrpn</i> and <i>Peg1</i> in oocytes is altered at 29 h after hCG injection	Liang <i>et al.</i> (2008)
Mouse	<i>Peg1</i> loss of methylation in oocytes during post-ovulatory ageing	Imamura <i>et al.</i> (2005)
Mouse	Acetylation of H4K8, H4K12 and H3K14 is altered in ageing oocytes	Huang <i>et al.</i> (2007) and Liu <i>et al.</i> (2009b)
Porcine	Histone of H4K12 is changed in post-ovulatory ageing oocytes	Cui <i>et al.</i> (2011)

oocytes (Liang *et al.* 2011). Although the DNA methylation of some imprinted genes is influenced by oocyte ageing, whether the whole-genome methylation patterns and histone modification are affected by oocyte post-ovulatory ageing is unclear. The histone modifications in post-ovulatory ageing of mouse oocytes are changed at 19 h of hCG injection compared with that at 14 h of hCG injection. When extending the time to 24 h of hCG injection, the fluorescence signals of acetylation of H3K14 also increased in oocytes (Huang *et al.* 2007). Another study reported that histone acetylation of H3K14 and H4K12 increased in mouse oocytes during post-ovulatory ageing (Liu *et al.* 2009b). This phenomenon is also observed in porcine oocytes during post-ovulatory ageing (Cui *et al.* 2011), although, the detailed underlying mechanism is still unknown.

Oocyte ageing and epigenetics: underlying mechanism(s)

Changes in enzymes related with epigenetic modifications. Epigenetic modifications are catalysed by numerous proteins, including DNMTs, ten-eleven-translocations (TETs), HDACs, ZFP57, Dicer, and KAP1/TRIM28. If their expression is affected in oocytes, the epigenetic modifications may be altered. DNMT1 and DNMT10 are crucial for maintaining proper methylation and DNMT3a, b and l are key *de novo* methylation enzymes. When the expression of DNMT1, which is necessary for maintaining DNA methylation, is disrupted in mouse oocytes, DNA methylation of imprinted genes is not maintained properly during early embryonic development (Hirasawa *et al.* 2008, Kurihara *et al.* 2008). The changed expression of DNMTs in oocytes from individuals of advanced maternal age (Hamatani *et al.* 2004, Grondahl *et al.* 2010) might be the direct reason for causing the DNA methylation alterations (Fig. 1). In MII oocytes of 35- to 40-week-old mice, the protein expression of DNMT1, DNMT3a, DNMT3b and DNMT3l is obviously lower than those in MII oocytes of 6- to 8-week-old mice (Yue *et al.* 2012). Anckaert *et al.* (2013) used 14-day *in vitro* follicle culture as a model to investigate pre-ovulatory intrafollicular

oocyte 'ageing' and found that the mRNA levels of *Dnmt3a*, *Dnmt3l* and *Zfp57* were altered compared with 12-day *in vitro* follicle culture in the mouse.

The expression of enzymes related with histone modification is also affected by advanced maternal ageing in oocytes. For example, *Hdac2* is downregulated and the expression of histone acetyltransferases (*Myst1* and *Mrgx*) decreases in aged mouse oocytes (Hamatani *et al.* 2004). In old mice, the expression of the histone methylation-related factors *Cbx1* and *Sirt2* was changed in GV oocyte (Manosalva & Gonzalez 2010). The protein and kinase activities of CDC2A decreased in the GV and MII oocytes of old mice (Manosalva & Gonzalez 2009). Although specific inhibitors of HDACs could delay post-ovulatory oocyte ageing in mice and pigs (Huang *et al.* 2007, Jeseta *et al.* 2008, Lee *et al.* 2013), it is still unclear whether the change in histone acetylation is caused by the altered level or activities of HDACs for these post-ovulatory ageing oocytes from young mice.

Other enzymes related with epigenetic modifications, for example TETs (Yamaguchi *et al.* 2012), Trim28 (Messerschmidt *et al.* 2012) and Dicer (Murchison *et al.* 2007), may also be critical for oocyte developmental potential. However, there are few studies to investigate whether their expression in oocytes is affected by ageing or not. The proteins of the TET family are not only involved in DNA demethylation during early embryo development (Tahiliani *et al.* 2009, Gu *et al.* 2011, Ma *et al.* 2012), they are also crucial for female germ cells to complete meiosis (Yamaguchi *et al.* 2012). If the expression of *Tet3* is suppressed in oocytes, the paternal global demethylation process at the zygote stage is impaired (Gu *et al.* 2011). Although TETs are associated with ageing and diseases (van den Hove *et al.* 2012), a role for them in pre-ovulatory ageing oocyte and post-ovulatory ageing oocyte is still undetermined.

Changes in mitochondrial activity. Mitochondrial dysfunction may be another crucial factor inducing epigenetic changes in the oocytes of advanced females (Fig. 1). Oocyte maturation includes nuclear maturation and cytoplasmic maturation, and there are many biological events involved in these two processes, such as gene expression and histone and chromatin modifications (Eppig 1996). During oocyte maturation, energy (ATP) required is supplied by mitochondria (Torner *et al.* 2004). Histone and DNMTs use S-adenosyl-L-methionine (SAM) as a donor of methyl groups. SAM is biosynthesised using methionine and ATP (Igarashi & Katoh 2013). Wellen *et al.* (2009) demonstrated that ATP-citrate lyase was necessary for histone modification. Therefore, we conclude that if mitochondrial function is compromised by maternal age, the epigenetic modification may also be affected. In aged mouse oocytes, ATP and mitochondrial genomes are reduced to 38.4 and 44% respectively (Simsek-Duran *et al.* 2013). In bovine oocytes, the number of mitochondria and content of ATP are also affected by maternal age (Iwata *et al.* 2011).

In post-ovulatory aged oocytes, reactive oxygen species (ROS) may be another factor inducing epigenetic changes. During *in vitro* ageing in porcine oocyte, H4K12 acetylation levels are related with ooplasmic ROS content (Cui *et al.* 2011). Glucose level is related with *de novo* purine and cAMP synthesis, which is associated with nuclear maturation of oocytes (Colton *et al.* 2003). The cumulus cells supply nutrition to oocytes in the final phase of oocyte maturation (Gilchrist *et al.* 2008). The function of cumulus cells is compromised by maternal age (Tatone & Amicarelli 2013). Thus, the energy transmitted from cumulus cells decreases, which may affect the establishment of epigenetic modifications in aged oocytes.

Nutrition effect. The changed nutrition supplies, especial one-carbon, may contribute to epigenetic changes in aged oocytes (Anckaert *et al.* 2010, Steegers-Theunissen *et al.* 2013; Fig. 1). For instance, the folate status declines with ageing, including decreased folate intake and altered folate availability (Jacob *et al.* 1998, Rampersaud *et al.* 2000). Therefore, the disruption of folate-mediated one-carbon metabolism by ageing may be another reason causing the abnormal DNA methylation in aged oocytes (Jacques *et al.* 2001).

Possible hormonal effects. A study demonstrated that oestrogen replacement therapy in menopause women reduced the total plasma homocysteine concentration and increased genomic DNA methylation of mononuclear cells (Friso *et al.* 2007). In rodent brain, the methylation percent on the promoter of oestrogen receptor alpha is modulated by the expression of oestrogen (Schwarz *et al.* 2010). The DNA methylation is not only affected by hormones present in tissues, may be also in oocytes. In humans and mice, the methylation patterns of *Peg1* and *H19* are changed by hormones used for superovulation of growing oocytes (Sato *et al.* 2007). With ageing, the androgen level, which can be converted to oestrogen, is reduced for females (Blevins *et al.* 2013). The DNMT proteins and transcripts in the livers of 3-, 12-, and 24-month-old Ames dwarf mice are drastically reduced compared with WT siblings, and growth hormone appears to modulate the expression of DNMT1 and 3a (Armstrong *et al.* 2014). The amount of oestrogen which is important for follicular development decreases with ageing (Olsen & Kovacs 1996). Although there is no direct evidence showing that the epigenetic modifications are affected by hormones in pre- and post-ovulatory aged oocytes, it might contribute to the changes in epigenetic modifications in aged oocytes, which needs experimental validation (Fig. 1).

Conclusion(s) and perspectives

The above discussions suggest that advanced maternal age and post-ovulatory oocyte ageing are deleterious to oocyte quality, including oocyte maturation, chromosome segregation, epigenetic modifications and mitochondrial function, and the health of the offspring

from advanced-age mothers may also be affected by compromised oocyte quality (Takeo *et al.* 2013a). In this review, we mainly discussed the possible relationship between advanced maternal age and epigenetic modifications in oocytes and the potential underlying mechanisms. Two major problems are still unresolved: i) the detailed mechanisms underlying compromised oocyte quality including epigenetic changes caused by advanced maternal age and ii) the prevention of the adverse effects of oocyte ageing on epigenetic changes. As discussed previously, changes in enzymes including methyltransferases (DNMTs) and demethylases (TETs) may be the direct reasons for epigenetic alterations in aged oocytes, but whether/how ageing induces the changes in their expression needs further clarification. With the application of new technologies, it has been possible to test the transcriptome, global DNA methylation, histone modifications and proteome in limited number of cells or even in a single cell (Guo *et al.* 2014, Lovatt *et al.* 2014, Smallwood *et al.* 2014), which may help to address this issue. Another important study still to pursue is how to prevent the age-related deleterious effects on oocytes. If the disulphide-reducing agent dithiothreitol (DTT), an antioxidant, is supplemented to the culture medium, the negative effects of post-ovulatory ageing of mouse oocytes *in vitro* on fertilisation, cellular fragmentation at 24 h post-insemination and the potential of embryos for development until the blastocyst stage are prevented, at least in part (Tarin *et al.* 1998b). *N*-acetyl-cysteine (NAC) supplemented to medium can also decrease ROS levels in post-ovulatory aged oocytes, but only resveratrol increased the fertilisation rate (Takeo *et al.* 2013b). Whether the epigenetic changes in these oocytes are prevented is unknown. But another study demonstrated that if adult female mice were subjected to caloric restriction, they did not exhibit age-related increases in oocyte aneuploidy, meiotic spindle abnormalities or mitochondrial dysfunction, all of which occurred in the oocytes of age-matched controls (Selesniemi *et al.* 2011). This indicates that the epigenetic modification in aged oocytes may also be affected by calorie restricted diet because energy is crucial for epigenetic modifications. In medium supplemented with pyruvate, post-ovulatory oocyte ageing is prevented and changes in histone acetylation are corrected in the mouse oocyte (Liu *et al.* 2009b). Some studies suggest that both DNA methylation and histone modification are associated with caloric restriction (Li *et al.* 2011, Chouliaras *et al.* 2012, Chen *et al.* 2013). One-carbon supplies methyl for methylation, so changing nutrients in diet might prevent abnormal methylation in oocytes. If the levels of methyl donor are lower in the medium during mouse follicle culture, the establishment of oocyte imprinting is affected (Anckaert *et al.* 2010). Steegers-Theunissen *et al.* (2013) reviewed the effects of one-carbon on reproduction and long-term health of

offspring. These studies suggest that age-related effects, including epigenetic changes, on oocytes might be prevented by diets, medicine or other methods. However, until now we still cannot effectively prevent the age-related deleterious effects on oocytes.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the review.

Funding

This work is supported by the National Natural Science Foundation of China (3147055) and the National Basic Research Program of China (2012CB944404, 2011CB944501).

Acknowledgements

The authors thank all the members of the Reproductive Medicine Center, Henan Provincial People's Hospital for their help with this work.

References

- Aagesen L, Grinstead J & Mikkelsen M 1984 Advanced grandmaternal age on the mother's side – a risk of giving rise to trisomy 21. *Annals of Human Genetics* **48** 297–301. (doi:10.1111/j.1469-1809.1984.tb00843.x)
- Abe K, Inoue A, Suzuki MG & Aoki F 2010 Global gene silencing is caused by the dissociation of RNA polymerase II from DNA in mouse oocytes. *Journal of Reproduction and Development* **56** 502–507. (doi:10.1262/jrd.10-068A)
- Abramov R, Fu G, Zhang Y & Peng C 2013 Expression and regulation of miR-17a and miR-430b in zebrafish ovarian follicles. *General and Comparative Endocrinology* **188** 309–315. (doi:10.1016/j.ygcen.2013.02.012)
- Akiyama T, Kim JM, Nagata M & Aoki F 2004 Regulation of histone acetylation during meiotic maturation in mouse oocytes. *Molecular Reproduction and Development* **69** 222–227. (doi:10.1002/mrd.20121)
- Akiyama T, Nagata M & Aoki F 2006 Inadequate histone deacetylation during oocyte meiosis causes aneuploidy and embryo death in mice. *PNAS* **103** 7339–7344. (doi:10.1073/pnas.0510946103)
- Anckaert E, Romero S, Adriaenssens T & Smits J 2010 Effects of low methyl donor levels in culture medium during mouse follicle culture on oocyte imprinting establishment. *Biology of Reproduction* **83** 377–386. (doi:10.1095/biolreprod.109.082164)
- Anckaert E, Sanchez F, Billooye K & Smits J 2013 Dynamics of imprinted DNA methylation and gene transcription for imprinting establishment in mouse oocytes in relation to culture duration variability. *Biology of Reproduction* **89** 130. (doi:10.1095/biolreprod.113.111641)
- Anwar SL & Lehmann U 2014 DNA methylation, microRNAs, and their crosstalk as potential biomarkers in hepatocellular carcinoma. *World Journal of Gastroenterology* **20** 7894–7913. (doi:10.3748/wjg.v20.i24.7894)
- Armstrong VL, Rakoczy S, Rojanathammanee L & Brown-Borg HM 2014 Expression of DNA methyltransferases is influenced by growth hormone in the long-living ames dwarf mouse *in vivo* and *in vitro*. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences* **69** 923–933. (doi:10.1093/gerona/glt133)
- Assou S, Al-edani T, Haouzi D, Philippe N, Lecellier CH, Piquemal D, Commes T, Ait-Ahmed O, Dechaud H & Hamamah S 2013 MicroRNAs: new candidates for the regulation of the human cumulus–oocyte complex. *Human Reproduction* **28** 3038–3049. (doi:10.1093/humrep/det321)
- Barker DJ 1995 Fetal origins of coronary heart disease. *BMJ* **311** 171–174. (doi:10.1136/bmj.311.6998.171)
- Bentov Y, Yavorska T, Esfandiari N, Jurisicova A & Casper RF 2011 The contribution of mitochondrial function to reproductive aging. *Journal of Assisted Reproduction and Genetics* **28** 773–783. (doi:10.1007/s10815-011-9588-7)
- van den Berg IM, Eleveld C, van der Hoeven M, Birnie E, Steegers EA, Galjaard RJ, Laven JS & van Doorninck JH 2011 Defective deacetylation of histone 4 K12 in human oocytes is associated with advanced maternal age and chromosome misalignment. *Human Reproduction* **26** 1181–1190. (doi:10.1093/humrep/der030)
- Bird A 2007 Perceptions of epigenetics. *Nature* **447** 396–398. (doi:10.1038/nature05913)
- Bjornsson HT, Sigurdsson MI, Fallin MD, Irizarry RA, Aspelund T, Cui H, Yu W, Rongione MA, Ekstrom TJ, Harris TB *et al.* 2008 Intra-individual change over time in DNA methylation with familial clustering. *Journal of the American Medical Association* **299** 2877–2883. (doi:10.1001/jama.299.24.2877)
- Blevins JK, Coxworth JE, Herndon JG & Hawkes K 2013 Brief communication: Adrenal androgens and aging: female chimpanzees (*Pan troglodytes*) compared with women. *American Journal of Physical Anthropology* **151** 643–648. (doi:10.1002/ajpa.22300)
- Bollati V, Schwartz J, Wright R, Litonjua A, Tarantini L, Suh H, Sparrow D, Vokonas P & Baccarelli A 2009 Decline in genomic DNA methylation through aging in a cohort of elderly subjects. *Mechanisms of Ageing and Development* **130** 234–239. (doi:10.1016/j.mad.2008.12.003)
- Brion MJ, Leary SD, Lawlor DA, Smith GD & Ness AR 2008 Modifiable maternal exposures and offspring blood pressure: a review of epidemiological studies of maternal age, diet, and smoking. *Pediatric Research* **63** 593–598. (doi:10.1203/PDR.0b013e31816fdbd3)
- Chen L, Hu X, Dai Y, Li Q, Wang X, Xue K, Li Y, Liang J, Wang Y, Liu X *et al.* 2012 MicroRNA-27a activity is not suppressed in porcine oocytes. *Frontiers in Bioscience* **4** 2679–2685. (doi:10.2741/E574)
- Chen PY, Ganguly A, Rubbi L, Orozco LD, Morselli M, Ashraf D, Jaroszewicz A, Feng S, Jacobsen SE, Nakano A *et al.* 2013 Intrauterine calorie restriction affects placental DNA methylation and gene expression. *Physiological Genomics* **45** 565–576. (doi:10.1152/physiol-genomics.00034.2013)
- Chouliaras L, van den Hove DL, Kenis G, Keitel S, Hof PR, van Os J, Steinbusch HW, Schmitz C & Rutten BP 2012 Prevention of age-related changes in hippocampal levels of 5-methylcytidine by caloric restriction. *Neurobiology of Aging* **33** 1672–1681. (doi:10.1016/j.neurobiolaging.2011.06.003)
- Colton SA, Humpherson PG, Leese HJ & Downs SM 2003 Physiological changes in oocyte–cumulus cell complexes from diabetic mice that potentially influence meiotic regulation. *Biology of Reproduction* **69** 761–770. (doi:10.1095/biolreprod.102.013649)
- Cui MS, Wang XL, Tang DW, Zhang J, Liu Y & Zeng SM 2011 Acetylation of H4K12 in porcine oocytes during *in vitro* aging: potential role of ooplasmic reactive oxygen species. *Theriogenology* **75** 638–646. (doi:10.1016/j.theriogenology.2010.09.031)
- Cui XS, Sun SC, Kang YK & Kim NH 2013 Involvement of microRNA-335-5p in cytoskeleton dynamics in mouse oocytes. *Reproduction, Fertility, and Development* **25** 691–699. (doi:10.1071/RD12138)
- Di Emidio G, Falone S, Vitti M, D'Alessandro AM, Vento M, Di Pietro C, Amicarelli F & Tatone C 2014 SIRT1 signalling protects mouse oocytes against oxidative stress and is deregulated during aging. *Human Reproduction* **29** 2006–2017. (doi:10.1093/humrep/deu160)
- Diederich MI, Hansmann T, Heinzmann J, Barg-Kues B, Herrmann D, Aldag P, Baulain U, Reinhard R, Kues W, Weissberger C, Haaf T & Niemann H 2012 DNA methylation and mRNA expression profiles in bovine oocytes derived from prepubertal and adult donors. *Reproduction* **144** 319–330. (doi:10.1530/REP-12-0134)
- Diez-Fraile A, Lammens T, Tillemann K, Witkowski W, Verhasselt B, De Sutter P, Benoit Y, Espeel M & D'Herde K 2014 Age-associated differential microRNA levels in human follicular fluid reveal pathways potentially determining fertility and success of *in vitro* fertilization. *Human Fertility* **17** 90–98. (doi:10.3109/14647273.2014.897006)
- Dorland M, van Kooij RJ & te Velde ER 1998 General ageing and ovarian ageing. *Maturitas* **30** 113–118. (doi:10.1016/S0378-5122(98)00066-8)
- Eichenlaub-Ritter U, Wiczorek M, Luke S & Seidel T 2011 Age related changes in mitochondrial function and new approaches to study redox regulation in mammalian oocytes in response to age or maturation conditions. *Mitochondrion* **11** 783–796. (doi:10.1016/j.mito.2010.08.011)

- Eppig JJ 1996 Coordination of nuclear and cytoplasmic oocyte maturation in eutherian mammals. *Reproduction, Fertility, and Development* **8** 485–489. (doi:10.1071/RD9960485)
- Flanagan JM, Popenklyte V, Pozdniakovaite N, Sobolev M, Assadzadeh A, Schumacher A, Zangeneh M, Lau L, Virtanen C, Wang SC *et al.* 2006 Intra- and interindividual epigenetic variation in human germ cells. *American Journal of Human Genetics* **79** 67–84. (doi:10.1086/504729)
- Fraga MF, Ballestar E, Paz MF, Ropero S, Setien F, Ballestar ML, Heine-Suner D, Cigudosa JC, Urioste M, Benitez J *et al.* 2005 Epigenetic differences arise during the lifetime of monozygotic twins. *PNAS* **102** 10604–10609. (doi:10.1073/pnas.0500398102)
- Fragouli E, Alfarawati S, Goodall NN, Sanchez-Garcia JF, Colls P & Wells D 2011 The cytogenetics of polar bodies: insights into female meiosis and the diagnosis of aneuploidy. *Molecular Human Reproduction* **17** 286–295. (doi:10.1093/molehr/gar024)
- Friso S, Lamón-Fava S, Jang H, Schaefer EJ, Corrocher R & Choi SW 2007 Oestrogen replacement therapy reduces total plasma homocysteine and enhances genomic DNA methylation in postmenopausal women. *British Journal of Nutrition* **97** 617–621. (doi:10.1017/S0007114507433013)
- Gale EA 2010 Maternal age and diabetes in childhood. *BMJ* **340** c623. (doi:10.1136/bmj.c623)
- Ge ZJ, Liang QX, Luo SM, Wei YC, Han ZM, Schatten H, Sun QY & Zhang CL 2013a Diabetic uterus environment may play a key role in alterations of DNA methylation of several imprinted genes at mid-gestation in mice. *Reproductive Biology and Endocrinology* **11** 119. (doi:10.1186/1477-7827-11-119)
- Ge ZJ, Liang XW, Guo L, Liang QX, Luo SM, Wang YP, Wei YC, Han ZM, Schatten H & Sun QY 2013b Maternal diabetes causes alterations of DNA methylation statuses of some imprinted genes in murine oocytes. *Biology of Reproduction* **88** 117. (doi:10.1095/biolreprod.112.105981)
- Ge ZJ, Zhang CL, Schatten H & Sun QY 2014a Maternal diabetes mellitus and the origin of non-communicable diseases in offspring: the role of epigenetics. *Biology of Reproduction* **90** 139. (doi:10.1095/biolreprod.114.118141)
- Ge ZJ, Luo SM, Lin F, Liang QX, Huang L, Wei YC, Hou Y, Han ZM, Schatten H & Sun QY 2014b DNA methylation in oocytes and liver of female mice and their offspring: effects of high-fat-diet-induced obesity. *Environmental Health Perspectives* **122** 159–164. (doi:10.1289/ehp.1307047)
- Gentilini D, Mari D, Castaldi D, Remondini D, Ogliari G, Ostan R, Bucci L, Sirchia SM, Tabano S, Cavagnini F *et al.* 2013 Role of epigenetics in human aging and longevity: genome-wide DNA methylation profile in centenarians and centenarians' offspring. *Age* **35** 1961–1973. (doi:10.1007/s11357-012-9463-1)
- Gilchrist RB, Lane M & Thompson JG 2008 Oocyte-secreted factors: regulators of cumulus cell function and oocyte quality. *Human Reproduction Update* **14** 159–177. (doi:10.1093/humupd/dmm040)
- Goldberg AD, Allis CD & Bernstein E 2007 Epigenetics: a landscape takes shape. *Cell* **128** 635–638. (doi:10.1016/j.cell.2007.02.006)
- Grondahl ML, Yding Andersen C, Bogstad J, Nielsen FC, Meinertz H & Borup R 2010 Gene expression profiles of single human mature oocytes in relation to age. *Human Reproduction* **25** 957–968. (doi:10.1093/humrep/deq014)
- Gu L, Wang Q & Sun QY 2010 Histone modifications during mammalian oocyte maturation: dynamics, regulation and functions. *Cell Cycle* **9** 1942–1950. (doi:10.4161/cc.9.10.11599)
- Gu TP, Guo F, Yang H, Wu HP, Xu GF, Liu W, Xie ZG, Shi L, He X, Jin SG *et al.* 2011 The role of Tet3 DNA dioxygenase in epigenetic reprogramming by oocytes. *Nature* **477** 606–610. (doi:10.1038/nature10443)
- Guglielmino MR, Santonocito M, Vento M, Ragusa M, Barbagallo D, Borzi P, Casciano I, Banelli B, Barbieri O, Astigiano S *et al.* 2011 Tap73 is downregulated in oocytes from women of advanced reproductive age. *Cell Cycle* **10** 3253–3256. (doi:10.4161/cc.10.19.17585)
- Guo H, Zhu P, Yan L, Li R, Hu B, Lian Y, Yan J, Ren X, Lin S, Li J *et al.* 2014 The DNA methylation landscape of human early embryos. *Nature* **511** 606–610. (doi:10.1038/nature13544)
- Hales BF, Grenier L, Lalancette C & Robaire B 2011 Epigenetic programming: from gametes to blastocyst. *Birth Defects Research. Part A, Clinical and Molecular Teratology* **91** 652–665. (doi:10.1002/bdra.20781)
- Hamatani T, Falco G, Carter MG, Akutsu H, Stagg CA, Sharov AA, Dudekula DB, VanBuren V & Ko MS 2004 Age-associated alteration of gene expression patterns in mouse oocytes. *Human Molecular Genetics* **13** 2263–2278. (doi:10.1093/hmg/ddh241)
- Han Y, Han D, Yan Z, Boyd-Kirkup JD, Green CD, Khaitovich P & Han JD 2012 Stress-associated H3K4 methylation accumulates during postnatal development and aging of rhesus macaque brain. *Aging Cell* **11** 1055–1064. (doi:10.1111/accel.12007)
- Handyside AH, Montag M, Magli MC, Repping S, Harper J, Schmutzler A, Vesela K, Gianaroli L & Geraedts J 2012 Multiple meiotic errors caused by predivision of chromatids in women of advanced maternal age undergoing *in vitro* fertilisation. *European Journal of Human Genetics* **20** 742–747. (doi:10.1038/ejhg.2011.272)
- Hassold T, Hall H & Hunt P 2007 The origin of human aneuploidy: where we have been, where we are going. *Human Molecular Genetics* **16** R203–R208. (doi:10.1093/hmg/ddm243)
- Henderson SA & Edwards RG 1968 Chiasma frequency and maternal age in mammals. *Nature* **218** 22–28. (doi:10.1038/218022a0)
- Herskind AM, McGue M, Holm NV, Sorensen TI, Harvald B & Vaupel JW 1996 The heritability of human longevity: a population-based study of 2872 Danish twin pairs born 1870–1900. *Human Genetics* **97** 319–323. (doi:10.1007/BF02185763)
- Heyn H, Li N, Ferreira HJ, Moran S, Pisano DG, Gomez A, Diez J, Sanchez-Mut JV, Setien F, Carmona FJ *et al.* 2012 Distinct DNA methylomes of newborns and centenarians. *PNAS* **109** 10522–10527. (doi:10.1073/pnas.1120658109)
- Hirasawa R, Chiba H, Kaneda M, Tajima S, Li E, Jaenisch R & Sasaki H 2008 Maternal and zygotic Dnmt1 are necessary and sufficient for the maintenance of DNA methylation imprints during preimplantation development. *Genes and Development* **22** 1607–1616. (doi:10.1101/gad.1667008)
- van den Hove DL, Chouliaras L & Rutten BP 2012 The role of 5-hydroxymethylcytosine in aging and Alzheimer's disease: current status and prospects for future studies. *Current Alzheimer Research* **9** 545–549.
- Howell CY, Bestor TH, Ding F, Latham KE, Mertineit C, Trasler JM & Chaillet JR 2001 Genomic imprinting disrupted by a maternal effect mutation in the Dnmt1 gene. *Cell* **104** 829–838. (doi:10.1016/S0092-8674(01)00280-X)
- Huang JC, Yan LY, Lei ZL, Miao YL, Shi LH, Yang JW, Wang Q, Ouyang YC, Sun QY & Chen DY 2007 Changes in histone acetylation during postovulatory aging of mouse oocyte. *Biology of Reproduction* **77** 666–670. (doi:10.1095/biolreprod.107.062703)
- Igarashi K & Katoh Y 2013 Metabolic aspects of epigenome: coupling of S-adenosylmethionine synthesis and gene regulation on chromatin by SAMIT module. *Sub-Cellular Biochemistry* **61** 105–118. (doi:10.1007/978-94-007-4525-4_5)
- Imamura T, Kerjean A, Heams T, Kupiec JJ, Thenevin C & Paldi A 2005 Dynamic CpG and non-CpG methylation of the Peg1/Mest gene in the mouse oocyte and preimplantation embryo. *Journal of Biological Chemistry* **280** 20171–20175. (doi:10.1074/jbc.M501749200)
- Inukai S, de Lencastre A, Turner M & Slack F 2012 Novel microRNAs differentially expressed during aging in the mouse brain. *PLoS ONE* **7** e40028. (doi:10.1371/journal.pone.0040028)
- Iwata H, Goto H, Tanaka H, Sakaguchi Y, Kimura K, Kuwayama T & Monji Y 2011 Effect of maternal age on mitochondrial DNA copy number, ATP content and IVF outcome of bovine oocytes. *Reproduction, Fertility, and Development* **23** 424–432. (doi:10.1071/RD10133)
- Jacob RA, Gretz DM, Taylor PC, James SJ, Pogribny IP, Miller BJ, Henning SM & Swendseid ME 1998 Moderate folate depletion increases plasma homocysteine and decreases lymphocyte DNA methylation in postmenopausal women. *Journal of Nutrition* **128** 1204–1212.
- Jacques PF, Bostom AG, Wilson PW, Rich S, Rosenberg IH & Selhub J 2001 Determinants of plasma total homocysteine concentration in the Framingham Offspring cohort. *American Journal of Clinical Nutrition* **73** 613–621.
- Jeseta M, Petr J, Krejcova T, Chmelikova E & Jilek F 2008 *In vitro* ageing of pig oocytes: effects of the histone deacetylase inhibitor trichostatin A. *Zygote* **16** 145–152. (doi:10.1017/S0967199408004668)
- Jessberger R 2012 Age-related aneuploidy through cohesion exhaustion. *EMBO Reports* **13** 539–546. (doi:10.1038/embor.2012.54)

- Jones KT & Lane SI 2013 Molecular causes of aneuploidy in mammalian eggs. *Development* **140** 3719–3730. (doi:10.1242/dev.090589)
- Kaneda M, Tang F, O'Carroll D, Lao K & Surani MA 2009 Essential role for Argonaute2 protein in mouse oogenesis. *Epigenetics & Chromatin* **2** 9. (doi:10.1186/1756-8935-2-9)
- Kemkes-Grottenthaler A 2004 Parental effects on offspring longevity – evidence from 17th to 19th century reproductive histories. *Annals of Human Biology* **31** 139–158. (doi:10.1080/03014460410001663407)
- Kim JM, Liu H, Tazaki M, Nagata M & Aoki F 2003 Changes in histone acetylation during mouse oocyte meiosis. *Journal of Cell Biology* **162** 37–46. (doi:10.1083/jcb.200303047)
- Kim M, Long TI, Arakawa K, Wang R, Yu MC & Laird PW 2010 DNA methylation as a biomarker for cardiovascular disease risk. *PLoS ONE* **5** e9692. (doi:10.1371/journal.pone.0009692)
- Kosubek A, Klein-Hitpass L, Rademacher K, Horsthemke B & Ryffel GU 2010 Aging of *Xenopus tropicalis* eggs leads to deadenylation of a specific set of maternal mRNAs and loss of developmental potential. *PLoS ONE* **5** e13532. (doi:10.1371/journal.pone.0013532)
- Kurihara Y, Kawamura Y, Uchijima Y, Amamo T, Kobayashi H, Asano T & Kurihara H 2008 Maintenance of genomic methylation patterns during preimplantation development requires the somatic form of DNA methyltransferase 1. *Developmental Biology* **313** 335–346. (doi:10.1016/j.ydbio.2007.10.033)
- Lee AR, Kishigami S, Amano T, Matsumoto K, Wakayama T & Hosoi Y 2013 Nicotinamide: a class III HDACi delays *in vitro* aging of mouse oocytes. *Journal of Reproduction and Development* **59** 238–244. (doi:10.1262/jrd.2012-171)
- Li Y, Daniel M & Tollefsbol TO 2011 Epigenetic regulation of caloric restriction in aging. *BMC Medicine* **9** 98. (doi:10.1186/1741-7015-9-98)
- Li L, Zhang T, Qin XS, Ge W, Ma HG, Sun LL, Hou ZM, Chen H, Chen P, Qin GQ *et al.* 2014 Exposure to diethylhexyl phthalate (DEHP) results in a heritable modification of imprint genes DNA methylation in mouse oocytes. *Molecular Biology Reports* **41** 1227–1235. (doi:10.1007/s11033-013-2967-7)
- Liang XW, Zhu JQ, Miao YL, Liu JH, Wei L, Lu SS, Hou Y, Schatten H, Lu KH & Sun QY 2008 Loss of methylation imprint of Snrpn in postovulatory aging mouse oocyte. *Biochemical and Biophysical Research Communications* **371** 16–21. (doi:10.1016/j.bbrc.2008.03.105)
- Liang XW, Ge ZJ, Guo L, Luo SM, Han ZM, Schatten H & Sun QY 2011 Effect of postovulatory oocyte aging on DNA methylation imprinting acquisition in offspring oocytes. *Fertility and Sterility* **96** 1479–1484. (doi:10.1016/j.fertnstert.2011.09.022)
- Liang X, Ma J, Schatten H & Sun Q 2012 Epigenetic changes associated with oocyte aging. *Science China. Life Sciences* **55** 670–676. (doi:10.1007/s11427-012-4354-3)
- Lim AS & Tsakok MF 1997 Age-related decline in fertility: a link to degenerative oocytes? *Fertility and Sterility* **68** 265–271. (doi:10.1016/S0015-0282(97)81513-0)
- Lister LM, Kouznetsova A, Hyslop LA, Kalleas D, Pace SL, Barel JC, Nathan A, Floros V, Adelfalk C, Watanabe Y *et al.* 2010 Age-related meiotic segregation errors in mammalian oocytes are preceded by depletion of cohesin and Sgo2. *Current Biology* **20** 1511–1521. (doi:10.1016/j.cub.2010.08.023)
- Liu L, Wylie RC, Andrews LG & Tollefsbol TO 2003 Aging, cancer and nutrition: the DNA methylation connection. *Mechanisms of Ageing and Development* **124** 989–998. (doi:10.1016/j.mad.2003.08.001)
- Liu L, van Groen T, Kadish I & Tollefsbol TO 2009a DNA methylation impacts on learning and memory in aging. *Neurobiology of Aging* **30** 549–560. (doi:10.1016/j.neurobiolaging.2007.07.020)
- Liu N, Wu YG, Lan GC, Sui HS, Ge L, Wang JZ, Liu Y, Qiao TW & Tan JH 2009b Pyruvate prevents aging of mouse oocytes. *Reproduction* **138** 223–234. (doi:10.1530/REP-09-0122)
- Liu L, van Groen T, Kadish I, Li Y, Wang D, James SR, Karpf AR & Tollefsbol TO 2011 Insufficient DNA methylation affects healthy aging and promotes age-related health problems. *Clinical Epigenetics* **2** 349–360. (doi:10.1007/s13148-011-0042-6)
- Lopes FL, Fortier AL, Darricarrere N, Chan D, Arnold DR & Trasler JM 2009 Reproductive and epigenetic outcomes associated with aging mouse oocytes. *Human Molecular Genetics* **18** 2032–2044. (doi:10.1093/hmg/ddp127)
- Lovatt D, Ruble BK, Lee J, Dueck H, Kim TK, Fisher S, Francis C, Spaethling JM, Wolf JA, Grady MS *et al.* 2014 Transcriptome *in vivo* analysis (TIVA) of spatially defined single cells in live tissue. *Nature Methods* **11** 190–196. (doi:10.1038/nmeth.2804)
- Ma JY, Liang XW, Schatten H & Sun QY 2012 Active DNA demethylation in mammalian preimplantation embryos: new insights and new perspectives. *Molecular Human Reproduction* **18** 333–340. (doi:10.1093/molehr/gas014)
- Malini SS & Ramachandra NB 2006 Influence of advanced age of maternal grandmothers on Down syndrome. *BMC Medical Genetics* **7** 4. (doi:10.1186/1471-2350-7-4)
- Manosalva I & Gonzalez A 2009 Aging alters histone H4 acetylation and CDC2A in mouse germinal vesicle stage oocytes. *Biology of Reproduction* **81** 1164–1171. (doi:10.1095/biolreprod.109.078386)
- Manosalva I & Gonzalez A 2010 Aging changes the chromatin configuration and histone methylation of mouse oocytes at germinal vesicle stage. *Theriogenology* **74** 1539–1547. (doi:10.1016/j.theriogenology.2010.06.024)
- McCauley BS & Dang W 2015 Histone methylation and aging: lessons learned from model systems. *Biochimica et Biophysica Acta*. In press. (doi:10.1016/j.bbagr.2014.05.008)
- Menken J, Trussell J & Larsen U 1986 Age and infertility. *Science* **233** 1389–1394. (doi:10.1126/science.3755843)
- Mertineit C, Yoder JA, Taketo T, Laird DW, Trasler JM & Bestor TH 1998 Sex-specific exons control DNA methyltransferase in mammalian germ cells. *Development* **125** 889–897.
- Messerschmidt DM, de Vries W, Ito M, Solter D, Ferguson-Smith A & Knowles BB 2012 Trim28 is required for epigenetic stability during mouse oocyte to embryo transition. *Science* **335** 1499–1502. (doi:10.1126/science.1216154)
- Miao YL, Kikuchi K, Sun QY & Schatten H 2009 Oocyte aging: cellular and molecular changes, developmental potential and reversal possibility. *Human Reproduction Update* **15** 573–585. (doi:10.1093/humupd/dmp014)
- Mitchell BD, Hsueh WC, King TM, Pollin TI, Sorkin J, Agarwala R, Schaffer AA & Shuldiner AR 2001 Heritability of life span in the Old Order Amish. *American Journal of Medical Genetics* **102** 346–352. (doi:10.1002/ajmg.1483)
- Moore LE, Pfeiffer RM, Poscablo C, Real FX, Kogevinas M, Silverman D, Garcia-Closas R, Chanock S, Tardon A, Serra C *et al.* 2008 Genomic DNA hypomethylation as a biomarker for bladder cancer susceptibility in the Spanish Bladder Cancer Study: a case-control study. *Lancet Oncology* **9** 359–366. (doi:10.1016/S1470-2045(08)70038-X)
- Murchison EP, Stein P, Xuan Z, Pan H, Zhang MQ, Schultz RM & Hannon GJ 2007 Critical roles for Dicer in the female germline. *Genes and Development* **21** 682–693. (doi:10.1101/gad.1521307)
- Navot D, Bergh PA, Williams MA, Garrisi GJ, Guzman I, Sandler B & Grunfeld L 1991 Poor oocyte quality rather than implantation failure as a cause of age-related decline in female fertility. *Lancet* **337** 1375–1377. (doi:10.1016/0140-6736(91)93060-M)
- Olsen NJ & Kovacs WJ 1996 Gonadal steroids and immunity. *Endocrine Reviews* **17** 369–384.
- Pan H, Ma P, Zhu W & Schultz RM 2008 Age-associated increase in aneuploidy and changes in gene expression in mouse eggs. *Developmental Biology* **316** 397–407. (doi:10.1016/j.ydbio.2008.01.048)
- Petronczki M, Siomos MF & Nasmyth K 2003 Un ménage à quatre: the molecular biology of chromosome segregation in meiosis. *Cell* **112** 423–440. (doi:10.1016/S0092-8674(03)00083-7)
- Pincus Z, Smith-Vikos T & Slack FJ 2011 MicroRNA predictors of longevity in *Caenorhabditis elegans*. *PLoS Genetics* **7** e1002306. (doi:10.1371/journal.pgen.1002306)
- Poulsen P, Esteller M, Vaag A & Fraga MF 2007 The epigenetic basis of twin discordance in age-related diseases. *Pediatric Research* **61** 38R–42R. (doi:10.1203/pdr.0b013e31803c7b98)
- Qiao J, Wang ZB, Feng HL, Miao YL, Wang Q, Yu Y, Wei YC, Yan J, Wang WH, Shen W *et al.* 2014 The root of reduced fertility in aged women and possible therapeutic options: current status and future prospects. *Molecular Aspects of Medicine* **38** 54–85. (doi:10.1016/j.mam.2013.06.001)
- Rambags BP, van Boxtel DC, Tharasani T, Lenstra JA, Colenbrander B & Stout TA 2014 Advancing maternal age predisposes to mitochondrial damage and loss during maturation of equine oocytes *in vitro*. *Theriogenology* **81** 959–965. (doi:10.1016/j.theriogenology.2014.01.020)

- Rampersaud GC, Kauwell GP, Hutson AD, Cerda JJ & Bailey LB 2000 Genomic DNA methylation decreases in response to moderate folate depletion in elderly women. *American Journal of Clinical Nutrition* **72** 998–1003.
- Reddy KC & Villeneuve AM 2004 *C. elegans* HIM-17 links chromatin modification and competence for initiation of meiotic recombination. *Cell* **118** 439–452. (doi:10.1016/j.cell.2004.07.026)
- Revenkova E, Eijpe M, Heyting C, Hodges CA, Hunt PA, Liebe B, Scherthan H & Jessberger R 2004 Cohesin SMC1 β is required for meiotic chromosome dynamics, sister chromatid cohesion and DNA recombination. *Nature Cell Biology* **6** 555–562. (doi:10.1038/ncb1135)
- Rocca WA, van Duijn CM, Clayton D, Chandra V, Fratiglioni L, Graves AB, Heyman A, Jorm AF, Kokmen E, Kondo K *et al.* 1991 Maternal age and Alzheimer's disease: a collaborative re-analysis of case-control studies. EURODEM Risk Factors Research Group. *International Journal of Epidemiology* **20** (Suppl 2) S21–S27. (doi:10.1093/ije/20.Supplement_2.S21)
- Romeu A, Muasher SJ, Acosta AA, Veeck LL, Diaz J, Jones GS, Jones HW Jr & Rosenwaks Z 1987 Results of *in vitro* fertilization attempts in women 40 years of age and older: the Norfolk experience. *Fertility and Sterility* **47** 130–136.
- Roth LW, McCallie B, Alvero R, Schoolcraft WB, Minjarez D & Katz-Jaffe MG 2014 Altered microRNA and gene expression in the follicular fluid of women with polycystic ovary syndrome. *Journal of Assisted Reproduction and Genetics* **31** 355–362. (doi:10.1007/s10815-013-0161-4)
- Santonocito M, Guglielmino MR, Vento M, Ragusa M, Barbagallo D, Borzi P, Casciano I, Scollo P, Romani M, Tatone C *et al.* 2013 The apoptotic transcriptome of the human MII oocyte: characterization and age-related changes. *Apoptosis* **18** 201–211. (doi:10.1007/s10495-012-0783-5)
- Sato A, Otsu E, Negishi H, Utsunomiya T & Arima T 2007 Aberrant DNA methylation of imprinted loci in superovulated oocytes. *Human Reproduction* **22** 26–35. (doi:10.1093/humrep/del316)
- Schwarz JM, Nugent BM & McCarthy MM 2010 Developmental and hormone-induced epigenetic changes to estrogen and progesterone receptor genes in brain are dynamic across the life span. *Endocrinology* **151** 4871–4881. (doi:10.1210/en.2010-0142)
- Schwarzer C, Siatkowski M, Pfeiffer MJ, Baeumer N, Drexler HC, Wang B, Fuellen G & Boiani M 2014 Maternal age effect on mouse oocytes: new biological insight from proteomic analysis. *Reproduction* **148** 55–72. (doi:10.1530/REP-14-0126)
- Sekhon LH, Gerber RS, Rebarber A, Saltzman DH, Klauser CK, Gupta S & Fox NS 2014 Effect of oocyte donation on pregnancy outcomes in *in vitro* fertilization twin gestations. *Fertility and Sterility* **101** 1326–1330. (doi:10.1016/j.fertnstert.2014.01.055)
- Selemani M, Mwanyangala MA, Mrema S, Shamte A, Kajungu D, Mkopi A, Mahande MJ & Nathan R 2014 The effect of mother's age and other related factors on neonatal survival associated with first and second birth in rural, Tanzania: evidence from Ifakara health and demographic surveillance system in rural Tanzania. *BMC Pregnancy and Childbirth* **14** 240. (doi:10.1186/1471-2393-14-240)
- Selesniemi K, Lee HJ, Muhlhauser A & Tilly JL 2011 Prevention of maternal aging-associated oocyte aneuploidy and meiotic spindle defects in mice by dietary and genetic strategies. *PNAS* **108** 12319–12324. (doi:10.1073/pnas.1018793108)
- Shao GB, Wang J, Zhang LP, Wu CY, Jin J, Sang JR, Lu HY, Gong AH, Du FY & Peng WX 2015 Aging alters histone H3 lysine 4 methylation in mouse germinal vesicle stage oocytes. *Reproduction, Fertility, and Development*. In press. (doi:10.1071/RD13293)
- Sher G, Keskinetepe L, Keskinetepe M, Ginsburg M, Maassarani G, Yakut T, Baltaci V, Kotze D & Unsal E 2007 Oocyte karyotyping by comparative genomic hybridization [correction of hybridization] provides a highly reliable method for selecting "competent" embryos, markedly improving *in vitro* fertilization outcome: a multiphase study. *Fertility and Sterility* **87** 1033–1040. (doi:10.1016/j.fertnstert.2006.08.108)
- Shimoda N, Izawa T, Yoshizawa A, Yokoi H, Kikuchi Y & Hashimoto N 2014 Decrease in cytosine methylation at CpG island shores and increase in DNA fragmentation during zebrafish aging. *Age* **36** 103–115. (doi:10.1007/s11357-013-9548-5)
- da Silveira JC, Veeramachaneni DN, Winger QA, Carnevale EM & Bouma GJ 2012 Cell-secreted vesicles in equine ovarian follicular fluid contain miRNAs and proteins: a possible new form of cell communication within the ovarian follicle. *Biology of Reproduction* **86** 71. (doi:10.1095/biolreprod.111.093252)
- Simsek-Duran F, Li F, Ford W, Swanson RJ, Jones HW Jr & Castora FJ 2013 Age-associated metabolic and morphologic changes in mitochondria of individual mouse and hamster oocytes. *PLoS ONE* **8** e64955. (doi:10.1371/journal.pone.0064955)
- Smallwood SA, Lee HJ, Angermueller C, Krueger F, Saadeh H, Peat J, Andrews SR, Stegle O, Reik W & Kelsey G 2014 Single-cell genome-wide bisulfite sequencing for assessing epigenetic heterogeneity. *Nature Methods* **11** 817–820. (doi:10.1038/nmeth.3035)
- Smits LJ, Zielhuis GA, Jongbloet PH & Van Poppel FW 2002 Mother's age and daughter's fecundity. An epidemiological analysis of late 19th to early 20th century family reconstitutions. *International Journal of Epidemiology* **31** 349–358. (doi:10.1093/ije/31.2.349)
- Sohel MM, Hoelker M, Noferesti SS, Salilew-Wondim D, Tholen E, Looft C, Rings F, Uddin MJ, Spencer TE, Schellander K *et al.* 2013 Exosomal and non-exosomal transport of extra-cellular microRNAs in follicular fluid: implications for bovine oocyte developmental competence. *PLoS ONE* **8** e78505. (doi:10.1371/journal.pone.0078505)
- Steegers-Theunissen RP, Twigt J, Pestinger V & Sinclair KD 2013 The periconceptual period, reproduction and long-term health of offspring: the importance of one-carbon metabolism. *Human Reproduction Update* **19** 640–655. (doi:10.1093/humupd/dmt041)
- Steuerwald NM, Bermudez MG, Wells D, Munne S & Cohen J 2007 Maternal age-related differential global expression profiles observed in human oocytes. *Reproductive Biomedicine Online* **14** 700–708. (doi:10.1016/S1472-6483(10)60671-2)
- Suh N & Billeloch R 2011 Small RNAs in early mammalian development: from gametes to gastrulation. *Development* **138** 1653–1661. (doi:10.1242/dev.056234)
- Suh N, Baehner L, Moltzahn F, Melton C, Shenoy A, Chen J & Billeloch R 2010 MicroRNA function is globally suppressed in mouse oocytes and early embryos. *Current Biology* **20** 271–277. (doi:10.1016/j.cub.2009.12.044)
- Suo L, Meng QG, Pei Y, Yan CL, Fu XW, Bunch TD & Zhu SE 2010 Changes in acetylation on lysine 12 of histone H4 (acH4K12) of murine oocytes during maternal aging may affect fertilization and subsequent embryo development. *Fertility and Sterility* **93** 945–951. (doi:10.1016/j.fertnstert.2008.12.128)
- Svoboda P & Flenr M 2010 The role of miRNAs and endogenous siRNAs in maternal-to-zygotic reprogramming and the establishment of pluripotency. *EMBO Reports* **11** 590–597. (doi:10.1038/embor.2010.102)
- Tahiliani M, Koh KP, Shen Y, Pastor WA, Bandukwala H, Brudno Y, Agarwal S, Iyer LM, Liu DR, Aravind L *et al.* 2009 Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* **324** 930–935. (doi:10.1126/science.1170116)
- Takeo S, Goto H, Kuwayama T, Monji Y & Iwata H 2013a Effect of maternal age on the ratio of cleavage and mitochondrial DNA copy number in early developmental stage bovine oocytes. *Journal of Reproduction and Development* **59** 174–179. (doi:10.1262/jrd.2012-148)
- Takeo S, Kawahara-Miki R, Goto H, Cao F, Kimura K, Monji Y, Kuwayama T & Iwata H 2013b Age-associated changes in gene expression and developmental competence of bovine oocytes, and a possible countermeasure against age-associated events. *Molecular Reproduction and Development* **80** 508–521. (doi:10.1002/mrd.22187)
- Tam OH, Aravin AA, Stein P, Girard A, Murchison EP, Cheloufi S, Hodges E, Anger M, Sachidanandam R, Schultz RM *et al.* 2008 Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature* **453** 534–538. (doi:10.1038/nature06904)
- Tarin JJ, Brines J & Cano A 1998a Long-term effects of delayed parenthood. *Human Reproduction* **13** 2371–2376. (doi:10.1093/humrep/13.9.2371)
- Tarin JJ, Ten J, Vendrell FJ & Cano A 1998b Dithiothreitol prevents age-associated decrease in oocyte/conceptus viability *in vitro*. *Human Reproduction* **13** 381–386. (doi:10.1093/humrep/13.2.381)
- Tarin JJ, Perez-Albala S, Aguilar A, Minarro J, Hermenegildo C & Cano A 1999 Long-term effects of postovulatory aging of mouse oocytes on offspring: a two-generational study. *Biology of Reproduction* **61** 1347–1355. (doi:10.1095/biolreprod61.5.1347)
- Tarin JJ, Perez-Albala S, Perez-Hoyos S & Cano A 2002 Postovulatory aging of oocytes decreases reproductive fitness and longevity of offspring. *Biology of Reproduction* **66** 495–499. (doi:10.1095/biolreprod66.2.495)
- Tatone C & Amicarelli F 2013 The aging ovary – the poor granulosa cells. *Fertility and Sterility* **99** 12–17. (doi:10.1016/j.fertnstert.2012.11.029)

- Tatone C, Amicarelli F, Carbone MC, Monteleone P, Caserta D, Marci R, Artini PG, Piomboni P & Focarelli R 2008 Cellular and molecular aspects of ovarian follicle ageing. *Human Reproduction Update* **14** 131–142. (doi:10.1093/humupd/dmm048)
- Tatone C, Eichenlaub-Ritter U & Amicarelli F 2014 Dicarbonyl stress and glyoxalases in ovarian function. *Biochemical Society Transactions* **42** 433–438. (doi:10.1042/BST20140023)
- Tomizawa S, Nowacka-Woszek J & Kelsey G 2012 DNA methylation establishment during oocyte growth: mechanisms and significance. *International Journal of Developmental Biology* **56** 867–875. (doi:10.1387/ijdb.120152gk)
- Torner H, Brussow KP, Alm H, Ratky J, Pohland R, Tuchscherer A & Kanitz W 2004 Mitochondrial aggregation patterns and activity in porcine oocytes and apoptosis in surrounding cumulus cells depends on the stage of pre-ovulatory maturation. *Theriogenology* **61** 1675–1689. (doi:10.1016/j.theriogenology.2003.09.013)
- Tripurani SK, Wee G, Lee KB, Smith GW, Wang L & Jianboyao Y 2013 MicroRNA-212 post-transcriptionally regulates oocyte-specific basic-helix-loop-helix transcription factor, factor in the germline alpha (FIGLA), during bovine early embryogenesis. *PLoS ONE* **8** e76114. (doi:10.1371/journal.pone.0076114)
- Truesdell SS, Mortensen RD, Seo M, Schroeder JC, Lee JH, LeTonqueze O & Vasudevan S 2012 MicroRNA-mediated mRNA translation activation in quiescent cells and oocytes involves recruitment of a nuclear microRNP. *Scientific Reports* **2** 842. (doi:10.1038/srep00842)
- Tsutsumi M, Fujiwara R, Nishizawa H, Ito M, Kogo H, Inagaki H, Ohye T, Kato T, Fujii T & Kurahashi H 2014 Age-related decrease of meiotic cohesins in human oocytes. *PLoS ONE* **9** e96710. (doi:10.1371/journal.pone.0096710)
- te Velde ER & Pearson PL 2002 The variability of female reproductive ageing. *Human Reproduction Update* **8** 141–154. (doi:10.1093/humupd/8.2.141)
- Wang YA, Farquhar C & Sullivan EA 2012 Donor age is a major determinant of success of oocyte donation/recipient programme. *Human Reproduction* **27** 118–125. (doi:10.1093/humrep/der359)
- Warburton D, Kline J, Stein Z, Hutzler M, Chin A & Hassold T 1987 Does the karyotype of a spontaneous abortion predict the karyotype of a subsequent abortion? Evidence from 273 women with two karyotyped spontaneous abortions. *American Journal of Human Genetics* **41** 465–483.
- Wellen KE, Hatzivassiliou G, Sachdeva UM, Bui TV, Cross JR & Thompson CB 2009 ATP-citrate lyase links cellular metabolism to histone acetylation. *Science* **324** 1076–1080. (doi:10.1126/science.1164097)
- Wilson VL & Jones PA 1983 DNA methylation decreases in aging but not in immortal cells. *Science* **220** 1055–1057. (doi:10.1126/science.6844925)
- Wnuk M, Lewinska A, Gurgul A, Zabeł T, Potocki L, Oklejewicz B, Bugno-Poniewierska M, Węgrzyn M & Słota E 2014 Changes in DNA methylation patterns and repetitive sequences in blood lymphocytes of aged horses. *Age* **36** 31–48. (doi:10.1007/s11357-013-9541-z)
- Wood JC & Helfand SL 2013 Chromatin structure and transposable elements in organismal aging. *Frontiers in Genetics* **4** 274. (doi:10.3389/fgene.2013.00274)
- Yamaguchi S, Hong K, Liu R, Shen L, Inoue A, Diep D, Zhang K & Zhang Y 2012 Tet1 controls meiosis by regulating meiotic gene expression. *Nature* **492** 443–447. (doi:10.1038/nature11709)
- Yang S, Wang S, Luo A, Ding T, Lai Z, Shen W, Ma X, Cao C, Shi L, Jiang J *et al.* 2013 Expression patterns and regulatory functions of microRNAs during the initiation of primordial follicle development in the neonatal mouse ovary. *Biology of Reproduction* **89** 126. (doi:10.1095/biolreprod.113.107730)
- Ye Y, Li D, Ouyang D, Deng L, Zhang Y, Ma Y & Li Y 2014 MicroRNA expression in the aging mouse thymus. *Gene* **547** 218–225. (doi:10.1016/j.gene.2014.06.039)
- Yue MX, Fu XW, Zhou GB, Hou YP, Du M, Wang L & Zhu SE 2012 Abnormal DNA methylation in oocytes could be associated with a decrease in reproductive potential in old mice. *Journal of Assisted Reproduction and Genetics* **29** 643–650. (doi:10.1007/s10815-012-9780-4)
- Zhang L, Hou X, Ma R, Moley K, Schedl T & Wang Q 2014a Sirt2 functions in spindle organization and chromosome alignment in mouse oocyte meiosis. *FASEB Journal* **28** 1435–1445. (doi:10.1096/fj.13-244111)
- Zhang H, Jiang X, Zhang Y, Xu B, Hua J, Ma T, Zheng W, Sun R, Shen W, Cooke HJ *et al.* 2014b microRNA 376a regulates follicle assembly by targeting PcnA in fetal and neonatal mouse ovaries. *Reproduction* **148** 43–54. (doi:10.1530/REP-13-0508)

Received 6 May 2014

First decision 15 July 2014

Revised manuscript received 7 November 2014

Accepted 12 November 2014