Epigenetic dynamics during preimplantation development

Chelsea Marcho*, Wei Cui* and Jesse Mager

Department of Veterinary and Animal Science, University of Massachusetts Amherst, 661 North Pleasant Street, Amherst, Massachusetts 01003, USA

Correspondence should be addressed to J Mager; Email: jmager@vasci.umass.edu

*(C Marcho and W Cui contributed equally to this work)

Abstract

Successful mammalian development requires descendants of single-cell zygotes to differentiate into diverse cell types even though they contain the same genetic material. Preimplantation dynamics are first driven by the necessity of reprogramming haploid parental epigenomes to reach a totipotent state. This process requires extensive erasure of epigenetic marks shortly after fertilization. During the few short days after formation of the zygote, epigenetic programs are established and are essential for the first lineage decisions and differentiation. Here we review the current understanding of DNA methylation and histone modification dynamics responsible for these early changes during mammalian preimplantation development. In particular, we highlight insights that have been gained through next-generation sequencing technologies comparing human embryos to other models as well as the recent discoveries of active DNA demethylation mechanisms at play during preimplantation.

Reproduction (2015) **150** R109–R120

From zygote to blastocyst

Mammalian preimplantation development is a time of dynamic change in which the fertilized egg undergoes cleavage divisions developing into a morula and then a blastocyst with the first two distinct cell lineages (inner cell mass (ICM) and trophectoderm (TE)). This developmental period is characterized by three major transitions, each of which entails pronounced changes in the pattern of gene expression. The first transition is the maternal-to-zygotic transition (MZT), which serves three functions: i) to destroy oocyte-specific transcripts (e.g., H1oo (Tanaka *et al.* 2001)), ii) to replace maternal transcripts that are common to the oocyte and early embryo with zygotic transcripts and iii) to facilitate the reprogramming of the early embryo by generating novel transcripts that are not expressed in the oocyte (Latham *et al.* 1991). In mice, zygotic gene activation initiates during the 1-cell stage, and is clearly evident by the 2-cell stage (Latham *et al.* 1991, Schultz *et al.* 1993). Coincident with genome activation is the implementation of a chromatin-based transcriptionally-repressive state (Nothias *et al.* 1995, Schultz 2002) and more efficient use of TATA-less promoters (Majumder & DePamphilis 1994), which are likely to play a major role in establishing the appropriate pattern of gene expression required for successful development.

The second developmental transition is compaction, which occurs during the 8-cell stage, when the first morphological differentiation occurs due to adhesive interactions between the blastomeres generating a tightly organized and less distinct mass of cells (Fleming *et al.* 2001). Accompanying compaction are pronounced biochemical changes through which blastomeres acquire characteristics resembling somatic cells, reflected in such features as ion transport, metabolism, cellular architecture and gene expression pattern (Fleming *et al.* 2001, Kidder & Winterhager 2001). Following compaction, cleavage divisions allocate cells to the inside of the developing morula. These inner cells are set aside between the 8-cell and 16-cell stage, and then again between the 16-cell stage and the 32-cell stage (Pedersen *et al.* 1986). The inner cells of the morula give rise to the ICM cells from which the embryo proper is derived, whereas the outer cells differentiate exclusively into the TE, which gives rise to extraembryonic tissues (Yamanaka *et al.* 2006). The TE is a fluid-transporting epithelium that is responsible for forming the blastocoel cavity and is essential for continued development and differentiation of the ICM (Biggers *et al.* 1988, Watson *et al.* 1990). Distinct differentiation first occurs in the blastocyst and is characterized by differences in gene expression between the ICM and

TE cells (Nichols & Gardner 1984, Pesce & Scholer 2001). Additionally, by the time of implantation the primitive endoderm has differentiated from the ICM/epiblast and resides as a single-cell layer on the blastocoel cavity side of the ICM/EPI (reviewed in Schrode *et al.* (2013)).

These dynamic morphological, cellular and molecular events are driven by gene expression changes facilitated by epigenetic phenomenon, including DNA methylation and histone modifications at sites throughout the genome. Below we review the current understanding of the mechanisms responsible for regulation of epigenetic programming and re-programming that occur during mammalian preimplantation.

DNA methylation dynamics in the preimplantation mouse embryo

In mammalian cells, the predominant form of DNA methylation occurs at CpG dinucleotides. Throughout the genome, non-promoter associated CpGs are generally found methylated. However, the majority of protein coding genes have regions of high-density CpG dinucleotides termed CpG islands. In most cell types the methylation status at these promoter associated CpG islands correlate with the transcriptional activity of the locus – actively transcribed genes generally are not methylated while silenced genes are often found to be heavily methylated in the promoter island. Additionally, there is growing evidence that CpG islands found outside of transcription start sites play functional roles (Saxonov *et al.* 2006, Illingworth *et al.* 2010, Maunakea *et al.* 2010). While DNA methylation at gene promoters is traditionally thought to act as a binary switch (methylated, silent; unmethylated, active), it appears that CpG density, not just presence of methylation alone, also contributes to regulation of expression. For example, methylation at low CpG dense promoters still allows for transcriptional activity (Fouse *et al.* 2008). Furthermore, there are numerous examples, particularly of non-coding RNAs, that are transcribed, although the allele is heavily methylated (Bartolomei *et al.* 1993, Takada *et al.* 2002, Sleutels *et al.* 2003). These examples highlight that while there are general correlations of methylation status and gene activity, individual loci vary greatly.

In mammals, the molecular machinery responsible for adding a methyl group to cytosine residues (resulting 5-methylcytosine (5mC)) has been identified as a family of DNA methyltransferases (Dnmt1, Dnmt3a, Dnmt3b, and Dnmt3l). Dnmt3a and Dnmt3b are responsible for *de novo* methylation and play partially redundant but independently essential roles during early development. This includes methylation of repeat regions, imprinted loci, as well as genes involved in lineage decisions (Okano *et al.* 1999, Bourc'his *et al.* 2001, Kaneda *et al.* 2004). More specifically, Dnmt3a

and Dnmt3b help to establish *de novo* methylation in the blastocyst, allowing global 5mC levels to increase to that of somatic cells following implantation (Smith *et al.* 2012). Dnmt3a is maternally loaded in the oocyte and is the predominate methyltransferase in the oocyte and zygote (Kaneda *et al.* 2004, Kato *et al.* 2007), whereas Dnmt3b is transcribed upon zygotic genome activation (Watanabe *et al.* 2002) and is the primary mediator of *de novo* methylation during implantation (Borgel *et al.* 2010). Knockout studies in mice show that each of the Dnmts is required for viability (Li *et al.* 1992, Okano *et al.* 1999), highlighting the essential nature of *de novo* and maintenance methylation during development.

Dnmt1 has two functional transcripts that are expressed during development – Dnmt1s is expressed in somatic cells while Dnmt1o is specifically expressed as an oocyte specific form (Rouleau *et al.* 1992, Gaudet *et al.* 1998, Mertineit *et al.* 1998). Unlike Dnmt3a and Dnmt3b, Dnmt1 maintains CpG methylation by recognizing hemimethylated DNA and methylating the unmethylated strand, ensuring 5mC is maintained through DNA synthesis (Leonhardt *et al.* 1992, Arand *et al.* 2012). Targeting of Dnmt1 to replication foci occurs in most proliferating cells (Kishikawa *et al.* 2003, Bostick *et al.* 2007), however, Dnmt1o/s is largely excluded from the nucleus during early preimplantation stages (Howell *et al.* 2001), likely to allow for the large-scale demethylation that occurs to both haploid genomes (Figs 1 and 2).

Both sperm and oocytes contain parent- of- origin specific 5mC patterns. Therefore, at the time of fertilization the two haploid genomes arrive with diverse epigenomic signatures. Both parental pronuclei undergo dramatic global demethylation, presumably to ensure similar epigenetic information at the two parental alleles of the majority of genes (imprinted loci being one exception) as well as to program the newly formed zygote to a totipotent state. The male haploid genome is heavily methylated in sperm, where between 80 and 90% of all CpG dinucleotides are methylated (Mayer *et al.* 2000, Oswald *et al.* 2000, Santos *et al.* 2002). Global DNA methylation levels in the maternal haploid genome are approximately half that of the sperm (Howlett & Reik 1991, Smallwood *et al.* 2011, Peat *et al.* 2014). Shortly after fertilization, the two parental genomes undergo distinct but equally dramatic waves of DNA demethylation. The paternal genome undergoes active, replication-independent demethylation within the first several hours post-fertilization. In contrast, the maternal genome largely undergoes passive, cell division-dependent diffusion of methylation, resulting in demethylation over the course of preimplantation development.

Active DNA demethylation during preimplantation development by Tet3 oxidation of 5mC

Demethylation begins immediately in the newly formed embryo, prior to the first cell division. By the time the

Figure 1 DNA demethylation in the zygote. Distinct demethylation dynamics occur in the maternal (pink) and paternal (blue) pronuclei prior to fusion. At fertilization, the maternal haploid genome has ~40% methylation compared to nearly 90% methylation in the paternal haploid genome. Upon fertilization and continuing through pronuclear stage PN2, the paternal genome undergoes Tet3-dependent demethylation. PGC7, which preferentially binds to H3K9me2-rich chromatin, protects the maternal genome from Tet3 activity during PN stages. By PN5 stage, the bulk of paternal 5mC is gone and little change has occurred to maternal methylation. PPN, paternal pronucleus; MPN, maternal pronucleus.

embryo reaches the morula stage, the genome is almost completely devoid of DNA methylation (Santos *et al.* 2002). Despite wide-spread global demethylation, a few regions of the genome are protected, including imprinted loci and active retrotransposons like intracisternal A particle (IAP) elements (Lane *et al.* 2003). The large-scale demethylation begins with the rapid, active demethylation of the paternal haploid genome.

The differences between demethylation dynamics within the maternal and paternal pronuclei are thought to arise from their distinct architecture. The paternal genome is packed mostly around protamines, which are disassembled after fertilization and re-organized with histone-containing nucleosomes (Braun 2001, Balhorn 2007). The maternal genome is largely assembled around H3K9me2-rich histones. These structural distinctions between the two haploid genomes at pronuclear stage 0 (PN0) is thought to greatly influence the timing of bulk genome-wide demethylation (Santos *et al.* 2005), the kinetics of which are different between the maternal and paternal pronuclei (Fig. 1). Examination of global DNA methylation by immunofluorescence showed that the paternal pronucleus undergoes division-independent demethylation (Santos *et al.* 2002). When the zygote reaches the PN3 stage (~4 h after fertilization), there is already a dramatic loss of 5mC observed in the paternal pronucleus but little change in the maternal pronucleus (Mayer *et al.* 2000, Oswald *et al.* 2000). By the time of the first cell division (24 h after fertilization), there is no 5mC signal detected in the paternal PN, indicating near-complete loss of 5mC methylation. Even though the two parental genomes occupy the same nucleus after PN fusion, the differences in 5mC levels are apparent beyond the 4-cell stage (Santos *et al.* 2002).

Early studies of demethylation dynamics in mice based on immunofluorescence conflicted somewhat with

bisulfite DNA sequencing data sets, which did not show as dramatic a loss of 5mC (Oswald *et al.* 2000). It was not until the realization that the 5mC oxidation product 5-hydroxymethylcytosine (5hmC) is present *in vivo* that this discrepancy was resolved. Traditional bisulfite treatment does not distinguish between 5mC and 5hmC (Huang *et al.* 2010), while the antibodies used for immunofluorescence specifically (and only) detect 5mC. It has been subsequently shown that TET enzymes mediate the oxidation of 5mC to 5hmC (as well as 5-formylcytosine (5fC) and 5-carboxylcytosine (5caC)) *in vivo* and that TET proteins are expressed and differentially localized during preimplantation development (Gu *et al.* 2011, He *et al.* 2011, Iqbal *et al.* 2011, Wossidlo *et al.* 2011). Specifically, TET3 primarily localizes to the paternal pronucleus (Fig. 1) and is thought to be responsible for the observed rapid demethylation. Importantly, Gu *et al.* (2011) showed that the loss of 5mC corresponds with a concomitant gain in 5hmC. In both pronuclei, 5mC is present until PN3, and by late PN3, there is a detectable decrease in 5mC and an increase in 5hmC (Fig. 1; Gu *et al.* 2011, Iqbal *et al.* 2011, Wossidlo *et al.* 2011). It was also shown that Tet proteins convert 5mC to 5fC and 5caC as well, suggesting that Tet-mediated oxidation results in three oxidative forms for cytosine *in vivo* (Inoue 2011), ultimately resulting in replacement of the oxidized base with unmethylated cytosine by base excision repair or replication-dependent diffusion. Additionally, oxidation of 5mC has been shown in other mammalian zygotes, indicating a conserved mechanism of demethylation (Wossidlo *et al.* 2011).

Surprisingly, deletion of Tet3 activity results in retention of 5mC in the paternal pronucleus and inappropriate gene activation at many loci, but only mild global phenotype (reduced viability (Gu *et al.* 2011)). Tet3 mediated

Figure 2 DNA methylation dynamics from fertilization through gastrulation. The three main phases of methylation change are illustrated. During zygotic stages, the paternal genome undergoes active demethylation (blue line). This active demethylation is evident by sharply increased levels of oxidative products (5mC to 5hmC, 5fC and 5caC) in the paternal pronucleus. Passive replication-dependent demethylation (predominantly in the maternal pronucleus) occurs through exclusion of DNMTs from the nucleus during early preimplantation (red line). Lowest levels of methylation are reached between the morula to blastocyst stage, when methylation levels begin to rise. During blastocyst formation and gastrulation, the genome becomes re-methylated to levels consistent with somatic cells (black line).

hydrolysis of 5mC also occurs during reprogramming after SCNT cloning – Tet3 localizes to the pseudo-pronucleus in recombined zygotes. In SCNT embryos made with Tet3-null host oocytes, there is no 5hmC present in the pseudo-pronucleus, further indicating the role of Tet3 in active demethylation (Wossidlo *et al.* 2011, Gu *et al.* 2011).

These recent studies offered the prevailing idea that demethylation during preimplantation development occurred via i) active DNA demethylation of the paternal pronuclei mediated by Tet3 and thymine DNA glycosylase (TDG)-mediated base excision repair (reviewed in Kohli & Zhang (2013)) and ii) passive, replication-dependent dilution loss of methylation of the maternal

genome due to lack of Dnmt1 in the nucleus (Howell *et al.* 2001).

However, recent work is shifting the hypotheses about the mechanisms responsible *in vivo*. Using whole genome approaches to assess cytosine methylation patterns, it has been shown that Tet3-mediated demethylation is only partially responsible for paternal demethylation and that active demethylation also occurs in the maternal PN (Guo *et al.* 2014a, Shen *et al.* 2014). Furthermore, although Tet3-mediated oxidation is required for active demethylation, TDG-mediated base excision repair is not (Guo *et al.* 2014a). Additionally, it appears that there is conflicting evidence regarding the

role that replication-dependent demethylation plays in removing methylation from the paternal genome. By blocking replication of the paternal pronucleus, Shen *et al.* (2014) showed diminished demethylation in the paternal pronucleus even though Tet3 activity is present, indicating that replication is also involved in the demethylation of the paternal PN. Adding additional ambiguity is the fact that that Tet3-mediated demethylation is largely dispensable for successful development (Peat *et al.* 2014, Shen *et al.* 2014, Inoue *et al.* 2015).

Taken together the mechanisms that reprogram sperm and oocyte specific DNA methylation are not mutually exclusive as once predicted, and these recent stories indicate that there are likely unknown mechanisms also contributing to DNA methylation dynamics during preimplantation. With these advances, there are three main modes of DNA demethylation: i) active Tet3-mediated oxidation (predominantly in the paternal pronucleus); ii) replication-dependent dilution of Tet3-oxidative products, which plays a major role in demethylation of the paternal pronucleus and iii) replication-dependent (Tet3-independent) dilution of 5mC (predominantly in the maternal pronucleus). As our technical abilities evolve, it will be interesting to determine the interplay of these mechanisms within the same cells *in vivo*, define the specific loci at which each occurs and identify whether there are differing roles influencing cell fate decisions.

If Tet3 is present in the oocyte but only acts primarily on the paternal genome, there must be a protective mechanism to prevent conversion of maternal 5mC. One candidate for this maternal genome protection is PGC7/Stella, a DNA-binding protein expressed during germ cell specification and in gonads and oocytes (Saitou *et al.* 2002, Sato *et al.* 2002). PGC7 null embryos fail to complete preimplantation and there is a loss of 5mC in both pronuclei, indicating a protective role in the maternal pronucleus. Additionally, PGC7/Stella is targeted to differentially methylated regions (DMRs) of imprinted genes in the early embryo (Nakamura *et al.* 2007), supporting a functional role in blocking Tet3-mediated demethylation.

PGC7/Stella is able to protect the maternal pronucleus by binding to H3K9me2, which is a distinguishing feature of the maternal pronucleus. Loss of H3K9me2 by ectopic Jndm2a, a H3K9 methylation/dimethylation-specific demethylase, leads to loss of 5mC in both the maternal and parental pronuclei. PGC7/Stella also binds to H3K9me2 regions of the paternal pronucleus, including DMRs, which are not subject to protamine replacement. Tet3 is inhibited by PGC7/Stella, thus offering protection from active demethylation (Nakamura *et al.* 2012).

Imprinted loci are protected from demethylation

While most of the genome undergoes global DNA demethylation, imprinted loci are protected and retain

parent of origin DMRs (Branco 2008, Cirio 2008, Hirasawa 2008). It is clear that Dnmt1 is required for the maintenance of these imprinted sites (Bourc'his *et al.* 2001), even though it is largely excluded from the nucleus (Hirasawa 2008). Stella is also known to protect these loci, including some imprinted sites of the paternal genome. Additionally, Zfp57, a KRAB zinc finger protein, and Trim28 have also been shown to be required for integrity of ICRs in the early embryo. Trim28 interacts with Zfp57 to target it to specific imprinted sites, resulting in recruitment of repressive complexes including NuRD, SETDB1 and DMNTs (Iyengar *et al.* 2011, Quenneville *et al.* 2011, Zuo *et al.* 2012). While loss of maternal Zfp57 can be rescued by paternal expression, loss of maternal Trim28 is lethal (Li *et al.* 2008, Messerschmidt *et al.* 2012), due in part to the variation in loss of imprinted expression (Messerschmidt *et al.* 2012).

Early DNA demethylation dynamics in other mammalian species

Preimplantation DNA demethylation dynamics are largely the same in mouse and human embryos. However, this is not the case in all mammals, indicating distinct epigenetic reprogramming in different species. During the PN stages and in the first cell divisions, human, mouse and rat zygotes lose the majority of their paternal 5mC (Dean *et al.* 2001, Zaitseva *et al.* 2007). In contrast, both bovine and goat embryos retain an intermediate level of 5mC in the paternal pronuclei (Park *et al.* 2010, Wossidlo *et al.* 2010). Strikingly, sheep, pig and rabbit embryos retain 5mC during the PN stages and throughout preimplantation development (Beaujean *et al.* 2004, Jeong *et al.* 2007, Reis e Silva *et al.* 2012). In sheep, levels of 5mC drop during the 2-cell stage, but then increase at the 16-cell stage, and the ICM maintains levels of DNA methylation but the TE levels decrease dramatically (Young & Beaujean 2004).

These comparative studies illustrate the differences in timing and degree of 5mC loss during preimplantation among different mammalian species. These differences may be due in part to variation in zygotic genome activation, but they may also hint at differences in methylation reprogramming requirements needed to reach a totipotent state. These data also support the idea that active, Tet3-mediated demethylation in mice is not required for normal preimplantation development (Peat *et al.* 2014, Shen *et al.* 2014).

Next-generation sequencing to assess global DNA methylation dynamics during preimplantation development

Next-generation sequencing, including reduced representation bisulfite sequencing (RRBS) and whole-genome bisulfite sequencing (WGBS) now allow

assessment of global DNA methylation reprogramming with high resolution even from limited numbers of cells. Confirming earlier work, methylation across the genome is observed at relatively low levels in oocytes and early preimplantation stages, while sperm and post-implantation embryos have methylation similar to that of somatic cells (Smith *et al.* 2012, Guo *et al.* 2014a,b, Peat *et al.* 2014). These newer technologies have allowed for refined assessment of methylation changes across the genome during precise developmental stages, examination of specific classes of DNA sequence elements and comparison of mouse and human preimplantation embryos.

DNA methylation dynamics in human preimplantation embryos

Two groups recently examined genome-wide DNA methylation changes in human oocyte, sperm, zygote, pre- and post-implantation stages, using RRBS and WGBS (Smith *et al.* 2012, Guo *et al.* 2014a). Similar to mouse, both groups found that human sperm is highly methylated (although less than mouse sperm) and human oocytes have intermediate levels of methylation. The post-fertilization demethylation kinetics are also similar in the human zygote (Smith *et al.* 2012). Guo *et al.* (2014a) note that the greatest loss of DNA methylation occurred between the 1- to 2-cell stage in human embryos, rather than during PN stages (as is the case in the mouse). This could indicate that differences in the rate of active demethylation also correlates with the timing of zygotic genome activation, which occurs later in humans (Beaujean *et al.* 2004). Because bisulfite sequencing used in these studies did not distinguish between 5mC and the oxidative products of Tet-mediated demethylation, the distinct timing in mouse and human embryos may reflect a difference in the oxidation rates of 5mC or Tet activity between species. Paternal genome demethylation in humans is similar to observations in mouse zygotes, in that the majority of methylation is rapidly lost and only low levels of methylation remain during preimplantation. Levels of methylation in the maternal pronucleus are similar to mice, but the genomic regions that are demethylated are divergent. Additionally, unlike in mice, the majority of sperm and oocyte-specific DMRs regain their full methylation following implantation (Smith *et al.* 2012, Guo *et al.* 2014a).

Comparison of genome-wide methylation with single-cell RNA sequencing data (Yan *et al.* 2013) confirmed the previously observed negative correlation between promoter methylation and gene expression and highlighted that this inverse relationship strengthens after the MZT in human embryos (Smith *et al.* 2012, Guo *et al.* 2014b). Genes that had increased promoter methylation after the blastocyst stage showed a predicted decrease in expression in post-implantation-stage embryos (Guo *et al.* 2014b). Also as expected, changes in DNA

methylation during preimplantation influence the repression of transposable elements. SINE/variable number of tandem repeat/Alu elements (SVAs) expression increases after the 2-cell stage, when rapid demethylation occurs. This expression is maintained until the morula stage, when expression decreases, presumably as the genome is re-methylated – a trend which continues post-implantation (Smith *et al.* 2012, Guo *et al.* 2014b).

While many repeat elements undergo loss of DNA methylation and increased expression, the evolutionary age of the transposable element appears to influence the retention of methylation during preimplantation development. Evolutionarily younger elements, which are still capable of transposition are relatively resistant to demethylation while their evolutionarily older counterparts that have lost the ability to jump are readily demethylated along with coding genes. This might hint at the evolutionary origins of methylation/demethylation dynamics in mammalian preimplantation development (Wang *et al.* 2014).

Histone modifications during preimplantation

In addition to DNA methylation changes, chromatin organization and histone modifications play a critical role in establishing a totipotent embryo, as well as directing the first lineage decisions. Chromatin is a highly organized and dynamic nuclear structure containing DNA, histones and many other proteins. Nucleosomes, the basic building block of chromatin are comprised of two each of histone H2A, H2B, H3 and H4. It is well established that the N-terminal tails of these core histones are subject to post-translational modifications (PTMs), which play a fundamental role in influencing gene expression patterns among disparate cell types (Fischle *et al.* 2003). Histone PTMs include acetylation, methylation, phosphorylation, ubiquitination and others, which occur at specific amino acid residues catalyzed by specific enzymes (Strahl & Allis 2000, Tan *et al.* 2011). Additional complexity arises in that methylation at lysines or arginines may exist in distinct forms: mono-, di-, or trimethyl for lysines and mono- or dimethyl on arginine residues (Kouzarides 2007). A general theme has emerged in which PTMs are catalyzed by opposite functional pairs of enzymes.

Many studies have revealed functional themes where histone PTMs correlate with gene expression patterns. For example, lysine acetylation is commonly considered to be an active mark that correlates with chromatin accessibility and active transcription, whereas histone lysine methylation can be either active or repressive depending on the particular lysine residue that is modified (Tsukada *et al.* 2006, Bernstein *et al.* 2007). Recent large-scale efforts supported by the Roadmap Epigenomics Project are defining 'chromatin states' in many diverse tissues – that is, combinations of histone modifications, DNA methylation and transcription factor binding that correlate with

functional property of a particular locus (<http://www.roadmapepigenomics.org/publications/>).

Histone modification during preimplantation embryo development

Studies of early embryonic development have shown that shortly after fertilization, many histone modifications are observed asymmetrically in the parental haploid genomes prior to PN fusion (summarized in Fig. 3). For example, in mice, H3K27ac, H4K5ac and H4K16ac are only detectable in the paternal PN of early zygotes (Adenot *et al.* 1997, Stein *et al.* 1997, Hayashi-Takanaka *et al.* 2011). Conversely, all forms of H3K4 methylation (me1, me2 and me3) are observed in maternal PN (Lepikhov & Walter 2004, Santenard *et al.* 2010), and H3K9me2 and me3 are also significantly higher in the maternal PN (Lepikhov & Walter 2004, Wongtawan *et al.* 2011, Beaujean 2014). H3K27me1 is present in both PNs, but H3K27me2 and -me3 occur extensively in the maternal PN (Erhardt *et al.* 2003, Santos *et al.* 2005, Santenard *et al.* 2010). Additionally, H3K9me3S10P, H3K36me3 and H4K20me3 are also found exclusively in the maternal PN at early post-fertilization stages (Boskovic *et al.* 2012, Ribeiro-Mason *et al.* 2012, Beaujean 2014). Although the functional significance of these asymmetric PTMs remains largely unknown, it highlights the distinct reprogramming that is required for the paternal and maternal PN for proper embryonic genome activation and embryo development (Ribeiro-Mason *et al.* 2012, Beaujean 2014).

Histone PTMs also play key roles in remodeling of chromatin configuration and DNA methylation. In mice, the increase of H3K79me by forced expression of DOT1L causes premature chromocenter formation and developmental arrest of 2-cell embryos (Ooga *et al.* 2013). Additionally, deletion of the methyltransferase Setdb1 in

mouse embryonic stem cells leads to the reduction of H3K9me3 and an overall decrease of DNA methylation levels at specific loci (Leung *et al.* 2014). In porcine embryos, disturbed H3K4me3-H3K27me3 balance after knockdown of demethylase Kdm5b can cause increased expressions of Tet family members (Huang *et al.* 2015), which are found to be crucial for the interactions between histone modification and DNA methylation in mouse embryonic stem cells (Sui *et al.* 2012).

Functional studies of the roles of specific modifications are just beginning, using genetic strategies to add or remove specific enzymatic activities to embryos. For example in mice, hyperacetylation of histone H4 mediated by knockdown of HDAC1 causes developmental delay (Ma & Schultz 2008). Knockdown of either *Ing2* (H3K4me3 methyltransferase activity) or RNF20 (histone H2B monoubiquitination) results in arrest at the morula stage (Ooga *et al.* 2015, Zhou *et al.* 2015). Depletion of H4K20me1 by knockout of the *PR-Set7* gene induces early embryonic lethality prior to the 8-cell stage (Oda *et al.* 2009), and our lab has shown critical roles of H3K36me3 during preimplantation development by knockdown of CTR9/PAF1 (Zhang *et al.* 2013a). Other recent examples include studies in mice showing that maternal-specific H3K9me3 is enriched at the *Xist* promoter region and prevents maternal *Xist* activation (Fukuda *et al.* 2014); increased H3K4me2 results in abnormal expression of *eIF-4C/Oct4* and arrest at the 2-cell stage (Shao *et al.* 2008); and that PRC1 binding to H3K27me3 plays an indispensable role in embryonic genome activation and developmental progression (Posfai *et al.* 2012). Although precise in the removal of specific gene function, these studies highlight the difficulty in assigning specific function to a particular modification or enzymatic activity since the phenotype is often developmental failure and misregulation of many genes. It is only very recently that next-generation technologies allow for very low input such that

Figure 3 Differential histone modification during preimplantation development. After fertilization, the parental pronuclei are differentially enriched with many distinct histone modifications (left side: list of PTMs). Little data is available regarding the timing and mechanisms resulting after pronuclear fusion that result in largely homogenous PTMs during cleavage stages (indicated by the question mark). By the time ICM and TE begin to differentiate, these cell lineages have acquired distinct epigenetic signatures and gene expression patterns (right side).

investigators can determine which loci across the genome are altered in these knockout/knockdown embryos (Brind'Amour *et al.* 2015).

Relevant to artificial reproductive technologies, histone modifications are sensitive to manipulations during preimplantation, potentially altering epigenomic patterns (Feil & Fraga 2012, Dupont *et al.* 2012). For example, in the mouse, H3K4me3 is significantly lower in the *in vitro* fertilized embryos compared with *in vivo* fertilized embryos (Wu *et al.* 2012). Similarly lower levels of H3K27me3 are found in the ICM of heated-sperm-derived blastocysts when compared to untreated-sperm-derived blastocysts (Chao *et al.* 2012), and cryopreservation can alter H4K12ac patterns in both oocytes and zygotes (Suo *et al.* 2010). Despite these observations, it remains unclear if altered PTM levels persist in offspring or if surviving individuals contain appropriate epigenomic information – possibly correcting the epigenome during cell lineage differentiation at post-implantation stages.

Histone modifications in ICM and TE lineage specification

In mouse embryos, transcription factors such as Oct4, Sox2 and Nanog are enriched in cells of the ICM and function to both promote pluripotency and resist differentiation. Conversely, in TE, transcription factors such as Cdx2 and Eomes become upregulated, promoting differentiation. In contrast to the mouse, Oct4 and Cdx2 are co-expressed in the ICM and TE of bovine and porcine embryos, and the mechanisms of molecular differentiation remain largely unknown (Kirchhof *et al.* 2000).

This first lineage specification is critical for implantation and successful development. DNA methylation has been shown to be dispensable for growth and differentiation of the extraembryonic lineages (Sakaue *et al.* 2010), suggesting that appropriate histone modifications may provide key epigenetic information directing gene expression and lineage specification. Once the TE and ICM become distinct, they exhibit asymmetries in specific histone PTMs. For example, in the mouse, H4- and H2AS1P are increased in the TE cells (Sarmiento *et al.* 2004), while H3K27me3 is enriched in the ICM (Erhardt *et al.* 2003). At the 4-cell stage, blastomeres have different levels of methylated H3R26me and those cells with higher H3R26me are more likely to result in ICM cell fate. Overexpression of the H3R26 methyltransferase CARM1 results in increased expression Nanog and Sox2, suggesting that pluripotency factor expression is influenced by locus-specific H3R26me (Torres-Padilla *et al.* 2007). Other examples include studies showing that repressive H3K9me3 at the *Cdx2* promoter is important for maintaining pluripotency and loss of associated methyltransferase ESET in early embryos results in ICM failure (Yeap *et al.* 2009). However, in TE lineage, Suv39h

methyltransferase mediates repressive H3K9me3 at ICM-specific gene promoters in the TE lineage (Alder *et al.* 2010, Rugg-Gunn *et al.* 2010). These studies highlight that even the same histone modification can be finely tuned by distinct enzymes to influence lineage specification in different cell populations.

There are ever growing observations of locus specific enrichment of histone modifications correlating with lineage decisions during preimplantation development. For example, H3K4me3 and H3K27me3 are enriched at promoters of genes exclusively expressed in ICM or TE in both murine and bovine embryos (Dahl *et al.* 2010, Herrmann *et al.* 2013). It was also recently shown that loss of repressive H3K27me3 participation at TE-specific genes is essential for TE lineage development and embryo implantation (Saha *et al.* 2013, Paul & Knott 2014). In addition to methylation of histone H3 residues, acetylation of histone H4 (H4K8ac and H4K12ac) has also been implicated in early lineage specification (VerMilyea *et al.* 2009, Zhang *et al.* 2013b).

A handful of histone-modifying proteins thought to be central to epigenetic programming during development have knock-out phenotypes only apparent after preimplantation. These include members of Polycomb Repressive Complex 2 (Eed, Ezh2 and Suz12) as well as the H3K9 methyltransferases G9a and Eset. It remains unclear if the timing of null phenotypes is due to functional redundancy with other genes or maternal loading of RNA/protein, or if the modifications they perform are in fact not required until gastrulation (or later). There are a few histone-modifying enzyme knock-out phenotypes in mice that do result in lethality during preimplantation, some of which show lineage-specific defects. Loss of the histone H3K9 demethylase Jmjd2C results in morula arrest and null embryos show reduced levels of ICM-specific gene transcription, suggesting a failure to maintain pluripotency (Wang *et al.* 2010). Similarly, null embryos of several members of the NuRD complex (Sin3A, Suds3, Arid4b (McDonel *et al.* 2009)) and PAF1 complex (Ctr9 and Rtf1 (Ding *et al.* 2009, Zhang *et al.* 2013a)) do form blastocysts but show defects in ICM proliferation as a major cause of developmental lethality and failure. It is perhaps not so surprising that knockout of genes with distinct functions (such as Sin3A and Ctr9) result in similar defects in maintenance of ICM potency, which is of the utmost importance for continued development and requires myriad proteins to accomplish.

Moving forward

As described above, a wide array of covalent histone modifications are now recognized to occur *in vivo* and correlate with distinct transcriptional states and/or chromatin conformation. However, knowledge about the role of histone modifications during development is mostly limited to reports of changes in global patterns – apparent by immunofluorescence with antibodies

directed against specific modifications (reviewed in [Beaujean \(2014\)](#)). While these descriptive studies are an essential beginning, little is known about the functional importance of these modifications. *In vivo* analysis of the role of histone modifications at specific loci during early development is only just beginning, and the relative lack of functional data is due to several factors including: i) limitations in our ability to efficiently generate maternal and zygote null embryos at the same time, ii) limitations in our ability to assess histone modifications at specific loci from very small numbers of cells and iii) an inability to alter specific modifications at specific loci. Due to the combinatorial nature of the histone code and the difficulty in functionally preventing one particular modification at one locus *in vivo*, it is currently not feasible to simply ask, 'What is the role of a specific histone modification at a specific genomic locus during development'. Fortunately, this type of epigenetic engineering has come to the fore and many groups are currently working to develop *in vivo* epigenetic targeting tools.

With greatly enhanced access to next-generation sequencing technologies, there is ever-growing opportunity to probe genome-wide methylation patterns at single-base/nucleosome resolution in diverse cell populations, and improved techniques are pushing WGBS towards single-cell sequencing. Additionally, multiple methods are now readily available for the discrimination of 5mC and 5hmC at a single base resolution. Combining DNA methylation analysis with ChIP-seq and RNA-seq during preimplantation development will allow for a comprehensive cataloguing of early epigenetic reprogramming dynamics. Cross-species comparison of these dynamics at specific loci and the capability to functionally test the importance of specific modifications will allow for deeper understanding of how epigenetic dynamics influence preimplantation development, the transition from gametes to totipotency and the requirements of lineage differentiation.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the review.

Funding

This was supported in part by March of Dime Research Grant #6-FY11-367 and NIH 1R21HD078942-01 to J Mager.

References

Adenot PG, Mercier Y, Renard JP & Thompson EM 1997 Differential H4 acetylation of paternal and maternal chromatin precedes DNA replication and differential transcriptional activity in pronuclei of 1-cell mouse embryos. *Development* **124** 4615–4625.

- Alder O, Laval F, Helness A, Brookes E, Pinho S, Chandrasekaran A, Arnaud P, Pombo A, O'Neill L & Azuara V 2010 Ring1B and Suv39h1 delineate distinct chromatin states at bivalent genes during early mouse lineage commitment. *Development* **137** 2483–2492. (doi:10.1242/dev.048363)
- Arand J, Spieler D, Karius T, Branco MR, Meilinger D, Meissner A, Jenwein T, Xu G, Leonhardt H, Wolf V *et al.* 2012 *In vivo* control of CpG and non-CpG DNA methylation by DNA methyltransferases. *PLoS Genetics* **8** e1002750. (doi:10.1371/journal.pgen.1002750)
- Balhorn R 2007 The protamine family of sperm nuclear proteins. *Genome Biology* **8** 227. (doi:10.1186/gb-2007-8-9-227)
- Bartolomei MS, Webber AL, Brunkow ME & Tilghman SM 1993 Epigenetic mechanisms underlying the imprinting of the mouse H19 gene. *Genes and Development* **7** 1663–1673. (doi:10.1101/gad.7.9.1663)
- Beaujean N 2014 Histone post-translational modifications in preimplantation mouse embryos and their role in nuclear architecture. *Molecular Reproduction and Development* **81** 100–112. (doi:10.1002/mrd.22268)
- Beaujean N, Hartshorne G, Cavilla J, Taylor J, Gardner J, Wilmut I, Meehan R & Young L 2004 Non-conservation of mammalian pre-implantation methylation dynamics. *Current Biology* **14** R266–R267. (doi:10.1016/j.cub.2004.03.019)
- Bernstein BE, Meissner A & Lander ES 2007 The mammalian epigenome. *Cell* **128** 669–681. (doi:10.1016/j.cell.2007.01.033)
- Biggers JD, Bell JE & Benos DJ 1988 Mammalian blastocyst: transport functions in a developing epithelium. *American Journal of Physiology* **255** C419–C432.
- Borgel J, Guibert S, Li Y, Chiba H, Schubeler D, Sasaki H, Forne T & Weber M 2010 Targets and dynamics of promoter DNA methylation during early mouse development. *Nature Genetics* **42** 1093–1100. (doi:10.1038/ng.708)
- Boskovic A, Bender A, Gall L, Ziegler-Birling C, Beaujean N & Torres-Padilla ME 2012 Analysis of active chromatin modifications in early mammalian embryos reveals uncoupling of H2A.Z acetylation and H3K36 trimethylation from embryonic genome activation. *Epigenetics* **7** 747–757. (doi:10.4161/epi.20584)
- Bostick M, Kim JK, Esteve PO, Clark A, Pradhan S & Jacobsen SE 2007 UHRF1 plays a role in maintaining DNA methylation in mammalian cells. *Science* **317** 1760–1764. (doi:10.1126/science.1147939)
- Bourc'his D, Xu GL, Lin CS, Bollman B & Bestor TH 2001 Dnmt3L and the establishment of maternal genomic imprints. *Science* **294** 2536–2539. (doi:10.1126/science.1065848)
- Branco MR, Oda M & Reik W 2008 Safeguarding parental identity: Dnmt1 maintains imprints during epigenetic reprogramming in early embryogenesis. *Genes and Development* **22** 1567–1571. (doi:10.1101/gad.1690508)
- Braun RE 2001 Packaging paternal chromosomes with protamine. *Nature Genetics* **28** 10–12.
- Brind'Amour J, Liu S, Hudson M, Chen C, Karimi MM & Lorincz MC 2015 An ultra-low-input native ChIP-seq protocol for genome-wide profiling of rare cell populations. *Nature Communications* **6** 6033. (doi:10.1038/ncomms7033)
- Chao SB, Chen L, Li JC, Ou XH, Huang XJ, Wen S, Sun QY & Gao GL 2012 Defective histone H3K27 trimethylation modification in embryos derived from heated mouse sperm. *Microscopy and Microanalysis* **18** 476–482. (doi:10.1017/S1431927612000396)
- Cirio MC, Ratnam S, Ding F, Reinhart B, Navara C & Chaillet JR 2008 Preimplantation expression of the somatic form of Dnmt1 suggests a role in the inheritance of genomic imprints. *BMC Developmental Biology* **8** 9. (doi:10.1186/1471-213X-8-9)
- Dahl JA, Reiner AH, Klungland A, Wakayama T & Collas P 2010 Histone H3 lysine 27 methylation asymmetry on developmentally-regulated promoters distinguish the first two lineages in mouse preimplantation embryos. *PLoS ONE* **5** e9150. (doi:10.1371/journal.pone.0009150)
- Dean W, Santos F, Stojkovic M, Zakhartchenko V, Walter J, Wolf E & Reik W 2001 Conservation of methylation reprogramming in mammalian development: aberrant reprogramming in cloned embryos. *PNAS* **98** 13734–13738. (doi:10.1073/pnas.241522698)
- Ding L, Paszkowski-Rogacz M, Nitzsche A, Slabicki MM, Heninger AK, de Vries I, Kittler R, Junqueira M, Shevchenko A, Schulz H *et al.* 2009 A genome-scale RNAi screen for Oct4 modulators defines a role of the Paf1 complex for embryonic stem cell identity. *Cell Stem Cell* **4** 403–415. (doi:10.1016/j.stem.2009.03.009)

- Dupont C, Cordier AG, Junien C, Mandon-Pepin B, Levy R & Chavatte-Palmer P 2012 Maternal environment and the reproductive function of the offspring. *Theriogenology* **78** 1405–1414. (doi:10.1016/j.theriogenology.2012.06.016)
- Erhardt S, Su IH, Schneider R, Barton S, Bannister AJ, Perez-Burgos L, Jenuwein T, Kouzarides T, Tarakhovsky A & Surani MA 2003 Consequences of the depletion of zygotic and embryonic enhancer of zeste 2 during preimplantation mouse development. *Development* **130** 4235–4248. (doi:10.1242/dev.00625)
- Feil R & Fraga MF 2012 Epigenetics and the environment: emerging patterns and implications. *Nature Reviews. Genetics* **13** 97–109. (doi:10.1038/nrg3142)
- Fischle W, Wang Y & Allis CD 2003 Binary switches and modification cassettes in histone biology and beyond. *Nature* **425** 475–479. (doi:10.1038/nature02017)
- Fleming TP, Sheth B & Fesenko I 2001 Cell adhesion in the preimplantation mammalian embryo and its role in trophectoderm differentiation and blastocyst morphogenesis. *Frontiers in Bioscience* **6** D1000–D1007. (doi:10.2741/Fleming)
- Fouse SD, Shen Y, Pellegrini M, Cole S, Meissner A, Van Neste L, Jaenisch R & Fan G 2008 Promoter CpG methylation contributes to ES cell gene regulation in parallel with Oct4/Nanog, PcG complex, and histone H3 K4/K27 trimethylation. *Cell Stem Cell* **2** 160–169. (doi:10.1016/j.stem.2007.12.011)
- Fukuda A, Tomikawa J, Miura T, Hata K, Nakabayashi K, Eggan K, Akutsu H & Umezawa A 2014 The role of maternal-specific H3K9me3 modification in establishing imprinted X-chromosome inactivation and embryogenesis in mice. *Nature Communications* **5** 5464. (doi:10.1038/ncomms6464)
- Gaudet F, Talbot D, Leonhardt H & Jaenisch R 1998 A short DNA methyltransferase isoform restores methylation *in vivo*. *Journal of Biological Chemistry* **273** 32725–32729. (doi:10.1074/jbc.273.49.32725)
- Gu TP, Guo F, Yang H, Wu HP, Xu GF, Liu W, Xie ZG, Shi L, He X, Jin SG *et al.* 2011 The role of Tet3 DNA dioxygenase in epigenetic reprogramming by oocytes. *Nature* **477** 606–610. (doi:10.1038/nature10443)
- Guo F, Li X, Liang D, Li T, Zhu P, Guo H, Wu X, Wen L, Gu TP, Hu B *et al.* 2014a Active and passive demethylation of male and female pronuclear DNA in the Mammalian zygote. *Cell Stem Cell* **15** 447–458. (doi:10.1016/j.stem.2014.08.003)
- Guo H, Zhu P, Yan L, Li R, Hu B, Lian Y, Yan J, Ren X, Lin S, Li J *et al.* 2014b The DNA methylation landscape of human early embryos. *Nature* **511** 606–610. (doi:10.1038/nature13544)
- Hayashi-Takanaka Y, Yamagata K, Wakayama T, Stasevich TJ, Kainuma T, Tsurimoto T, Tachibana M, Shinkai Y, Kurumizaka H, Nozaki N *et al.* 2011 Tracking epigenetic histone modifications in single cells using Fab-based live endogenous modification labeling. *Nucleic Acids Research* **39** 6475–6488. (doi:10.1093/nar/gkr343)
- He YF, Li BZ, Li Z, Liu P, Wang Y, Tang Q, Ding J, Jia Y, Chen Z, Li L *et al.* 2011 Tet-mediated formation of 5-carboxylcytosine and its excision by TDG in mammalian DNA. *Science* **333** 1303–1307. (doi:10.1126/science.1210944)
- Herrmann D, Dahl JA, Lucas-Hahn A, Collas P & Niemann H 2013 Histone modifications and mRNA expression in the inner cell mass and trophectoderm of bovine blastocysts. *Epigenetics* **8** 281–289. (doi:10.4161/epi.23899)
- Hirasawa R, Chiba H, Kaneda M, Tajima S, Li E, Jaenisch R & Sasaki H 2008 Maternal and zygotic Dnmt1 are necessary and sufficient for the maintenance of DNA methylation imprints during preimplantation development. *Genes and Development* **22** 1607–1616. (doi:10.1101/gad.1667008)
- Howell CY, Bestor TH, Ding F, Latham KE, Mertineit C, Trasler JM & Chaillet JR 2001 Genomic imprinting disrupted by a maternal effect mutation in the Dnmt1 gene. *Cell* **104** 829–838. (doi:10.1016/S0092-8674(01)00280-X)
- Howlett SK & Reik W 1991 Methylation levels of maternal and paternal genomes during preimplantation development. *Development* **113** 119–127.
- Huang Y, Pastor WA, Shen Y, Tahiliani M, Liu DR & Rao A 2010 The behaviour of 5-hydroxymethylcytosine in bisulfite sequencing. *PLoS ONE* **5** e8888. (doi:10.1371/journal.pone.0008888)
- Huang J, Zhang H, Wang X, Dobbs KB, Yao J, Qin G, Whitworth K, Walters EM, Prather RS & Zhao J 2015 Impairment of preimplantation porcine embryo development by histone demethylase KDM5B knockdown through disturbance of bivalent H3K4me3-H3K27me3 modifications. *Biology of Reproduction* **92** 72. (doi:10.1095/biolreprod.114.122762)
- Illingworth RS, Gruenewald-Schneider U, Webb S, Kerr AR, James KD, Turner DJ, Smith C, Harrison DJ, Andrews R & Bird AP 2010 Orphan CpG islands identify numerous conserved promoters in the mammalian genome. *PLoS Genetics* **6** e1001134. (doi:10.1371/journal.pgen.1001134)
- Inoue A & Zhang Y 2011 Replication-dependent loss of 5-hydroxymethylcytosine in mouse preimplantation embryos. *Science* **334** 194. (doi:10.1126/science.1212483)
- Inoue A, Shen L, Matoba S & Zhang Y 2015 Haploinsufficiency, but not defective paternal 5mC oxidation, accounts for the developmental defects of maternal Tet3 knockouts. *Cell Reports* **10** 463–470. (doi:10.1016/j.celrep.2014.12.049)
- Iqbal K, Jin SG, Pfeifer GP & Szabo PE 2011 Reprogramming of the paternal genome upon fertilization involves genome-wide oxidation of 5-methylcytosine. *PNAS* **108** 3642–3647. (doi:10.1073/pnas.1014033108)
- Iyengar S, Ivanov AV, Jin VX, Rauscher FJ III & Farnham PJ 2011 Functional analysis of KAP1 genomic recruitment. *Molecular and Cellular Biology* **31** 1833–1847. (doi:10.1128/MCB.01331-10)
- Jeong YS, Yeo S, Park JS, Koo DB, Chang WK, Lee KK & Kang YK 2007 DNA methylation state is preserved in the sperm-derived pronucleus of the pig zygote. *International Journal of Developmental Biology* **51** 707–714. (doi:10.1387/ijdb.072450yj)
- Kaneda M, Okano M, Hata K, Sado T, Tsujimoto N, Li E & Sasaki H 2004 Essential role for *de novo* DNA methyltransferase Dnmt3a in paternal and maternal imprinting. *Nature* **429** 900–903. (doi:10.1038/nature02633)
- Kato Y, Kaneda M, Hata K, Kumaki K, Hisano M, Kohara Y, Okano M, Li E, Nozaki M & Sasaki H 2007 Role of the Dnmt3 family in *de novo* methylation of imprinted and repetitive sequences during male germ cell development in the mouse. *Human Molecular Genetics* **16** 2272–2280. (doi:10.1093/hmg/ddm179)
- Kidder GM & Winterhager E 2001 Intercellular communication in preimplantation development: the role of gap junctions. *Frontiers in Bioscience* **6** D731–D736. (doi:10.2741/Kidder)
- Kirchhof N, Carnwath JW, Lemme E, Anastassiadis K, Scholer H & Niemann H 2000 Expression pattern of Oct-4 in preimplantation embryos of different species. *Biology of Reproduction* **63** 1698–1705. (doi:10.1095/biolreprod63.6.1698)
- Kishikawa S, Murata T, Ugai H, Yamazaki T & Yokoyama KK 2003 Control elements of Dnmt1 gene are regulated in cell-cycle dependent manner. *Nucleic Acids Research. Supplement (2001)* **3** 307–308. (doi:10.1093/nass/3.1.307)
- Kohli RM & Zhang Y 2013 TET enzymes, TDG and the dynamics of DNA demethylation. *Nature* **502** 472–479. (doi:10.1038/nature12750)
- Kouzarides T 2007 Chromatin modifications and their function. *Cell* **128** 693–705. (doi:10.1016/j.cell.2007.02.005)
- Lane N, Dean W, Erhardt S, Hajkova P, Surani A, Walter J & Reik W 2003 Resistance of IAPs to methylation reprogramming may provide a mechanism for epigenetic inheritance in the mouse. *Genesis* **35** 88–93. (doi:10.1002/gene.10168)
- Latham KE, Solter D & Schultz RM 1991 Activation of a two-cell stage-specific gene following transfer of heterologous nuclei into enucleated mouse embryos. *Molecular Reproduction and Development* **30** 182–186. (doi:10.1002/mrd.1080300303)
- Leonhardt H, Page AW, Weier HU & Bestor TH 1992 A targeting sequence directs DNA methyltransferase to sites of DNA replication in mammalian nuclei. *Cell* **71** 865–873. (doi:10.1016/0092-8674(92)90561-P)
- Lepikhov K & Walter J 2004 Differential dynamics of histone H3 methylation at positions K4 and K9 in the mouse zygote. *BMC Developmental Biology* **4** 12. (doi:10.1186/1471-213X-4-12)
- Leung D, Du T, Wagner U, Xie W, Lee AY, Goyal P, Li Y, Szulwach KE, Jin P, Lorincz MC *et al.* 2014 Regulation of DNA methylation turnover at LTR retrotransposons and imprinted loci by the histone methyltransferase Setdb1. *PNAS* **111** 6690–6695. (doi:10.1073/pnas.1322273111)

- Li E, Bestor TH & Jaenisch R 1992 Targeted mutation of the DNA methyltransferase gene results in embryonic lethality. *Cell* **69** 915–926. (doi:10.1016/0092-8674(92)90611-F)
- Li X, Ito M, Zhou F, Youngson N, Zuo X, Leder P & Ferguson-Smith AC 2008 A maternal-zygotic effect gene, *Zfp57*, maintains both maternal and paternal imprints. *Developmental Cell* **15** 547–557. (doi:10.1016/j.devcel.2008.08.014)
- Ma P & Schultz RM 2008 Histone deacetylase 1 (HDAC1) regulates histone acetylation, development, and gene expression in preimplantation mouse embryos. *Developmental Biology* **319** 110–120. (doi:10.1016/j.ydbio.2008.04.011)
- Majumder S & DePamphilis ML 1994 Requirements for DNA transcription and replication at the beginning of mouse development. *Journal of Cellular Biochemistry* **55** 59–68. (doi:10.1002/jcb.240550107)
- Maunakea AK, Nagarajan RP, Bilenky M, Ballinger TJ, D'Souza C, Fouse SD, Johnson BE, Hong C, Nielsen C, Zhao Y *et al.* 2010 Conserved role of intragenic DNA methylation in regulating alternative promoters. *Nature* **466** 253–257. (doi:10.1038/nature09165)
- Mayer W, Niveleau A, Walter J, Fundele R & Haaf T 2000 Demethylation of the zygotic paternal genome. *Nature* **403** 501–502. (doi:10.1038/35000656)
- McDonel P, Costello I & Hendrich B 2009 Keeping things quiet: roles of NuRD and Sin3 co-repressor complexes during mammalian development. *International Journal of Biochemistry & Cell Biology* **41** 108–116. (doi:10.1016/j.biocel.2008.07.022)
- Mertineit C, Yoder JA, Taketo T, Laird DW, Trasler JM & Bestor TH 1998 Sex-specific exons control DNA methyltransferase in mammalian germ cells. *Development* **125** 889–897.
- Messerschmidt DM, de Vries W, Ito M, Solter D, Ferguson-Smith A & Knowles BB 2012 Trim28 is required for epigenetic stability during mouse oocyte to embryo transition. *Science* **335** 1499–1502. (doi:10.1126/science.1216154)
- Nakamura T, Arai Y, Umehara H, Masuhara M, Kimura T, Taniguchi H, Sekimoto T, Ikawa M, Yoneda Y, Okabe M *et al.* 2007 PGC7/Stella protects against DNA demethylation in early embryogenesis. *Nature Cell Biology* **9** 64–71. (doi:10.1038/ncb1519)
- Nakamura T, Liu YJ, Nakashima H, Umehara H, Inoue K, Matoba S, Tachibana M, Ogura A, Shinkai Y & Nakano T 2012 PGC7 binds histone H3K9me2 to protect against conversion of 5mC to 5hmC in early embryos. *Nature* **486** 415–419. (doi:10.1038/nature11093)
- Nichols J & Gardner RL 1984 Heterogeneous differentiation of external cells in individual isolated early mouse inner cell masses in culture. *Journal of Embryology and Experimental Morphology* **80** 225–240.
- Nothias JY, Majumder S, Kaneko KJ & DePamphilis ML 1995 Regulation of gene expression at the beginning of mammalian development. *Journal of Biological Chemistry* **270** 22077–22080. (doi:10.1074/jbc.270.38.22077)
- Oda H, Okamoto I, Murphy N, Chu J, Price SM, Shen MM, Torres-Padilla ME, Heard E & Reinberg D 2009 Monomethylation of histone H4-lysine 20 is involved in chromosome structure and stability and is essential for mouse development. *Molecular and Cellular Biology* **29** 2278–2295. (doi:10.1128/MCB.01768-08)
- Okano M, Bell DW, Haber DA & Li E 1999 DNA methyltransferases Dnmt3a and Dnmt3b are essential for *de novo* methylation and mammalian development. *Cell* **99** 247–257. (doi:10.1016/S0092-8674(00)81656-6)
- Ooga M, Suzuki MG & Aoki F 2013 Involvement of DOT1L in the remodeling of heterochromatin configuration during early preimplantation development in mice. *Biology of Reproduction* **89** 145. (doi:10.1095/biolreprod.113.113258)
- Ooga M, Suzuki MG & Aoki F 2015 Involvement of histone H2B monoubiquitination in the regulation of mouse preimplantation development. *Journal of Reproduction and Development* **61** 179–184.
- Oswald J, Engemann S, Lane N, Mayer W, Olek A, Fundele R, Dean W, Reik W & Walter J 2000 Active demethylation of the paternal genome in the mouse zygote. *Current Biology* **10** 475–478. (doi:10.1016/S0960-9822(00)00448-6)
- Park JS, Lee D, Cho S, Shin ST & Kang YK 2010 Active loss of DNA methylation in two-cell stage goat embryos. *International Journal of Developmental Biology* **54** 1323–1328. (doi:10.1387/ijdb.092973jp)
- Paul S & Knott JG 2014 Epigenetic control of cell fate in mouse blastocysts: the role of covalent histone modifications and chromatin remodeling. *Molecular Reproduction and Development* **81** 171–182. (doi:10.1002/mrd.22219)
- Peat JR, Dean W, Clark SJ, Krueger F, Smallwood SA, Ficiz G, Kim JK, Marioni JC, Hore TA & Reik W 2014 Genome-wide bisulfite sequencing in zygotes identifies demethylation targets and maps the contribution of TET3 oxidation. *Cell Reports* **9** 1990–2000. (doi:10.1016/j.celrep.2014.11.034)
- Pedersen RA, Wu K & Balakier H 1986 Origin of the inner cell mass in mouse embryos: cell lineage analysis by microinjection. *Developmental Biology* **117** 581–595. (doi:10.1016/0012-1606(86)90327-1)
- Pesce M & Scholer HR 2001 Oct-4: gatekeeper in the beginnings of mammalian development. *Stem Cells* **19** 271–278. (doi:10.1634/stemcells.19-4-271)
- Posfai E, Kunzmann R, Brochard V, Salvaing J, Cabuy E, Roloff TC, Liu Z, Tardat M, van Lohuizen M, Vidal M *et al.* 2012 Polycomb function during oogenesis is required for mouse embryonic development. *Genes and Development* **26** 920–932. (doi:10.1101/gad.188094.112)
- Quenneville S, Verde G, Corsinotti A, Kappoulou A, Jakobsson J, Offner S, Baglivo I, Pedone PV, Grimaldi G, Riccio A *et al.* 2011 In embryonic stem cells, ZFP57/KAP1 recognize a methylated hexanucleotide to affect chromatin and DNA methylation of imprinting control regions. *Molecular Cell* **44** 361–372. (doi:10.1016/j.molcel.2011.08.032)
- Reis e Silva AR, Bruno C, Fleuret R, Daniel N, Archilla C, Peynot N, Lucci CM, Beaujean N & Duranthon V 2012 Alteration of DNA demethylation dynamics by *in vitro* culture conditions in rabbit pre-implantation embryos. *Epigenetics* **7** 440–446. (doi:10.4161/epi.19563)
- Ribeiro-Mason K, Boulesteix C, Brochard V, Aguirre-Lavin T, Salvaing J, Fleuret R, Adenot P, Maalouf WE & Beaujean N 2012 Nuclear dynamics of histone H3 trimethylated on lysine 9 and/or phosphorylated on serine 10 in mouse cloned embryos as new markers of reprogramming? *Cellular Reprogramming* **14** 283–294. (doi:10.1089/cell.2011.0071)
- Rouleau J, Tanigawa G & Szyf M 1992 The mouse DNA methyltransferase 5'-region. A unique housekeeping gene promoter. *Journal of Biological Chemistry* **267** 7368–7377.
- Rugg-Gunn PJ, Cox BJ, Ralston A & Rossant J 2010 Distinct histone modifications in stem cell lines and tissue lineages from the early mouse embryo. *PNAS* **107** 10783–10790. (doi:10.1073/pnas.0914507107)
- Saha B, Home P, Ray S, Larson M, Paul A, Rajendran G, Behr B & Paul S 2013 EED and KDM6B coordinate the first mammalian cell lineage commitment to ensure embryo implantation. *Molecular and Cellular Biology* **33** 2691–2705. (doi:10.1128/MCB.00069-13)
- Saitou M, Barton SC & Surani MA 2002 A molecular programme for the specification of germ cell fate in mice. *Nature* **418** 293–300. (doi:10.1038/nature00927)
- Sakaue M, Ohta H, Kumaki Y, Oda M, Sakaide Y, Matsuoka C, Yamagiwa A, Niwa H, Wakayama T & Okano M 2010 DNA methylation is dispensable for the growth and survival of the extraembryonic lineages. *Current Biology* **20** 1452–1457. (doi:10.1016/j.cub.2010.06.050)
- Santenard A, Ziegler-Birling C, Koch M, Tora L, Bannister AJ & Torres-Padilla ME 2010 Heterochromatin formation in the mouse embryo requires critical residues of the histone variant H3.3. *Nature Cell Biology* **12** 853–862. (doi:10.1038/ncb2089)
- Santos F, Hendrich B, Reik W & Dean W 2002 Dynamic reprogramming of DNA methylation in the early mouse embryo. *Developmental Biology* **241** 172–182. (doi:10.1006/dbio.2001.0501)
- Santos F, Peters AH, Otte AP, Reik W & Dean W 2005 Dynamic chromatin modifications characterise the first cell cycle in mouse embryos. *Developmental Biology* **280** 225–236. (doi:10.1016/j.ydbio.2005.01.025)
- Sarmento OF, Digilio LC, Wang Y, Perlin J, Herr JC, Allis CD & Coonrod SA 2004 Dynamic alterations of specific histone modifications during early murine development. *Journal of Cell Science* **117** 4449–4459. (doi:10.1242/jcs.01328)
- Sato M, Kimura T, Kurokawa K, Fujita Y, Abe K, Masuhara M, Yasunaga T, Ryo A, Yamamoto M & Nakano T 2002 Identification of PGC7, a new gene expressed specifically in preimplantation embryos and germ cells. *Mechanisms of Development* **113** 91–94. (doi:10.1016/S0925-4773(02)00002-3)
- Saxonov S, Berg P & Brutlag DL 2006 A genome-wide analysis of CpG dinucleotides in the human genome distinguishes two distinct classes of promoters. *PNAS* **103** 1412–1417. (doi:10.1073/pnas.0510310103)
- Schrode N, Xenopoulos P, Piliszek A, Frankenberg S, Plusa B & Hadjantonakis AK 2013 Anatomy of a blastocyst: cell behaviors driving cell fate choice and morphogenesis in the early mouse embryo. *Genesis* **51** 219–233. (doi:10.1002/dvg.22368)

- Schultz RM 2002 The molecular foundations of the maternal to zygotic transition in the preimplantation embryo. *Human Reproduction Update* **8** 323–331. (doi:10.1093/humupd/8.4.323)
- Schultz GA, Hahnel A, Arcellana-Panlilio M, Wang L, Goubau S, Watson A & Harvey M 1993 Expression of IGF ligand and receptor genes during preimplantation mammalian development. *Molecular Reproduction and Development* **35** 414–420. (doi:10.1002/mrd.1080350416)
- Shao GB, Ding HM & Gong AH 2008 Role of histone methylation in zygotic genome activation in the preimplantation mouse embryo. *In Vitro Cellular & Developmental Biology. Animal* **44** 115–120. (doi:10.1007/s11626-008-9082-4)
- Shen L, Inoue A, He J, Liu Y, Lu F & Zhang Y 2014 Tet3 and DNA replication mediate demethylation of both the maternal and paternal genomes in mouse zygotes. *Cell Stem Cell* **15** 459–470. (doi:10.1016/j.stem.2014.09.002)
- Seutels F, Tjon G, Ludwig T & Barlow DP 2003 Imprinted silencing of Slc22a2 and Slc22a3 does not need transcriptional overlap between Igf2r and Air. *EMBO Journal* **22** 3696–3704. (doi:10.1093/emboj/cdg341)
- Smallwood SA, Tomizawa S, Krueger F, Ruf N, Carli N, Segonds-Pichon A, Sato S, Hata K, Andrews SR & Kelsey G 2011 Dynamic CpG island methylation landscape in oocytes and preimplantation embryos. *Nature Genetics* **43** 811–814. (doi:10.1038/ng.864)
- Smith ZD, Chan MM, Mikkelsen TS, Gu H, Gnirke A, Reggev A & Meissner A 2012 A unique regulatory phase of DNA methylation in the early mammalian embryo. *Nature* **484** 339–344. (doi:10.1038/nature10960)
- Stein P, Worrad DM, Belyaev ND, Turner BM & Schultz RM 1997 Stage-dependent redistributions of acetylated histones in nuclei of the early preimplantation mouse embryo. *Molecular Reproduction and Development* **47** 421–429. (doi:10.1002/(SICI)1098-2795(199708)47:4<421::AID-MRD8>3.0.CO;2-M)
- Strahl BD & Allis CD 2000 The language of covalent histone modifications. *Nature* **403** 41–45. (doi:10.1038/47412)
- Sui X, Price C, Li Z & Chen J 2012 Crosstalk between DNA and histones: Tet's new role in embryonic stem cells. *Current Genomics* **13** 603–608. (doi:10.2174/138920212803759730)
- Suo L, Meng Q, Pei Y, Fu X, Wang Y, Bunch TD & Zhu S 2010 Effect of cryopreservation on acetylation patterns of lysine 12 of histone H4 (acH4K12) in mouse oocytes and zygotes. *Journal of Assisted Reproduction and Genetics* **27** 735–741. (doi:10.1007/s10815-010-9469-5)
- Takada S, Paulsen M, Tevendale M, Tsai CE, Kelsey G, Cattanaach BM & Ferguson-Smith AC 2002 Epigenetic analysis of the Dlk1-Gtl2 imprinted domain on mouse chromosome 12: implications for imprinting control from comparison with Igf2-H19. *Human Molecular Genetics* **11** 77–86. (doi:10.1093/hmg/11.1.77)
- Tan M, Luo H, Lee S, Jin F, Yang JS, Montellier E, Buchou T, Cheng Z, Rousseaux S, Rajagopal N *et al.* 2011 Identification of 67 histone marks and histone lysine crotonylation as a new type of histone modification. *Cell* **146** 1016–1028. (doi:10.1016/j.cell.2011.08.008)
- Tanaka M, Hennebold JD, Macfarlane J & Adashi EY 2001 A mammalian oocyte-specific linker histone gene H1oo: homology with the genes for the oocyte-specific cleavage stage histone (cs-H1) of sea urchin and the B4/H1M histone of the frog. *Development* **128** 655–664.
- Torres-Padilla ME, Parfitt DE, Kouzarides T & Zernicka-Goetz M 2007 Histone arginine methylation regulates pluripotency in the early mouse embryo. *Nature* **445** 214–218. (doi:10.1038/nature05458)
- Tsukada Y, Fang J, Erdjument-Bromage H, Warren ME, Borchers CH, Tempst P & Zhang Y 2006 Histone demethylation by a family of JmjC domain-containing proteins. *Nature* **439** 811–816. (doi:10.1038/nature04433)
- VerMilyea MD, O'Neill LP & Turner BM 2009 Transcription-independent heritability of induced histone modifications in the mouse preimplantation embryo. *PLoS ONE* **4** e6086. (doi:10.1371/journal.pone.0006086)
- Wang J, Zhang M, Zhang Y, Kou Z, Han Z, Chen DY, Sun QY & Gao S 2010 The histone demethylase JMJD2C is stage-specifically expressed in preimplantation mouse embryos and is required for embryonic development. *Biology of Reproduction* **82** 105–111. (doi:10.1095/biolreprod.109.078055)
- Wang J, Xie G, Singh M, Ghanbarian AT, Rasko T, Szvetnik A, Cai H, Besser D, Prigione A, Fuchs NV *et al.* 2014 Primate-specific endogenous retrovirus-driven transcription defines naive-like stem cells. *Nature* **516** 405–409. (doi:10.1038/nature13804)
- Watanabe D, Suetake I, Tada T & Tajima S 2002 Stage- and cell-specific expression of Dnmt3a and Dnmt3b during embryogenesis. *Mechanisms of Development* **118** 187–190. (doi:10.1016/S0925-4773(02)00242-3)
- Watson AJ, Damsky CH & Kidder GM 1990 Differentiation of an epithelium: factors affecting the polarized distribution of Na⁺,K⁺-ATPase in mouse trophoblast. *Developmental Biology* **141** 104–114. (doi:10.1016/0012-1606(90)90105-R)
- Wongtawan T, Taylor JE, Lawson KA, Wilmut I & Pennings S 2011 Histone H4K20me3 and HP1 α are late heterochromatin markers in development, but present in undifferentiated embryonic stem cells. *Journal of Cell Science* **124** 1878–1890. (doi:10.1242/jcs.080721)
- Wossidlo M, Arand J, Sebastiano V, Lepikhov K, Boiani M, Reinhardt R, Scholer H & Walter J 2010 Dynamic link of DNA demethylation, DNA strand breaks and repair in mouse zygotes. *EMBO Journal* **29** 1877–1888. (doi:10.1038/emboj.2010.80)
- Wossidlo M, Nakamura T, Lepikhov K, Marques CJ, Zakhartchenko V, Boiani M, Arand J, Nakano T, Reik W & Walter J 2011 5-Hydroxymethylcytosine in the mammalian zygote is linked with epigenetic reprogramming. *Nature Communications* **2** 241. (doi:10.1038/ncomms1240)
- Wu FR, Liu Y, Shang MB, Yang XX, Ding B, Gao JG, Wang R & Li WY 2012 Differences in H3K4 trimethylation in *in vivo* and *in vitro* fertilization mouse preimplantation embryos. *Genetics and Molecular Research* **11** 1099–1108. (doi:10.4238/2012.April.27.9)
- Yamanaka Y, Ralston A, Stephenson RO & Rossant J 2006 Cell and molecular regulation of the mouse blastocyst. *Developmental Dynamics* **235** 2301–2314. (doi:10.1002/dvdy.20844)
- Yan L, Yang M, Guo H, Yang L, Wu J, Li R, Liu P, Lian Y, Zheng X, Yan J *et al.* 2013 Single-cell RNA-Seq profiling of human preimplantation embryos and embryonic stem cells. *Nature Structural & Molecular Biology* **20** 1131–1139. (doi:10.1038/nsmb.2660)
- Yeap LS, Hayashi K & Surani MA 2009 ERG-associated protein with SET domain (ESET)-Oct4 interaction regulates pluripotency and represses the trophoblast lineage. *Epigenetics & Chromatin* **2** 12. (doi:10.1186/1756-8935-2-12)
- Young LE & Beaujean N 2004 DNA methylation in the preimplantation embryo: the differing stories of the mouse and sheep. *Animal Reproduction Science* **82–83** 61–78.
- Zaitseva I, Zaitsev S, Alenina N, Bader M & Krivokharchenko A 2007 Dynamics of DNA-demethylation in early mouse and rat embryos developed *in vivo* and *in vitro*. *Molecular Reproduction and Development* **74** 1255–1261. (doi:10.1002/mrd.20704)
- Zhang K, Haversat JM & Mager J 2013a CTR9/PAF1c regulates molecular lineage identity, histone H3K36 trimethylation and genomic imprinting during preimplantation development. *Developmental Biology* **383** 15–27. (doi:10.1016/j.ydbio.2013.09.005)
- Zhang K, Dai X, Wallingford MC & Mager J 2013b Depletion of Suda3 reveals an essential role in early lineage specification. *Developmental Biology* **373** 359–372. (doi:10.1016/j.ydbio.2012.10.026)
- Zhou L, Wang P, Zhang J, Heng BC & Tong GQ 2015 ING2 (inhibitor of growth protein-2) plays a crucial role in preimplantation development. *Zygote* (In press). (doi:10.1017/S0967199414000768)
- Zuo X, Sheng J, Lau HT, McDonald CM, Andrade M, Cullen DE, Bell FT, Iacovino M, Kyba M, Xu G *et al.* 2012 Zinc finger protein ZFP57 requires its co-factor to recruit DNA methyltransferases and maintains DNA methylation imprint in embryonic stem cells via its transcriptional repression domain. *Journal of Biological Chemistry* **287** 2107–2118. (doi:10.1074/jbc.M111.322644)

Received 17 April 2015

First decision 11 May 2015

Revised manuscript received 26 May 2015

Accepted 1 June 2015