Histone H3 lysine 27 trimethylation acts as an epigenetic barrier in porcine nuclear reprogramming

Bingteng Xie*, Heng Zhang*, Renyue Wei, Qiannan Li, Xiaogang Weng, Qingran Kong and Zhonghua Liu

Laboratory of Embryo Biotechnology, College of Life Science, Northeast Agricultural University, No. 59 Mucai Street, Harbin, 150030 Heilongjiang, China

Correspondence should be addressed to Z Liu; Email: liu086@126.com

*(B Xie and H Zhang contributed equally to this work)

Abstract

Aberrant epigenetic reprogramming is the main obstacle to the development of somatic cell nuclear transfer (SCNT) embryos and the generation of induced pluripotent stem (iPS) cells, which results in the low reprogramming efficiencies of SCNT and iPS. Histone H3 lysine 27 trimethylation (H3K27me3), as a repressive epigenetic mark, plays important roles in mammalian development and iPS induction. However, the reprogramming of H3K27me3 in pig remains elusive. In this study, we showed that H3K27me3 levels in porcine early cloned embryos were higher than that in IVF embryos. Then GSK126 and GSK-J4, two small molecule inhibitors of H3K27me3 methylase (EZH2) and demethylases (UTX/JMJD3), were used to regulate the H3K27me3 level. The results showed that H3K27me3 level was reduced in cloned embryos after treatment of PEF with 0.75 μ M GSK126 for 48 h, incubation of one-cell reconstructed oocytes with 0.1 μ M GSK126 and injection of antibody for EZH2 into oocyte. Meanwhile, the development of the cloned embryos was significantly improved after these treatments. On the contrary, GSK-J4 treatment increased the H3K27me3 level in cloned embryos and decreased the cloned embryonic development. Furthermore, iPS efficiency was both increased after reducing the H3K27me3 level in donor cells and in early reprogramming phase. In summary, our results suggest that H3K27me3 acts as an epigenetic barrier in SCNT and iPS reprogramming, and reduction of H3K27me3 level in donor cells and in early reprogramming phase can enhance both porcine SCNT and iPS efficiency.

Reproduction (2016) **151** 9–16

Introduction

Somatic cell nuclear transfer (SCNT) and induced pluripotent stem (iPS) cells are two of the primary routes to reprogram differentiated cells back to pluripotent state. Both of the two reprogramming methods have great significance: SCNT holds great potential for agriculture, biomedical industry and endangered species conservation as this technique allows the generation of a whole organism from a single differentiated somatic cell (Yang *et al.* 2007b). iPS cells can serve as a valuable patient specific cell source for *in vitro* disease modeling and cell/tissue replacement therapies because they do not have ethical issues (Park *et al.* 2008). However, the efficiencies of SCNT and iPS induction are extremely low, which limit their practical use. For example, the overall reproductive cloning efficiency in all species is remarkably low (1–5%) except bovine species (5–20%) (Rodriguez-Osorio *et al.* 2012). And in general, the efficiency of iPS colony formation with the four Yamanaka's factors (*Oct4*, *Sox2*, *Klf4* and *c-Myc*; OSKM)

is less than 1% of the starting population (Silva *et al.* 2008).

Aberrant epigenetic reprogramming is believed to be the main reason for the low reprogramming efficiencies of SCNT and iPS (Mason *et al.* 2012, Buganim *et al.* 2013). It could occur on many aspects, including DNA methylation and histone modifications (Dean *et al.* 2001, Kang *et al.* 2001, Santos *et al.* 2003, Ohgane *et al.* 2004). For example, cloned embryos always show higher DNA methylation level compared to their fertilized counterparts (Dean *et al.* 2001, Beaujean *et al.* 2004, Yang *et al.* 2007a). And more recently, H3K9me3 was identified as a critical epigenetic barrier in SCNT reprogramming. The reprogramming resistant regions (RRRs) that could not be activated in cloned embryos as that in fertilized embryos were enriched for H3K9me3 in donor cells, and its removal by deletion of H3K9me3 methylase in donor cells and by ectopically expressed H3K9me3 demethylase in cloned embryos not only reactivated the majority of RRRs, but greatly improved SCNT efficiency as well (Matoba *et al.* 2014).

Similarly, abnormal epigenetics reprogramming also often occur in these areas during iPS induction, and the reprogramming efficiency could be improved by regulating these epigenetic status both in SCNT and iPS (Kishigami *et al.* 2006, Huangfu *et al.* 2008, Onder *et al.* 2012, Huan *et al.* 2013, Matoba *et al.* 2014).

Histone H3 lysine 27 trimethylation (H3K27me3) is another repressive epigenetic mark. It is catalyzed by enhancer of zeste homolog 2 (EZH2), and can be removed by H3K27me3-specific demethylase of Jumonji domain containing protein 3 (JMJD3) and ubiquitously transcribed tetratricopeptide repeat X (UTX) (Bogliotti & Ross 2012). And EZH2 and JMJD3/UTX can be specifically suppressed by two small molecule inhibitors of GSK126 and GSK-J4 respectively (Kruidenier *et al.* 2012, McCabe *et al.* 2012). H3K27me3 has a great effect on embryonic development. For example, the embryonic development was reduced after inhibition of H3K27me3 removal in bovine (Canovas *et al.* 2012). Similarly, iPS reprogramming process was also accompanied by genome wide H3K27me3 changes, and the aberrant dynamics of H3K27me3 would impair iPS cells induction (Mikkelsen *et al.* 2008, Hawkins *et al.* 2010, Mansour *et al.* 2012). Recently, defective reprogramming of H3K27me3 has been observed in cloned mouse embryos (Zhang *et al.* 2009). However, the reprogramming of H3K27me3 has not been reported in other species of cloned embryos.

In this study, we investigated the reprogramming of H3K27me3 in porcine cloned embryos, and assessed the impact of H3K27me3 level on SCNT and iPS efficiency through regulating H3K27me3 level by treatment of small molecules GSK126 and GSK-J4 respectively. We believe the results would advance the understanding of reprogramming mechanism of SCNT and iPS.

Materials and methods

The reagents and media used in our study were purchased from Life Technologies, R&D, Millipore and Bioind (Kibbutz Beit Haemek, Israel) unless otherwise stated.

Animals

The PEF lines were derived from Landrace which were from Northeast Agricultural University Embryo Engineering Laboratory Experimental Pig Base. ICR mice for mouse embryonic fibroblast cells were purchased from Vital River Company, Beijing, China. The porcine ovaries were obtained from DaZhongRouLian Slaughterhouse Company (Harbin, Heilongjiang, China). All studies adhered to procedures consistent with the Northeast Agriculture University of Biological Sciences Guide for the care and use of laboratory animals.

SCNT, parthenogenetic activation and IVF

The procedure for porcine SCNT was the same as we described previously (Liu *et al.* 2008). After 42 h of maturation culture, the oocytes were treated with 1 mg/ml hyaluronidase to remove the surrounding cumulus cells. Oocytes with a clearly extruded first polar body were selected as recipient cytoplasts. Cumulus free oocytes were enucleated by aspirating the first polar body and adjacent cytoplasm with a glass pipette 25 µm in diameter in TCM199-Hepes plus 0.3% BSA and 7.5 µg/ml cytochalasin B. A single donor cell was injected into the perivitelline space and electrically fused using two direct pulses of 120 V/mm for 30 µs in fusion medium. Fused eggs were cultured in porcine zygote medium-3 (PZM-3) medium for 6.5 days in an atmosphere of 5% CO₂ and 95% air at 39 °C. The cleavage and blastocyst rates were assessed at 48 and 156 h after activation, and the number of blastocyst cells was examined by nuclear staining with 5 µg/ml Hoechst 33342. For anti-EZH2 group, ten picoliters of EZH2 antibody (3147, Cell signaling) was injected into matured oocytes before NT. After injection, oocytes were kept for at least 2 h before manipulations, which allows the antibody to bind endogenous EZH2. Cumulus free oocytes were directly activated by the same parameters as SCNT procedure for producing parthenogenetic activation (PA) embryos.

For IVF, freshly ejaculated sperm rich fractions were collected from fertile boars, and following a short incubation at 39 °C, the semen was resuspended and washed three times in DPBS supplemented with 0.1% (w/v) BSA by centrifugation at 1500g for 4 min. The spermatozoa concentration was measured using a hemocytometer, and the proportion of motile sperm was determined. The spermatozoa were diluted with modified Tris-buffered medium (mTBM) to an optimal concentration. Cumulus free oocytes were washed three times in mTBM. Approximately 30 oocytes were inseminated in 50-µl drops of mTBM at a final sperm concentration of 3×10^5 /ml for 6 h. Then embryos were washed and cultured in PZM-3 medium for 6.5 days in an atmosphere of 5% CO₂ and 95% air at 39 °C. SCNT and IVF embryos, one-cell, two-cell, four-cell and blastocyst (B) stages, were collected at 6/12, 24, 48 and 144 h respectively.

Immunofluorescence analysis

PEF, sperm, oocytes and embryos without zona pellucidas were washed twice in PBS, then fixed in freshly prepared 4% paraformaldehyde in PBS, permeabilized in 1% Triton X-100 in PBS, and left in blocking solution (1% BSA in PBS) for 1 h. For immunolabeling, the embryos were incubated overnight at 4 °C with primary antibody for H3K27me3 (07449, Millipore) washed three times and incubated for 1 h with secondary

antibody, FITC-labeled donkey anti-rabbit IgG (A21206, Invitrogen), diluted 1:1000 with blocking solution. Samples were washed and stained with 5 µg/ml of Hoechst 33342. Fluorescence was detected and imaged using a Nikon fluorescence microscope. The average optical intensity (AOI) of the H3K27me3 staining and the number of PEF cells were calculated using the freely available Image J software. The AOI of H3K27me3 nuclear area staining in embryos were normalized by subtracting the AOI of the background areas.

Western blot

PEF or embryos removed zona pellucida, stored at -80°C , were lysed with cell lysis buffer for western (Beyotime, Jiangsu, China). Lysates were separated in SDS-PAGE for the routine western blot assay and transferred to PVDF membranes (Invitrogen); nonspecific binding was blocked by incubation in 1% BSA in PBS at room temperature for 1 h. Blots were then probed for overnight at 4°C with primary antibody for H3K27me3 (07449, Millipore). β -Actin (A1978, Sigma) served as loading control. After 1 h incubation at room temperature with secondary antibodies, protein bands were detected by HRP Chemiluminescent Kit (Invitrogen) and Kodak BioMax Light film. The average optical intensity of the H3K27me3 and β -actin bands were calculated by Image J.

iPSCs generation

Inducible iPSCs were generated as described previously (Wang *et al.* 2013). Briefly, GP293T cells were transfected with the pMX plasmids which contained mouse *Oct4*, *Sox2*, *Klf4*, *c-Myc*, *Nr5a2* and *Tbx3* together with packaging plasmids VSVG. The medium was replaced 12 h after transfection, and the virus supernatants were harvested after subsequent 24 h. PEF at passage 3 or 4 in 24-well plate dishes at a density of 10^4 cells/dish were incubated with filtrated viral supernatants containing 8 µg/ml polybrene. The infection medium was replaced after 24 h with PEF medium. When the confluence of PEF was 90%, we passed them into 12-well plate as 1:8, and the medium was replaced with MX medium (which was developed in our Lab for porcine iPSCs culture) everyday until the mouse ESCs like colonies appeared at around seven days after infection (Wang *et al.* 2013). The colonies were picked and were cultured with MX medium. And then, the line was passaged with tryp LE (Gibco, Grand Island, US) digestion.

Alkaline phosphatase staining

iPSCs was fixed with 4% paraformaldehyde at room temperature for 1 min and washed three times with

PBS. Alkaline phosphatase (AP) staining was performed with BCIP/NBT Alkaline Phosphatase Color Development Kit (Beyotime) following the manufacturer's instructions.

GSK126 and GSK-J4 treatment

PEF cells (2×10^5 per well) were seeded into six well tissue culture plates 24 h before treatment. Cells were then exposed to 0.1% DMSO and varying concentrations of GSK126 and GSK-J4 (range 0–3 µM) for 48 h respectively, then the cells were harvested for SCNT or iPS induction. During cloned and PA embryonic development 0.1% DMSO, 0.1 µM GSK126 and GSK-J4 incubated one-cell embryos for 24 h respectively. During iPS induction, 0.75 µM GSK126 incubated PEF for 2–7 days after infection respectively.

Statistical analysis

Differences of data (mean $\pm$ S.E.M.) were analyzed by SPSS statistical software 19.0. The data were analyzed using one way ANOVA. For all analyses, differences were considered to be statistically significant when $P < 0.05$.

Results

The H3K27me3 levels of early cloned embryos were higher than that of IVF embryos

The global H3K27me3 levels were determined at varying stages of IVF and cloned embryos (Fig. 1A). In pronuclear stage of IVF embryos (6 and 12 h), H3K27me3 signal was found in female pronucleus (red arrowhead). Since the late stage of one-cell, H3K27me3 staining intensity was dramatically reduced and faint in two-cell, four-cell and blastocyst stage. In cloned embryos, similar to IVF embryos, H3K27me3 level gradually decreased with the cleavage, and became faint after four-cell stage. However, in one-cell and two-cell cloned embryos, H3K27me3 levels were higher than that of IVF embryos (Fig. 1A and C).

Beside that, the H3K27me3 level of PEF cells, sperm and oocyte were also determined. The results showed that H3K27me3 could be detected on metaphase chromosomes of MII oocytes but not in sperm. And PEF possessed strong H3K27me3 signal in nucleus (Fig. 1B). In order to compare the H3K27me3 level of PEF and oocyte more clearly, nuclear transfer was performed without enucleation, and H3K27me3 was detected within 1 h after the fusion. It could be observed clearly that H3K27me3 level of the nucleus derived from PEF (red arrowhead) was obviously higher than that of maternal chromosomes (white arrows) (Fig. 1B).

Figure 1 The H3K27me3 levels of early cloned embryos and IVF embryos (A) The distribution of H3K27me3 in cloned and IVF embryos: two-cell (6 h/12 h), one-cell, four-cell, and blastocyst. Red arrowheads refer to the female pronucleus in IVF one-cell embryos. Bar, 50 μ m. Results are representative of about 20 embryos for each stage. (B) The distribution of H3K27me3 in oocyte, sperm, PEF cells and 1 h cloned embryo without enucleation. Red arrowhead refers to the nucleus derived from PEF, and white arrows refer to maternal chromosomes. Bar, 50 μ m. (C) The western blot analysis of H3K27me3 in two-cell of IVF and cloned embryos. One hundred embryos for each group were analyzed. The below image is relative quantity of H3K27me3 standardized to β -actin.

Improvement of cloned embryonic development by reducing the H3K27me3 level in donor PEF cells

In order to regulate the H3K27me3 level in donor cells, PEF were treated with GSK126 and GSK-J4 for 48 h. It was shown that treatment with GSK126 in the dosage more than 0.5 μ M significantly reduced the H3K27me3 level in PEF cells (Supplementary Figure 1, see section on supplementary data given at the end of this article, Fig. 2). And interestingly, when treated with 0.75 μ M GSK126, the blastocyst rate of PEF cells was significantly higher than that of control group (DMSO vs GSK126 0.75 μ M and 21.99 vs 31.82%, $P < 0.05$; Table 1). Meanwhile, the H3K27me3 level was lower in one-cell and two-cell of treatment group than that of control (Fig. 3A and C). On the other hand, treatment with

GSK-J4 in the dosage more than 0.5 μ M increased the H3K27me3 level in PEF cells (Supplementary Figure 1, Fig. 2). However, there was no significant difference in the blastocyst rate between 0.75 μ M GSK-J4 treatment group and control group (DMSO vs GSK-J4 0.75 μ M and 21.99 vs 20.93%, $P > 0.05$) (Table 1). The H3K27me3 level was higher in one-cell but not in two-cell of treatment group than that of control group (Fig. 3A and C).

Improvement of cloned embryonic development by reducing the H3K27me3 level in early embryos

We also incubated one-cell cloned embryos with GSK126 and GSK-J4 to regulate H3K27me3 level.

Figure 2 The H3K27me3 level of PEF cells incubated with 0.75 μ M GSK126 and GSK-J4 for 48 h respectively. (A) Immunofluorescence analysis of H3K27me3 level in PEF cells incubated with 0.75 μ M GSK126 and GSK-J4 for 48 h respectively. Bar, 100 μ m. (B) Western blot of PEF cells incubated with 0.75 μ M GSK126 and GSK-J4 for 48 h respectively. The western blot was repeated three times. (C) Relative quantity of H3K27me3 standardized to β -actin. Different superscript indicates $P < 0.05$.

Table 1 The effect of treatment of donor PEF cells with GSK126 or GSK-J4 on cloned embryonic development

Groups	Replicates	Embryos	Cleavage (%)	Blastocyst (%)	No. of blastocyst cells
DMSO	4	162	133 (82.42 ± 2.87)	35 (21.99 ± 1.54) ^a	36.29 ± 12.65
GSK126 0.75 µM	4	192	167 (86.86 ± 5.73)	61 (31.82 ± 4.04) ^b	35.00 ± 16.92
GSK-J4 0.75 µM	4	211	176 (83.44 ± 1.30)	44 (20.93 ± 2.29) ^a	35.91 ± 8.84

Different superscript indicates $P < 0.05$.

The results showed that there were no significant effects on parthenogenetic embryonic development after incubating one-cell with 0.1 µM GSK126 or GSK-J4 for 24 h (Supplementary Table 1). When one-cell cloned embryos were incubated with 0.1 µM GSK126 for 24 h, the H3K27me3 level in two-cell of treatment group were significantly lower than that of control group (Fig. 3B and D). Meanwhile, the blastocyst rate of treatment group was significantly higher than that of control group (DMSO vs GSK126 0.1 µM, 21.99 vs 31.33%, $P < 0.05$; Table 2). On the contrary, when one-cell cloned embryos were incubated with 0.1 µM GSK-J4 for 24 h, the H3K27me3 level in two-cell of treatment group was significantly higher than that of control group (Fig. 3B and D). The blastocyst rate of treatment group was significant lower than that of control group (DMSO vs GSK-J4 0.1 µM, 21.99 vs 8.05%, $P < 0.05$; Table 2).

In order to verify the influence of H3K27me3 level on the cloned embryonic development, we injected EZH2 antibody in oocyte to interfere EZH2 function.

Consistent with GSK126 treatment, the H3K27me3 level in two-cell of anti-EZH2 group was significantly lower than that of control group (Fig. 3B and D). Meanwhile, the blastocyst rate of anti-EZH2 group was significantly higher than that of the control group (IgG vs anti-EZH2 and 20.60 vs 30.63%, $P < 0.05$; Supplementary Table 2).

Improvement of iPS efficiency by reducing the H3K27me3 level in donor cells or during early reprogramming phase

We also examined the iPS efficiency after reducing H3K27me3 level. After incubation with 0.75 µM GSK126 for 72 h, PEF cells (PEF1) were infected by *Oct4*, *Sox2*, *Klf4*, *c-Myc*, *Nr5a2*, and *Tbx3* (m-pMX-OSKMNT). On the 8th day after infection, the numbers of AP-positive colonies were detected. The results showed that the AP-positive colonies of GSK126 treatment group were about 4 times more than that of

Figure 3 The H3K27me3 level of one-cell and two-cell cloned embryos treated with GSK126 and GSK-J4 respectively (A) The H3K27me3 level of one-cell and two-cell cloned embryos derived from PEF cells incubated with 0.75 µM GSK126 and GSK-J4 for 48 h respectively. (B) The H3K27me3 level of two-cell cloned embryos after treating one-cell embryos with 0.1 µM GSK126 and GSK-J4 for 24 h respectively. Images shown are representative of about 20 embryos for each group. Bar, 50 µm. (C and D) The normalized AOI of H3K27me3 nuclear area staining in one-cell and two-cell cloned embryos. Different superscript in the same figure indicates $P < 0.05$.

Table 2 The effect of GSK126 or GSK-J4 treatment on cloned embryonic development

Groups	Replicates	Embryos	Cleavage (%)	Blastocyst (%)	No. of blastocyst cells
DMSO	3	89	81 (91.03 ± 3.78) ^a	18 (20.27 ± 3.74) ^b	37.70 ± 15.67
GSK126 0.1 μM	3	102	90 (88.26 ± 2.75) ^a	32 (31.33 ± 2.60) ^a	33.20 ± 13.18
GSK-J4 0.1 μM	3	87	125 (71.26 ± 5.26) ^b	7 (8.05 ± 5.26) ^c	33.60 ± 14.01

Different superscript in same array indicates $P < 0.05$.

control group ($P < 0.05$; Fig. 4A and B). When PEF1 was incubated with 0.75 μM GSK126 for 2–7 days after infection respectively, the results showed that incubation with GSK126 enhanced iPS induction, which achieved the highest efficiency by incubating for 4 days ($P < 0.05$; Fig. 4A and B). We validated the results on the other PEF line (PEF2), and the results showed that GSK126 treatment group had more AP-positive colonies than DMSO treatment group ($P < 0.05$; Fig. 4C), which were similar with that in PEF1.

Discussion

Epigenetic modifications are important factors that affect the cellular reprogramming efficiencies of SCNT and iPS. But few specific epigenetic locus have been evaluated carefully to determine its role on reprogramming. In this study, our results suggest H3K27me3 acts as an epigenetic barrier in porcine nuclear reprogramming, and reduction of H3K27me3 level in donor cells and in early reprogramming phase can enhance both porcine SCNT and iPS efficiency. This will advance our understanding of the reprogramming mechanisms of both SCNT and iPS.

H3K27me3 is an important repressive epigenetic mark and its level changes dynamically during mammalian embryonic development (Ross *et al.* 2008, Park *et al.* 2009, Zhang *et al.* 2009, 2012, Gao *et al.* 2010). Here we showed that in porcine IVF embryos, H3K27me3 level decreased drastically after fertilization, which is consistent with previous works (Park *et al.* 2009, Gao *et al.* 2010); this was believed to be necessary for normal embryonic genome activation. However, in cloned embryos H3K27me3 levels were significantly higher in one-cell and two-cell cloned embryos compared to that in IVF embryos. And this should be closely related with higher H3K27me3 levels in donor PEF cells indicating that H3K27me3 was failed to be reprogrammed normally in porcine cloned embryos, and thus impaired expression of some genes and the development potential of cloned porcine embryos.

Various abnormal epigenetic modifications in cloned embryos have been found. In a recent study, H3K9me3 was identified as a critical epigenetic barrier in murine SCNT reprogramming, which impeded cloned embryonic development. And the SCNT efficiency was greatly improved after reducing H3K9me3 level. Similarly, in this study, we regulated H3K27me3 level by GSK126 and GSK-J4, which have been used in several studies

(McCabe *et al.* 2012, Rao *et al.* 2015). The results manifested that when the H3K27me3 level of cloned embryos were reduced, the embryonic development was significantly improved, including incubation of

Figure 4 Improvement of iPS efficiency by reducing the H3K27me3 level in donor cells or during early reprogramming phase (A) the numbers of AP-positive colonies. -3: PEF1 cells were incubated with 0.75 μM GSK126 for 3 days, and then infected by *Oct4*, *Sox2*, *Klf4*, *c-Myc*, *Nr5a2*, and *Tbx3* (OSKMNT). 0, +2 - +7: PEF1 cells were incubated with 0.75 μM GSK126 for 0, 2 to 7 days after infection. Data represent mean and s.e.m. of 3 replicates experiments; * $P < 0.05$. (B) Representative images of AP-staining of -3, 0, +4 groups. (C) The numbers of AP-positive colonies. -3: PEF2 cells were incubated with DMSO and GSK126 (0.75 μM) respectively for 3 days, and then infected by OSKMNT. +4: PEF2 cells were incubated with DMSO and GSK126 (0.75 μM) respectively for 4 days after infection. Data represent mean and s.e.m. of 3 replicates experiments. * $P < 0.05$.

PEF/one-cell with GSK126 and injection of EZH2 antibody into oocyte. On the contrary, when the H3K27me3 level of cloned embryos was increased, the cloned embryonic development was impaired. This demonstrated that high H3K27me3 level is detrimental to porcine embryonic development, and H3K27me3 acts as an epigenetic barrier to porcine SCNT reprogramming. We can speculate that H3K27me3 enriched domain may include genes important for embryonic development, the remained high level of H3K27me3 after SCNT would certainly impede the development. So when H3K27me3 levels were artificially down regulated, the expressions of these genes could be derepressed, as a result, the development of cloned embryos could be improved. However, the improvement of embryonic development could be due to the changes of complicated gene networks, rather than a single gene. For example, in the recent study, it was found that numerous candidate genes expressed differently after regulation of H3K9me3, but most of these genes function were unclear (Matoba *et al.* 2014). So it was great difficulty, if possible, to determine which genes were responsible for the improvement of embryonic development.

It had been reported that regulating epigenetic status can improve the reprogramming efficiency both in SCNT and iPS (Kishigami *et al.* 2006, Huangfu *et al.* 2008, Onder *et al.* 2012, Huan *et al.* 2013, Matoba *et al.* 2014). Therefore, we also examined the iPS efficiency after reducing H3K27me3 level. As expected, we found that iPS induction efficiency could be enhanced by reducing the H3K27me3 level in donor cells and in early reprogramming phase. This suggests that H3K27me3 also acts as an epigenetic barrier to porcine iPS reprogramming. Previous works found that genome wide H3K27me3 changes were accompanied with iPS reprogramming process, and removal of H3K27me3 was required for activation of some genes expression during iPS induction (Hawkins *et al.* 2010). In this research, during porcine iPS induction, the expressions of these genes might also be facilitated by reduction of H3K27me3 level, and in turn enhance the iPS reprogramming.

In summary, our results suggest that H3K27me3 acts as an epigenetic barrier in porcine SCNT and iPS reprogramming, and reduction of H3K27me3 level in donor cells and in early reprogramming phase can enhance both porcine SCNT and iPS efficiency. In the future, it is exciting to determine whether our results could be generally applied to other animal species, even human.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/REP-15-0338>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by grants from the National Basic Research Program of China (2011CB944202, 2011CBA01006) and the National Natural Science Foundation of China (J1210069, 31470079).

References

- Beaujean N, Taylor J, Gardner J, Wilmut I, Meehan R & Young L 2004 Effect of limited DNA methylation reprogramming in the normal sheep embryo on somatic cell nuclear transfer. *Biology of Reproduction* **71** 185–193. (doi:10.1095/biolreprod.103.026559)
- Bogliotti YS & Ross PJ 2012 Mechanisms of histone H3 lysine 27 trimethylation remodeling during early mammalian development. *Epigenetics* **7** 976–981. (doi:10.4161/epi.21615)
- Buganim Y, Faddah DA & Jaenisch R 2013 Mechanisms and models of somatic cell reprogramming. *Nature Review Genetics* **14** 427–439. (doi:10.1038/nrg3473)
- Canovas S, Cibelli JB & Ross PJ 2012 Jumoni domain-containing protein 3 regulates histone 3 lysine 27 methylation during bovine preimplantation development. *PNAS* **109** 2400–2405. (doi:10.1073/pnas.1119112109)
- Dean W, Santos F, Stojkovic M, Zakhartchenko V, Walter J, Wolf E & Reik W 2001 Conservation of methylation reprogramming in mammalian development: aberrant reprogramming in cloned embryos. *PNAS* **98** 13734–13738. (doi:10.1073/pnas.241522698)
- Gao Y, Hyttel P & Hall VJ 2010 Regulation of H3K27me3 and H3K4me3 during early porcine embryonic development. *Molecular Reproduction and Development* **77** 540–549. (doi:10.1002/mrd.21180)
- Hawkins RD, Hon GC, Lee LK, Ngo Q, Lister R, Pelizzola M, Edsall LE, Kuan S, Luu Y, Klugman S *et al.* 2010 Distinct epigenomic landscapes of pluripotent and lineage-committed human cells. *Cell Stem Cell* **6** 479–491. (doi:10.1016/j.stem.2010.03.018)
- Huan YJ, Zhu J, Xie BT, Wang JY, Liu SC, Zhou Y, Kong QR, He HB & Liu ZH 2013 Treating cloned embryos, but not donor cells, with 5-aza-2'-deoxycytidine enhances the developmental competence of porcine cloned embryos. *Journal of Reproduction and Development* **59** 442–449. (doi:10.1262/jrd.2013-026)
- Huangfu D, Maehr R, Guo W, Eijkelenboom A, Snitow M, Chen AE & Melton DA 2008 Induction of pluripotent stem cells by defined factors is greatly improved by small-molecule compounds. *Nature Biotechnology* **26** 795–797. (doi:10.1038/nbt1418)
- Kang YK, Koo DB, Park JS, Choi YH, Chung AS, Lee KK & Han YM 2001 Aberrant methylation of donor genome in cloned bovine embryos. *Nature Genetics* **28** 173–177. (doi:10.1038/88903)
- Kishigami S, Mizutani E, Ohta H, Hikichi T, Thuan NV, Wakayama S, Bui HT & Wakayama T 2006 Significant improvement of mouse cloning technique by treatment with trichostatin A after somatic nuclear transfer. *Biochemical and Biophysical Research Communications* **340** 183–189. (doi:10.1016/j.bbrc.2005.11.164)
- Kruidenier L, Chung CW, Cheng Z, Liddle J, Che K, Joberty G, Bantscheff M, Bountra C, Bridges A, Diallo H *et al.* 2012 A selective jumoni H3K27 demethylase inhibitor modulates the proinflammatory macrophage response. *Nature* **488** 404–408. (doi:10.1038/nature11262)
- Liu ZH, Song J, Wang ZK, Tian JT, Kong QR, Zheng Z, Yin Z, Gao L, Ma HK, Sun S *et al.* 2008 Green fluorescent protein (GFP) transgenic pig produced by somatic cell nuclear transfer. *Chinese Science Bulletin* **53** 1035–1039. (doi:10.1007/s11434-008-0168-x)
- Mansour AA, Gafni O, Weinberger L, Zviran A, Ayyash M, Rais Y, Krupalnik V, Zerbib M, Amann-Zalcenstein D, Maza I, Hanna JH *et al.* 2012 The H3K27 demethylase Utx regulates somatic and germ cell epigenetic reprogramming. *Nature* **488** 409–413. (doi:10.1038/nature11272)

- Mason K, Liu Z, Aguirre-Lavin T & Beaujean N 2012 Chromatin and epigenetic modifications during early mammalian development. *Animal Reproduction Science* **134** 45–55. (doi:10.1016/j.anireprosci.2012.08.010)
- Matoba S, Liu Y, Lu F, Iwabuchi KA, Shen L, Inoue A & Zhang Y 2014 Embryonic Development following Somatic Cell Nuclear Transfer Impeded by Persisting Histone Methylation. *Cell* **159** 884–895. (doi:10.1016/j.cell.2014.09.055)
- McCabe MT, Ott HM, Ganji G, Korenchuk S, Thompson C, Van Aller GS, Liu Y, Graves AP, Della Pietra A III, Diaz E *et al.* 2012 EZH2 inhibition as a therapeutic strategy for lymphoma with EZH2-activating mutations. *Nature* **492** 108–112. (doi:10.1038/nature11606)
- Mikkelsen TS, Hanna J, Zhang X, Ku M, Wernig M, Schorderet P, Bernstein BE, Jaenisch R, Lander ES & Meissner A 2008 Dissecting direct reprogramming through integrative genomic analysis. *Nature* **454** 49–55. (doi:10.1038/nature07056)
- Ohgane J, Wakayama T, Senda S, Yamazaki Y, Inoue K, Ogura A, Marh J, Tanaka S, Yanagimachi R & Shiota K 2004 The Sall3 locus is an epigenetic hotspot of aberrant DNA methylation associated with placentomegaly of cloned mice. *Genes to Cells* **9** 253–260. (doi:10.1111/j.1356-9597.2004.00720.x)
- Onder TT, Kara N, Cherry A, Sinha AU, Zhu N, Bernt KM, Cahan P, Marcacci BO, Unternaehrer J, Gupta PB *et al.* 2012 Chromatin-modifying enzymes as modulators of reprogramming. *Nature* **483** 598–602. (doi:10.1038/nature10953)
- Park IH, Arora N, Huo H, Maherali N, Ahfeldt T, Shimamura A, Lensch MW, Cowan C, Hochedlinger K & Daley GQ 2008 Disease-specific induced pluripotent stem cells. *Cell* **134** 877–886. (doi:10.1016/j.cell.2008.07.041)
- Park KE, Magnani L & Cabot RA 2009 Differential remodeling of mono- and trimethylated H3K27 during porcine embryo development. *Molecular Reproduction and Development* **76** 1033–1042. (doi:10.1002/mrd.21061)
- Rao RA, Dhele N, Cheemadan S, Ketkar A, Jayandharan GR, Palakodeti D & Rampalli S 2015 Ezh2 mediated H3K27me3 activity facilitates somatic transition during human pluripotent reprogramming. *Scientific Reports* **5** 8229. (doi:10.1038/srep08229)
- Rodriguez-Orsorio N, Urrego R, Cibelli JB, Eilertsen K & Memili E 2012 Reprogramming mammalian somatic cells. *Theriogenology* **78** 1869–1886. (doi:10.1016/j.theriogenology.2012.05.030)
- Ross PJ, Ragina NP, Rodriguez RM, Lager AE, Siripattaraprat K, Lopez-Corrales N & Cibelli JB 2008 Polycomb gene expression and histone H3 lysine 27 trimethylation changes during bovine preimplantation development. *Reproduction* **136** 777–785. (doi:10.1530/REP-08-0045)
- Santos F, Zakhartchenko V, Stojkovic M, Peters A, Jenuwein T, Wolf E, Reik W & Dean W 2003 Epigenetic marking correlates with developmental potential in cloned bovine preimplantation embryos. *Current Biology* **13** 1116–1121. (doi:10.1016/S0960-9822(03)00419-6)
- Silva J, Barrandon O, Nichols J, Kawaguchi J, Theunissen TW & Smith A 2008 Promotion of reprogramming to ground state pluripotency by signal inhibition. *PLoS Biology* **6** e253. (doi:10.1371/journal.pbio.0060253)
- Wang J, Gu Q, Hao J, Jia Y, Xue B, Jin H, Ma J, Wei R, Hai T, Kong Q *et al.* 2013 Tbx3 and Nr5a2 play important roles in pig pluripotent stem cells. *Stem Cell Reviews* **9** 700–708. (doi:10.1007/s12015-013-9439-2)
- Yang J, Yang S, Beaujean N, Niu Y, He X, Xie Y, Tang X, Wang L, Zhou Q & Ji W 2007a Epigenetic marks in cloned rhesus monkey embryos: comparison with counterparts produced *in vitro*. *Biology of Reproduction* **76** 36–42. (doi:10.1095/biolreprod.106.051383)
- Yang X, Smith SL, Tian XC, Lewin HA, Renard JP & Wakayama T 2007b Nuclear reprogramming of cloned embryos and its implications for therapeutic cloning. *Nature Genetics* **39** 295–302. (doi:10.1038/ng1973)
- Zhang M, Wang F, Kou Z, Zhang Y & Gao S 2009 Defective chromatin structure in somatic cell cloned mouse embryos. *Journal of Biological Chemistry* **284** 24981–24987. (doi:10.1074/jbc.M109.011973)
- Zhang A, Xu B, Sun Y, Lu X, Gu R, Wu L, Feng Y & Xu C 2012 Dynamic changes of histone H3 trimethylated at positions K4 and K27 in human oocytes and preimplantation embryos. *Fertility and Sterility* **98** 1009–1016. (doi:10.1016/j.fertnstert.2012.06.034)

Received 23 July 2015

First decision 25 August 2015

Revised manuscript received 23 October 2015

Accepted 29 October 2015