Progesterone effects on mouse sperm kinetics in conditions of viscosity

S Pérez-Cerezales¹, A P López-Cardona^{1,2} and A Gutiérrez-Adán¹

¹Departamento de Reproducción Animal, INIA, Avda. Puerta de Hierro, Madrid, Spain and ²Grupo de Investigación en Biogénesis, Universidad de Antioquia, Antioquia, Colombia

Correspondence should be addressed to S Pérez-Cerezales; Email: perez.serafin@inia.es

Abstract

The spermatozoa delivered to the female genital tract need to swim towards the oocyte through viscous secretions. Once close to the oocyte, the spermatozoa are guided by a gradient of progesterone (P_4) and other unknown chemoattractants via a process known as chemotaxis. Using polyvinylpyrrolidone to establish the conditions of viscosity, we examined the response of mouse spermatozoa to P_4 . Herein, we show that in low-viscous media, P_4 induces hyperactive-like motility whereby sperm show erratic trajectories and non-progressive movement. However, an opposite response is produced in viscous medium in that trajectories are linear and motility is more progressive and less erratic. Our observations provide a behavioural explanation for the chemotaxis of spermatozoa swimming under viscous conditions in a spatial gradient of the chemoattractant P_4 . They also highlight the importance of using viscous solutions to mimic *in vivo* conditions when analysing sperm behaviour in response to any stimulus.

Reproduction (2016) **151** 501–507

Introduction

Once in the female genital tract, to reach the fertilization site, spermatozoa need to tackle several anatomical structures, each with specific and complex physicochemical properties (Suarez & Pacey 2006). Active mechanisms will guide the spermatozoa to the proximity of the oocyte (Eisenbach & Giojalas 2006). So far, three such guidance mechanisms have been proposed: temperature gradient (thermotaxis), against fluid flow (rheotaxis) and chemoattractant(s) gradient (chemotaxis). All the three mechanisms are functional in the oviduct (see Pérez-Cerezales *et al.* (2015) for a review).

A feature shared by the three guidance mechanisms is that spermatozoa show a hyperactivation-like motility (Armon & Eisenbach 2011, Miki & Clapham 2013, Boryshpolets *et al.* 2015). Hyperactivation is a property of capacitated spermatozoa that, under low-viscous conditions, involves vigorous movements produced by asymmetrical and high-amplitude waves in the flagella, resulting in erratic swimming trajectories (Suarez 2008). Currently, the model suggested for spermatozoa swimming in a spatial gradient of temperature (Boryshpolets *et al.* 2015) or chemoattractant (Gakamsky *et al.* 2009, Armon & Eisenbach 2011) is similar to that described for bacterial chemotaxis (Macnab & Koshland 1972). According to this model, a spermatozoon swimming in a gradient towards a lower temperature or lower chemoattractant concentration responds by increasing the frequency of hyperactivation-like events and turns to modify its direction of swimming. Conversely, when

swimming towards a higher temperature or chemoattractant concentration, this constant stimulation causes the suppression of turns and hyperactive-like events. If no changes are perceived, the spermatozoon recovers its non-stimulated swimming mode, consisting of a straighter movement pattern with occasional hyperactivation events and turns. Although the model has been empirically validated for chemotaxis and thermotaxis, the behavioural response of spermatozoa to progesterone (P_4) as the main known chemoattractant has shown some discrepancies with the model. In studies addressing the response of a population of human spermatozoa to P_4 (1–10 nM), enhanced hyperactivation-like motility events have been observed (Uhlér *et al.* 1992, Gakamsky *et al.* 2009). This argues against the proposed model as increasing concentrations of chemoattractant should have a damping effect on hyperactivation-like motility. In an elegant series of experiments, Armon *et al.* (2011) showed that hyperactive-like motility of human spermatozoa was effectively reduced in the sperm subpopulation swimming towards higher concentrations of P_4 . However, the physiological meaning of P_4 's main effect of increasing the hyperactive-like motility of the general sperm population remains unknown.

It has been established that CatSper mediates P_4 -induced Ca^{2+} influx (Lishko *et al.* 2011, Strünker *et al.* 2011) as well as hyperactivation (Carlson *et al.* 2003) in human spermatozoa. However, studies conducted so far have failed to detect CatSper activation by P_4 in mouse spermatozoa (Lishko *et al.* 2011), neither P_4 -mediated chemotaxis (Chang *et al.* 2013). It has been shown in

mice that the initial Ca^{2+} influx provoked by P_4 occurs at the base of the sperm head (Fukami *et al.* 2003), whereas CatSper activation initiates Ca^{2+} influx in the main stretch of the flagellum (Xia *et al.* 2007). Moreover, P_4 elicits transient Ca^{2+} influx in spermatozoa from *Catsper1*-null mice (Ren *et al.* 2001), suggesting that P_4 might activate another Ca^{2+} influx pathway besides CatSper in mammalian sperm. It has been reported that P_4 fails to induce Ca^{2+} fluctuations in *Plcδ4*-null mice (Fukami *et al.* 2003), suggesting PLC pathway involvement in this response, possibly via Ca^{2+} mobilization from internal Ca^{2+} stores at the base of the sperm head through the IP3 receptor (Darszon *et al.* 2011). Thus, given the localization of the P_4 membrane receptor in mouse spermatozoa (Pietrobon *et al.* 2003), its activation by P_4 might be the mechanism mediating the hyperactivation response, perhaps through the PLC pathway.

Interestingly, it has been noted that the behaviour of human spermatozoa in response to temperature changes varies under conditions of viscosity or low viscosity. Under viscous conditions considered closer to those of the physiological environment, the temperature-dependent hyperactive-like behaviour observed in low-viscous conditions changes to one showing a higher frequency of turns when the temperature is reduced (Boryshpolets *et al.* 2015). This indicates the different behaviour of spermatozoa in response to a stimulus depending on the viscosity of the surrounding medium. Indeed, hyperactivation-like motility in low-viscous conditions has been proposed to be the manifestation of a more efficient motility mode for sperm propagation under viscous conditions in hamster (Suarez *et al.* 1991) and mouse spermatozoa (Suarez & Dai 1992). Thus, the hyperactive-like motility shown by spermatozoa swimming in low-viscous conditions in response to P_4 could also indicate a more efficient swimming mode under physiological conditions of viscosity.

This study was designed to address the inconsistencies observed in the P_4 chemotaxis model by determining the behaviour of mouse spermatozoa in response to this chemoattractant under conditions simulating the viscosity of the physiological environment of the female genital tract.

Materials and methods

Reagents

All media components were purchased from Sigma–Aldrich, except where otherwise stated.

Spermatozoa

Experiments with mice were carried out in strict accordance with recommendations of the guidelines of European Community Council Directive 86/609/EEC. Every effort was made to

minimize suffering. Experiments were approved by the Committee on the Ethics of Animal Experiments of the INIA (permit number CEEA 2014/025).

For sperm analysis, F1 generation males from crosses between C57BL/6J (B6) female × DBA/2J (D2) male mice were used. The animals were kept in conditions of a 14 h light:10 h darkness cycle with no food or water restrictions. At the age of 4 months, males were individually isolated for 2 weeks before killing by cervical dislocation. Spermatozoa were collected from the caudal epididymis and suspended in human tubal fluid (HTF) medium (2.04 mM $\text{CaCl}_2 \times 2\text{H}_2\text{O}$, 101.6 mM NaCl, 4.69 mM KCl, 0.37 mM KH_2PO_4 , 0.2 mM $\text{MgSO}_4 \times 7\text{H}_2\text{O}$, 21.4 mM sodium lactate, 0.33 mM sodium pyruvate, 2.78 mM glucose, 25 mM NaHCO_3 , 100 U/mL penicillin, 50 µg/mL streptomycin SO_4 and 0.001% (w/v) phenol red, supplemented with 1% BSA and stabilized overnight at 37°C in a 5% CO_2 atmosphere). Once collected, the sperm samples were incubated for 1 h under an atmosphere of 5% CO_2 at 37°C for capacitation.

Sperm processing and motility

After capacitation, spermatozoa were diluted to a final concentration of $\sim 2 \times 10^6/\text{mL}$ in HTF with or without polyvinylpyrrolidone (PVP) of average MW360000 at 4, 6 or 7.5% (w/v). For each experiment with P_4 , 20 µL of P_4 or DMSO (control) pre-diluted in HTF (1:10) were added to 100 µL of spermatozoa diluted in HTF with or without PVP. Motility was then determined within 20 s of adding P_4 or DMSO by placing 20 µL of sperm suspension onto a pre-warmed slide on the stage heated to 37°C of a Nikon Eclipse 50i microscope (Nikon) fitted with a digital camera Basler A312f (Basler AG, Ahrensburg, Germany) capable of recording 25 frames/s. In each repetition, five videos of 1 s (20–60 moving spermatozoa) were recorded in different fields on the top of the drop and analysed using the Integrated Semen Analysis System (ISAS; Projectes i Serveis R+D S.L., Valencia, Spain). In each experiment, three replicates for each condition/treatment were set up. The parameters analysed were as described by Mortimer *et al.* (2000): straight-line velocity (VSL; time-averaged velocity of the sperm head along a straight line from its first position to its last position, expressed in µm/s); curvilinear velocity (VCL; time-averaged velocity of the sperm head along its actual curvilinear path, expressed in µm/s); average path velocity (VAP; velocity over an average path generated by a roaming average between frames, expressed in µm/s); linearity (LIN) (defined as $(\text{VSL}/\text{VCL}) \times 100$); straightness (STR) (defined as $(\text{VSL}/\text{VAP}) \times 100$); wobble (WOB) (defined as $(\text{VAP}/\text{VCL}) \times 100$); amplitude of lateral head (ALH) displacement (width of the lateral movement of the sperm head, expressed in µm) and beat-cross frequency (BCF; number of times the sperm head crosses the direction of movement per second, expressed in Hz).

Acrosome reaction

The method used was based on acrosome staining using *Pisum sativum* lectin conjugated with FITC (PSA–FITC) following a standard protocol as described previously (Lybaert *et al.* 2009) with minor modifications. Briefly, P_4 at 0.015 pM,

15 pM, 15 nM, 15 μ M or the same volume of DMSO (as a negative control) was added to capacitated spermatozoa and these were immediately processed for acrosome staining. In parallel, spermatozoa were diluted in HTF with or without 6% PVP (w/v) and also immediately processed for acrosome staining. As a positive control, spermatozoa were incubated for 30 min with 10 μ M of the ionophore A23187 to induce the acrosome reaction. The spermatozoa were washed twice in PBS by centrifugation (1 min at 500 *g*) and subsequently smeared on a microscope glass slide and air-dried for 15 min. Next, the slides were immersed in absolute methanol for 30 s, air-dried and rinsed in PBS twice for 5 min before incubation with PSA-FITC (15 μ g/mL in PBS) in a humid box for 30 min. Finally, the slides were washed with distilled water for 15 min and mounted with an aqueous mounting medium (Fluoromount G; EMS, Hatfield, PA, USA). Slides were examined in a fluorescence microscope and numbers of acrosome-reacted and non-acrosome-reacted spermatozoa were counted by randomly moving across different fields of the slide (counting 100 cells per slide). For each treatment, experiments were run in triplicate using two sperm samples from two different animals.

Statistical analysis

All statistical tests were performed using the software package SPSS Statistics 22.0 for Windows (IBM, Armonk, NY, USA). Results are provided as means $\pm$ S.E.M. Means were compared and analysed by three-way ANOVA followed by Tukey's *post hoc* test. Significance was set at $P < 0.05$.

Results

To establish the swimming pattern of mouse spermatozoa in viscous conditions, we examined sperm kinetics at four concentrations of PVP (0 (low-viscous medium), 4, 6 and 7.5% (viscous media)). Differences in most of the variables recorded were detected between the viscous media and low-viscous medium. Velocities (VCL, VSL and VAP) decreased with increasing percentages of PVP in the medium ($P < 0.05$, three-way ANOVA; Fig. 1A). The velocity ratios LIN and STR were only affected at the highest PVP concentration (7.5%; $P < 0.05$, three-way ANOVA; Fig. 1B), indicating equally proportional reductions in the three velocities when spermatozoa swam in a medium containing 4 and 6% PVP. The other velocity ratio analysed, WOB, did not vary with PVP concentration. In addition, viscosity effects were also observed on motility patterns such that in the viscous media, the lateral displacement of the cell was reduced in an inverse manner with respect to the PVP concentration, reflected as a reduction in BCF and ALH with increasing PVP percentage ($P < 0.05$, three-way ANOVA; Fig. 1C). Accordingly, PVP led to the reduced velocity of spermatozoa and reduced side-to-side movement of the head and frequency of this movement, leading to a slower, more regular swimming trajectory (Fig. 1D). In the absence of

PVP, the swimming pattern was erratic, including more irregular side-to-side head movements.

We then went on to assess the dose-dependent effect of P_4 on spermatozoa in low-viscous vs viscous medium. For these experiments, 6% PVP was selected as the highest concentration for which relationships among the three velocities defined were unaffected, determining that STR, WOB and LIN remained unchanged. This means that 6% PVP does not affect the STR of the sperm swimming trajectory and that any change in sperm behaviour could be attributed to their response to P_4 . Moreover, this percentage of PVP did not induce the acrosome reaction (Fig. 2), which could in itself affect the behaviour of the spermatozoa.

In low-viscous medium, P_4 reduced the LIN of sperm movement indicated by reduced values of LIN, STR and WOB ($P < 0.05$, three-way ANOVA; Fig. 3). These effects were already observed at the lowest P_4 concentration tested ($P < 0.05$, three-way ANOVA). Such changes produced in LIN, STR and WOB (Mortimer 2000) reflect the effect of P_4 on sperm, provoking a slight increase in VCL and slight decreases in VSL and VAP (these effects alone were insignificant; three-way ANOVA; Fig. 3). In addition, 15 nM and 15 μ M concentrations of P_4 led to a greater ALH ($P < 0.05$, three-way ANOVA), whereas no effects were observed on BCF. These observations indicate that the vigorous motility induced by P_4 consisted of a greater sperm head

Figure 1 Sperm kinetics in viscous media. (A) Velocities: curvilinear (VCL), straight line (VSL) and average path (VAP). (B) Velocity ratios: linearity (LIN), straightness (STR) and wobble (WOB). (C) Head movements: beat-cross frequency (BCF, expressed in Hz) and amplitude of lateral head displacement (ALH, expressed in μ m). (A, B and C) Different letters indicate significant differences between percentages of PVP according to three-way ANOVA ($P < 0.05$). Mean $\pm$ S.E.M., $n = 5$ determinations; each determination is the average of measurements in five fields on 21–34 spermatozoa per field. (D) Sperm trajectories tracked in 1 s in the presence of different concentrations of PVP (from left to right: 0, 4, 6 and 7.5%).

Figure 2 Acrosome-reacted spermatozoa visualized with *Pisum sativum* agglutinin conjugated to FITC (PSA-FITC) under different conditions. A23187 was used as a positive control to induce the acrosome reaction. 6% PVP was used for the viscosity conditions. As a negative control for progesterone (P_4 ; concentration 0), DMSO (solvent) was added instead of P_4 at the same volume. Varying concentrations of P_4 were used to induce the acrosome reaction (0.015 pM, 15 pM, 15 nM and 15 μ M). * $P < 0.05$ and ** $P < 0.02$ according to three-way ANOVA. Mean $\pm$ S.E.M., $n = 8$ determinations; each determination is the average of measurements in 52–60 spermatozoa.

displacement distance but did not affect the frequency at which the head crossed the swimming trajectory. These changes in sperm kinetics are characteristic of a hyperactive-like motility, as described by others for human spermatozoa in response to P_4 (Uhler *et al.* 1992, Gakamsky *et al.* 2009). P_4 at 15 pM, 15 nM and 15 μ M induced the acrosome reaction (Fig. 2; $P < 0.05$, three-way ANOVA), meaning that the motility changes induced by P_4 were accompanied by the release of acrosome contents.

In the viscous medium (6% PVP), contrary effects of P_4 to those produced in the low-viscous medium were observed. The P_4 dose-dependent curves between low viscosity vs viscosity were significantly different for VSL, VAP, LIN, STR, WOB and ALH ($P < 0.01$, three-way ANOVA; Fig. 3). Thus, sperm velocities (VSL and VAP) were increased (in response to 15 nM and 15 μ M P_4 ; $P < 0.05$, three-way ANOVA) and movement became linear as indicated by LIN and WOB ratios with increasing P_4 concentrations (significant for 15 nM and 15 μ M at $P < 0.05$, three-way ANOVA) (Fig. 3). Moreover, P_4 led to a higher BCF (significant for 15 μ M; $P < 0.05$, three-way ANOVA), whereas ALH was unaltered. These results suggest that the P_4 -induced non-progressive sperm motility (hyperactivation-like) observed in

low-viscous medium is equivalent to the progressive motility observed under the more physiological conditions of the viscous media.

Discussion

Once in the female genital tract, spermatozoa have to swim in different mucous secretions whose viscosity differs according to anatomical site (Jansen 1978, Smith *et al.* 2009) and oestrous cycle stage (Jansen 1978, Miki & Clapham 2013). Thus, the averaged viscosity of mouse uterus and oviduct is 81 and 1.8 cP respectively (Miki & Clapham 2013), in the range of viscosity obtained with the percentages of PVP (Hyakutake *et al.* 2015) used in our study. The effects of viscosity on mouse sperm kinetics observed here including a reduced velocity but more regular trajectory are in agreement with observations in hamster (Suarez *et al.* 1991), mice (Suarez & Dai 1992) and cattle (Hyakutake *et al.* 2015). The different behaviour shown by spermatozoa in low-viscous compared with viscous media (Fig. 4A) suggests a need to study those factors determining the sperm motility mode under the latter conditions. P_4 is among the most studied modulators of sperm function (Baldi *et al.* 2009). This hormone is secreted by cumulus cells and is a physiological sperm chemoattractant (Teves *et al.* 2006, Oren-Benaroya *et al.* 2008). In agreement with earlier data for human spermatozoa (Uhler *et al.* 1992, Gakamsky *et al.* 2009), under low-viscous conditions, we also observed hyperactivation-like motility in mouse spermatozoa in response to acute P_4 treatment. Thus, P_4 reduced LIN, STR and WOB and increased lateral displacement of the head (ALH; Figs 3 and 4A). As a consequence of these motility changes, the P_4 -activated spermatozoa acquired a more erratic swimming pattern, moving from side to side with higher amplitude (Fig. 4A). This motility type is known to facilitate spermatozoon progression under viscous conditions (Suarez *et al.* 1991, Suarez & Dai 1992). Remarkably, under viscous conditions, we observed the opposite effects of P_4 on sperm kinetics (Figs 3 and 4A). Thus, sperm LIN and BCF increased, whereas head displacement was reduced (Figs 3 and 4A). These results indicate that, contrary to what occurs in low-viscous media, the motility type induced by P_4 is one of the propagation under viscous conditions. Therefore, it seems logical that the known non-progressive motility mode induced by P_4 is actually an artefact of analysing the response under low-viscous conditions. This means that spermatozoa behave as expected for the model suggested for sperm chemotaxis when swimming in a spatial gradient of the chemoattractant P_4 (Armon & Eisenbach 2011). Thereby, when a spermatozoon swims towards a higher concentration of P_4 , its motility becomes linearized, propagating the cell towards the higher concentrations (Fig. 4B). In contrast, a spermatozoon swimming

Figure 3 Sperm kinetics in response to progesterone (P₄) in low-viscous vs viscous medium. Different letters indicate significant differences between P₄ concentrations in each viscosity condition according to three-way ANOVA ($P < 0.05$). Asterisks indicate significant differences between the slopes of the kinetic parameters for low viscosity vs viscosity according to three-way ANOVA (* $P < 0.05$ and ** $P < 0.001$). Mean $\pm$ S.E.M., viscous medium ($n = 10$ determinations); low-viscous medium ($n = 7$ determinations). Each determination is the average of measurements in five fields (20–41 spermatozoa per field).

towards lower concentrations of P₄ would show a more circular trajectory until higher concentrations of P₄ linearized its motility again.

Chang *et al.* (2007) did not succeed in detecting spatial chemotaxis in response to gradient of P₄ in mouse spermatozoa when using conditions of low viscosity. Our results provide an explanation to this observation, as only under viscous conditions, the spermatozoa would migrate towards a P₄ gradient in a chemotaxis spatial assay. Accordingly, the chemotaxis of human spermatozoa to P₄ was reported using a chemotaxis spatial assay under viscous conditions (3.5% PVP 25 K; Armon & Eisenbach 2011). This demonstrates the importance of using conditions closer to physiological when studying the behavioural response of spermatozoa to any kind of stimuli. Complementary studies would include analyses of the response to P₄ under other rheological conditions of the surrounding medium such as the viscoelasticity that characterize some fluids present along the female genital tract (Suarez & Dai 1992). It has been reported that at similar viscosity, the VSL and flagellar beat frequency of cattle spermatozoa are higher in non-Newtonian fluids leading to a more linear motility type (Hyakutake *et al.* 2015) compared with Newtonian fluids. Accordingly, it has been shown that under viscoelastic conditions, hyperactive samples of mouse spermatozoa linearize the motility, whereas non-hyperactive spermatozoa behave in an opposite way (Suarez & Dai 1992).

Our results also show that at a certain concentration (higher than 15 pM), P₄ induces the acrosome reaction (Fig. 2). This observation is consistent with the dual effect of P₄ on human spermatozoa, whereby it acts as a chemoattractant at lower concentrations and induces the acrosome reaction at higher concentrations (Uhler

et al. 1992). This release of acrosome contents could be related to the P₄ gradient existing within the cumulus cells (Teves *et al.* 2006). Thus, we propose that P₄ could linearize motility when sperm swimming up the P₄ gradient penetrate the cumulus and this could simultaneously provoke the release of acrosome contents to facilitate this penetration (Fig. 4B). This dual effect of P₄ could only apply to certain sperm populations, as recently described in human spermatozoa (Uñates *et al.* 2014). Accordingly, the acrosome-reacting population would eliminate the barrier formed by the cumulus cells and the chemotactic population would progress towards the oocyte by acquiring a more linear path in response to the P₄ gradient. It is also possible that both acrosome-reacting and chemotactic populations are comprised in part by the same spermatozoa. Indeed, it seems that most fertilizing spermatozoa undergo the acrosome reaction before reaching the zona pellucida of cumulus-enclosed oocytes (Jin *et al.* 2011). This would be an interesting case of altruism between 'sister' cells that needs to be further explored to extend the concept of cooperation between spermatozoa of a single ejaculate. Despite no descriptions to date in the laboratory mouse, heterogeneity among different sperm populations of the same ejaculate and cooperation phenomena have been observed in the wood mouse *Apodemus sylvaticus* (Moore *et al.* 2002) and species of *Peromyscus* (Fisher & Hoekstra 2010).

In conclusion, our findings suggest that the hyperactive-like motility induced by P₄ in mouse spermatozoa swimming in low-viscous media is a manifestation of the chemotactic behaviour of spermatozoa that propagates them towards a chemoattractant gradient in viscous conditions. In view of the dramatically different responses to P₄ produced in spermatozoa swimming in

Figure 4 Models of the behavioural response of mouse spermatozoa to progesterone (P_4). (A) Trajectories of mouse spermatozoa swimming in viscous or low-viscous medium with or without P_4 . In low-viscous medium, the trajectory is more irregular than in viscous medium and P_4 induces erratic motility. Under viscosity conditions, the trajectory is more regular and the addition of P_4 linearizes motility facilitating sperm progression. (B) Dual effect of P_4 on spermatozoa swimming in viscous medium towards the oocyte. (1) The trajectory becomes linear when spermatozoa swim towards higher P_4 concentrations and is therefore guided by chemotaxis to the oocyte. (2) Once the spermatozoon reaches the granulosa, the higher concentration of P_4 induces the acrosome reaction, facilitating penetration of the oocyte.

low-viscous or viscous media, we propose the viscosity of the medium is an essential factor to consider in any study designed to address sperm behaviour.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was funded by Grant AGL2012-39652-C02-01 awarded by the Spanish Ministry of Economy and Competitiveness. S Pérez-Cereales was supported by a postdoctoral grant from the same ministry. A P López-Cardona was supported by COLCIENCIAS (Francisco José de Caldas fellowship 512/2010).

References

- Armon L & Eisenbach M 2011 Behavioral mechanism during human sperm chemotaxis: involvement of hyperactivation. *PLoS ONE* **6** e28359. (doi:10.1089/scd.2015.0322)
- Baldi E, Luconi M, Muratori M, Marchiani S, Tamburrino L & Forti G 2009 Nongenomic activation of spermatozoa by steroid hormones: facts and fictions. *Molecular and Cellular Endocrinology* **308** 39–46. (doi:10.2741/3793)
- Boryshpolets S, Pérez-Cereales S & Eisenbach M 2015 Behavioral mechanism of human sperm in thermotaxis: a role for hyperactivation. *Human Reproduction* **30** 884–892. (doi:10.1038/srep16146)
- Carlson AE, Westenbroek RE, Quill T, Ren D, Clapham DE, Hille B, Garbers DL & Babcock DF 2003 CatSper1 required for evoked Ca^{2+} entry and control of flagellar function in sperm. *Proceedings of the National Academy of Sciences* **100** 14864–14868. (doi:10.1073/pnas.2536658100)
- Chang H, Kim BJ, Kim YS, Suarez SS & Wu M 2013 Different migration patterns of sea urchin and mouse sperm revealed by a microfluidic chemotaxis device. *PLoS ONE* **8** e60587. (doi:10.1371/journal.pone.0060587)
- Darszon A, Nishigaki T, Beltran C & Treviño CL 2011 Calcium channels in the development, maturation, and function of spermatozoa. *Physiological Reviews* **91** 1305–1355. (doi:10.1016/j.ydbio.2014.07.017)
- Eisenbach M & Giojalas LC 2006 Sperm guidance in mammals – an unpaved road to the egg. *Nature Reviews. Molecular Cell Biology* **7** 276–285. (doi:10.1074/jbc.M115.690396)
- Fisher HS & Hoekstra HE 2010 Competition drives cooperation among closely related sperm of deer mice. *Nature* **463** 801–803. (doi:10.1038/nature08736)
- Fukami K, Yoshida M, Inoue T, Kurokawa M, Fissore RA, Yoshida N, Mikoshiba K & Takenawa T 2003 Phospholipase Cdelta4 is required for Ca^{2+} mobilization essential for acrosome reaction in sperm. *Journal of Cell Biology* **161** 79–88. (doi:10.1083/jcb.200210057)
- Gakamsky A, Armon L & Eisenbach M 2009 Behavioral response of human spermatozoa to a concentration jump of chemoattractants or intracellular cyclic nucleotides. *Human Reproduction* **24** 1152–1163. (doi:10.1093/humrep/den409)
- Hyakutake T, Suzuki H & Yamamoto S 2015 Effect of non-Newtonian fluid properties on bovine sperm motility. *Journal of Biomechanics* **48** 2941–2947. (doi:10.1016/j.jbiomech.2015.08.005)
- Jansen RP 1978 Fallopian tube isthmus mucus and ovum transport. *Science* **201** 349–351. (doi:10.1126/science.580814)
- Jin M, Fujiwara E, Kakiuchi Y, Okabe M, Satouh Y, Baba SA, Chiba K & Hirohashi N 2011 Most fertilizing mouse spermatozoa begin their acrosome reaction before contact with the zona pellucida during in vitro fertilization. *Proceedings of the National Academy of Sciences* **108** 4892–4896. (doi:10.1073/pnas.1018202108)
- Lishko PV, Botchkina IL & Kirichok Y 2011 Progesterone activates the principal Ca^{2+} channel of human sperm. *Nature* **471** 387–391. (doi:10.1038/nature09767)
- Lybaert P, Danguy A, Leleux F, Meuris S & Lebrun P 2009 Improved methodology for the detection and quantification of the acrosome reaction in mouse spermatozoa. *Histology and Histopathology* **24** 999–1007.
- Macnab RM & Koshland DE 1972 The gradient-sensing mechanism in bacterial chemotaxis. *Proceedings of the National Academy of Sciences* **69** 2509–2512. (doi:10.1073/pnas.69.9.2509)
- Miki K & Clapham DE 2013 Rheotaxis guides mammalian sperm. *Current Biology* **23** 443–452. (doi:10.1016/j.cub.2013.02.007)
- Moore H, Dvoráková K, Jenkins N & Breed W 2002 Exceptional sperm cooperation in the wood mouse. *Nature* **418** 174–177. (doi:10.1038/nature00832)
- Mortimer ST 2000 CASA – practical aspects. *Journal of Andrology* **21** 515–524.
- Oren-Benaroya R, Orvieto R, Gakamsky A, Pinchasov M & Eisenbach M 2008 The sperm chemoattractant secreted from human cumulus cells is progesterone. *Human Reproduction* **23** 2339–2345. (doi:10.1093/humrep/den265)
- Pérez-Cereales S, Boryshpolets S & Eisenbach M 2015 Behavioral mechanisms of mammalian sperm guidance. *Asian Journal of Andrology* **17** 628–632. (doi:10.1038/srep16146)

- Pietrobon EO, de los Angeles Monclus M, Alberdi AJ & Fornés MW** 2003 Progesterone receptor availability in mouse spermatozoa during epididymal transit and capacitation: ligand blot detection of progesterone-binding protein. *Journal of Andrology* **24** 612–620. (doi:10.1002/j.1939-4640.2003.tb02713.x)
- Ren D, Navarro B, Perez G, Jackson AC, Hsu S, Shi Q, Tilly JL & Clapham DE** 2001 A sperm ion channel required for sperm motility and male fertility. *Nature* **413** 603–609. (doi:10.1038/35098027)
- Smith DJ, Gaffney EA, Gadêlha H, Kapur N & Kirkman-Brown JC** 2009 Bend propagation in the flagella of migrating human sperm, and its modulation by viscosity. *Cell Motility and the Cytoskeleton* **66** 220–236. (doi:10.1002/cm.v66:4)
- Strünker T, Goodwin N, Brenker C, Kashikar ND, Weyand I, Seifert R & Kaupp UB** 2011 The CatSper channel mediates progesterone-induced Ca^{2+} influx in human sperm. *Nature* **471** 382–386. (doi:10.1038/nature09769)
- Suarez SS** 2008 Control of hyperactivation in sperm. *Human Reproduction Update* **14** 647–657. (doi:10.1093/humupd/dmn029)
- Suarez SS & Dai X** 1992 Hyperactivation enhances mouse sperm capacity for penetrating viscoelastic media. *Biology of Reproduction* **46** 686–691. (doi:10.1007/s00441-015-2244-2)
- Suarez SS & Pacey AA** 2006 Sperm transport in the female reproductive tract. *Human Reproduction Update* **12** 23–37. (doi:10.1007/s00441-015-2244-2)
- Suarez SS, Katz DF, Owen DH, Andrew JB & Powell RL** 1991 Evidence for the function of hyperactivated motility in sperm. *Biology of Reproduction* **44** 375–381. (doi:10.1095/biolreprod44.2.375)
- Teves ME, Barbano F, Guidobaldi HA, Sanchez R, Miska W & Giojalas LC** 2006 Progesterone at the picomolar range is a chemoattractant for mammalian spermatozoa. *Fertility and Sterility* **86** 745–749. (doi:10.1016/j.fertnstert.2006.02.080)
- Uhler ML, Leung A, Chan SY & Wang C** 1992 Direct effects of progesterone and antiprogesterone on human sperm hyperactivated motility and acrosome reaction. *Fertility and Sterility* **58** 1191–1198.
- Uñates DR, Guidobaldi HA, Gatica LV, Cubilla MA, Teves ME, Moreno A & Giojalas LC** 2014 Versatile action of picomolar gradients of progesterone on different sperm subpopulations. *PloS ONE* **9** e91181. (doi:10.1371/journal.pone.0091181)
- Xia J, Reigada D, Mitchell CH & Ren D** 2007 CATSPER channel-mediated Ca^{2+} entry into mouse sperm triggers a tail-to-head propagation. *Biology of Reproduction* **77** 551–559. (doi:10.1095/biolreprod.107.061358)

Received 10 December 2015

First decision 8 January 2016

Revised manuscript received 31 January 2016

Accepted 22 February 2016