Human granulosa–luteal cells initiate an innate immune response to pathogen-associated molecules

Laila A Ibrahim^{1,3}, Joseph M Kramer^{1,2}, R Stan Williams^{1,2} and John J Bromfield^{1,3}

¹D. H. Barron Reproductive and Perinatal Biology Research Program, Gainesville, Florida, USA, ²Department of Obstetrics and Gynecology, College of Medicine, University of Florida, Gainesville, Florida, USA, and

³Department of Animal Sciences, University of Florida, Gainesville, Florida, USA

Correspondence should be addressed to J J Bromfield; Email: jbromfield@ufl.edu

Abstract

The microenvironment of the ovarian follicle is key to the developmental success of the oocyte. Minor changes within the follicular microenvironment can significantly disrupt oocyte development, compromising the formation of competent embryos and reducing fertility. Previously described as a sterile environment, the ovarian follicle of women has been shown to contain colonizing bacterial strains, whereas in domestic species, pathogen-associated molecules are concentrated in the follicular fluid of animals with uterine infection. The aim of this study is to determine whether human granulosa–luteal cells mount an innate immune response to pathogen-associated molecules, potentially disrupting the microenvironment of the ovarian follicle. Human granulosa–luteal cells were collected from patients undergoing assisted reproduction. Cells were cultured in the presence of pathogen-associated molecules (LPS, FSL-1 and Pam3CSK4) for 24 h. Supernatants and total RNA were collected for assessment by PCR and ELISA. Granulosa–luteal cells were shown to express the molecular machinery required to respond to a range of pathogen-associated molecules. Expression of *TLR4* varied up to 15-fold between individual patients. Granulosa–luteal cells increased the expression of the inflammatory mediators *IL1B*, *IL6* and *CXCL8* in the presence of the TLR4 agonist *E. coli* LPS. Similarly, the TLR2/6 ligand, FSL-1, increased the expression of *IL6* and *CXCL8*. Although no detectable changes in *CYP19A1* or *STAR* expression were observed in granulosa–luteal cells following challenge, a significant reduction in progesterone secretion was measured after treatment with FSL-1. These findings demonstrate the ability of human granulosa–luteal cells to respond to pathogen-associated molecules and generate an innate immune response.

Reproduction (2016) **152** 261–270

Introduction

Sexually transmitted infections in women of reproductive age are very common in the USA, with most infections occurring in young women aged between 15 and 24 years. The Gram-negative bacteria *Chlamydia trachomatis* and *Neisseria gonorrhoeae* are the leading causes of sexually transmitted bacterial infections with a combined total of 3.6 million new cases/year in the USA (277 million worldwide; Centers for Disease Control and Prevention). Pelvic inflammatory disease (PID) is caused primarily by untreated chlamydia or gonorrhea of the upper reproductive tract, with approximately 10–15% of women with chlamydia developing PID (reviewed in (Mitchell & Prabhu 2013)). Direct medical expenditures for PID and its sequelae were estimated at \$1.88 billion in 1998 (Rein *et al.* 2000). Although PID is a leading cause of gynecological hospitalization of reproductive aged women in the USA, 10% of PID patients become infertile primarily due to pelvic adhesions and tubal obstruction (Veale *et al.* 1995). Although PID is a cause of pelvic pain and infertility, little is known about the effects on the ovary and female gametes. However, it has been shown that patients with PID have reduced

follicle numbers and increased cystic follicles and corpora lutea after the resolution of the initial infection (Weiner & Wallach 1974, Bychkov 1990). Retrospective analysis of patients undergoing fertility treatment suggest that increased endometrial LPS reduces IVF success (Kamiyama *et al.* 2004). More recently, it has been demonstrated that bacteria can be isolated from the follicular fluid of women undergoing infertility treatment, with the extent of bacterial colonization of the follicle being correlated with the success of fertility treatment (Neuer *et al.* 1997, Pelzer *et al.* 2013).

The initial immune response to pathogens is driven by the innate immune system. Innate immune responses to bacteria depend on conserved pathogen-associated molecules being recognized by receptors of the innate immune system, such as Toll-like receptors (TLRs). Toll-like receptors are typically found in hematopoietic immune cells in addition to non-immune cells of the gut, which readily encounter bacteria. Previous work has demonstrated the ability of primary human granulosa–luteal cells (GLCs) to bind LPS (Sancho *et al.* 1992), whereas the granulosa cell line COV434

Table 1 Primer sequences for PCR.

Gene	Sequence (5'–3')		Accession number
	Sense	Antisense	
<i>ACTB</i>	ACAGAGCCTCGCCTTTGCCG	TTGCACATGCCGGAGCCGTT	NM_001101.3
<i>CD14</i>	CAGAACCTAGATGCCCTGC	CGCGCTCCATGGTCGATA	NM_000591.3
<i>CYP19A1</i>	GACGTCGCGACTCTAAATTGC	TGGGAGATGAGGGGTCCAAT	NM_000103.3
<i>IL1B</i>	AACCTCTTCGAGGCACAAGG	GTCCTGGAAGGAGCACTTCAT	NM_000576.2
<i>IL6</i>	CAGTTCCTGCAGAAAAAGGCAA	GCTGCGCAGATGAGATGAG	NM_000600.3
<i>CXCL8</i>	CAGAGACAGCAGAGCACACA	GGCAAACTGCACCTTCACA	NM_000584.3
<i>LY96 (MD2)</i>	ACACCTACTGTGGGAGAGAT	CGTCATCAGATCCTCGGCAA	NM_001195797.1
<i>MYD88</i>	TCTGTCTGCTGCTTCAAGAT	GACTGCTCGAGCTGCTTACC	NM_001172567.1
<i>PTPRC (CD45)</i>	TGAAAGAGTGAGAGTGGACGA	TTCTGGTGTCTGCCTGCTTC	NM_001267798.1
<i>STAR</i>	TGTCATCAGGGCGGAGCAC	CAGGACCTGGTTGATGATGCT	NM_000349.2
<i>TLR4</i>	CTGCCACATGTCAGGCCTTAT	AATGCCACCTGGAAGACTCT	NM_138554.4

possess TLRs and initiate an innate immune response to pathogen-associated molecules (Price *et al.* 2012). Whether human GLCs can initiate an innate immune response to pathogen-associated molecules and potentially alter the microenvironment of the follicle remains unknown.

We hypothesized that human GLCs isolated from infertility patients would initiate an innate immune response to pathogen-associated molecules similar to those found on sexually transmitted pathogens. Innate immune-mediated changes to the microenvironment of the ovarian follicle may help to further explain infertility associated with PID or sexually transmitted infections.

Materials and methods

Tissue collection and granulosa–luteal cell isolation

Human follicular aspirates were collected from women attempting conception through IVF, as well as from oocyte donors, at the University of Florida Reproductive Medicine

Clinic. The collection of follicular aspirates for this study was reviewed and approved by the Institutional Review Board at the University of Florida (#IRB201400399). All patients provided non-written consent to provide tissues as the University of Florida Institutional Review Board considered the follicular aspirates to be non-human waste collected during normal procedures and subsequently discarded and no patient information was obtained. Consenting patients underwent ovarian stimulation using standard long Lupron suppression or GnRH antagonist protocols. Oocyte retrieval was scheduled for 36 h after administration of an ovulation trigger. After oocytes were collected from the aspirate, the remaining follicular cells were collected by centrifugation at 800 *g* for 10 min. Cell pellets were then resuspended in warm PBS and washed by centrifugation at 800 *g* for 10 min. Cell pellets were resuspended in complete culture medium containing DMEM: Hams F12 (1:1), 10% fetal bovine serum, ITS, and 1% penicillin/streptomycin (all Thermo Fisher Scientific). A 1:1 (vol:vol) Percoll:PBS solution was prepared and the cell suspension layered on top of the Percoll solution and centrifuged at 800 *g* for 10 min. Cells were aspirated from a single layer in Percoll and washed in 10 mL warm PBS by

Figure 1 Granulosa–luteal cells express the TLR4 co-receptors and Toll-like receptors required to initiate an innate immune response. GLCs were collected from patients undergoing IVF. (A) After cellular purification, expression of the TLR4 co-receptors *MYD88*, *LY96*, and *CD14* were assessed by RT-PCR in addition to the granulosa cell-specific marker *CYP19A1* (aromatase). (B) Patient GLCs were assessed for the expression of all ten TLRs using RT-PCR after cellular purification. Commercial primer sets were utilized for the detection of individual TLR members. dsDNA was used as positive controls to demonstrate target amplification. RNA collected from whole blood was used as a positive control for all PCR. Each amplification was performed on a minimum of four patient samples.

Figure 2 Expression of granulosa–luteal cell *TLR4* varies between individual fertility patients. Following cellular purification, expression of *TLR4* was assessed in individual patients using real-time RT-PCR. A total of 18 patients were assessed for the expression of *TLR4* after confirming cultures were free of immune cell marker. Data were normalized to the housekeeper *ACTB*.

Figure 3 Granulosa-luteal cells increase expression of *IL1B*, *IL6*, and *CXCL8* in response to pathogen-associated molecules. Following a 24-h treatment of GLCs with tenfold increasing concentrations of *E. coli* LPS (A, B and C), FSL-1 (D, E and F), or Pam3CSK4 (G, H and I), expression of *IL1B* (A, D and G), *IL6* (B, E and H), and *CXCL8* (C, F and I) was assessed using real-time RT-PCR. Data are presented as mean $\pm$ S.E.M. from four to eight independent experiments. ** $P \leq 0.05$ compared with untreated controls; analysis by ANOVA followed by Dunnett's pairwise *post hoc* tests.

centrifugation at 800g. Cellular aggregates of GLCs were disassociated in 1 mL of 300 μ g/mL of hyaluronidase for 10 min at 37°C following a brief vortex. Cells were washed twice in complete medium by centrifugation at 800g for 10 min. Cells were resuspended and plated at 10^5 cells/mL in 500 μ L aliquots in 24-well plates (TPP; Switzerland) and incubated at 37°C with 5% CO₂ in a humidified environment.

Granulosa-luteal cell culture and challenge

Granulosa-luteal cells were allowed a 48-h period of equilibration before treatment. Each treatment was performed in a single well and repeated a minimum of four times on different days from different patients. Following 48 h of culture, medium was replaced with fresh medium containing either ultrapure *Escherichia coli* O111:B4 LPS (TLR4 agonist; 1 mg/mL is equivalent to 10^6 EU/mL), FSL-1 (TLR2/6 agonist; LPS ≤ 0.001 EU/ μ g), Pam3CSK4 (*N*-palmitoyl-S-[2,3-bis(palmitoyloxy)-(2RS)-propyl]-[R]-cysteiny]-[S]-seryl-[S]-lysyl-[S]-lysyl-[S]-lysyl-[S]-lysine) (synthetic ligand for TLR1/2; LPS ≤ 0.001 EU/ μ g), or ultrapure *Rhodobacter sphaeroides* LPS (LPS-RS; TLR4 antagonist) (all from Invivogen, San Diego, CA, USA). Treatments were performed at tenfold increasing doses between 100 pg/mL and 10 μ g/mL. After culture for 24 h, cell-free supernatants and total RNA were collected and stored at -20°C . A time-course experiment involved the treatment of cells with 100 ng/mL *E. coli* LPS between 15 and 360 min.

THP-1 cell line culture

The human monocyte cell line THP-1 (ATCC, Manassas, VA, USA) was used as a positive control throughout. THP-1 cells were authenticated by ATCC using short tandem repeat (STR) profiling and confirmed a 100% identity with THP-1 cells

according to ATCCs STR database. Cells were cultured in RPMI-1640 containing 10% FCS and 0.05 mM 2-mercaptoethanol. When required, cells were plated at a density of 10^5 cells/mL in 500 μ L aliquots in 24-well plates and incubated at 37°C with 5% CO₂ in a humidified environment. The addition of 50 ng/mL phorbol myristate acetate (PMA) facilitated the differentiation of monocytes to adherent macrophage-like cells that were used in experimental treatments (Park *et al.* 2007). Cells were allowed 48 h for differentiation and equilibration before treatment as per the methods described above.

RNA isolation and PCR

Cells were washed in warm PBS and lysed in RLT buffer to collect total RNA. Extraction of RNA was performed using the RNA Easy Mini kit according to the manufacturer's instruction (Qiagen). Total RNA was subjected to reverse transcription using the Verso cDNA synthesis kit according to the manufacturer's instruction (Thermo Fisher Scientific). Primers were designed using the NCBI database and initial specificity verified by BLAST to ensure no cross-reactivity with other loci (Table 1). Real-time PCR was performed in 25 μ L reactions using Maxima SYBR Green chemistry containing 0.5 μ M of each forward and reverse primer (Thermo Fisher Scientific). A CFX Connect light cycler was employed to perform quantitative PCR (Bio-Rad). Data were normalized independently to *ACTB* mRNA expression and plotted as expression relative to the mean of the untreated control group using the $\Delta\Delta\text{Ct}$ method. To examine cell-specific markers (*PTPRC* and *CYP19A1*) and the presence of receptors required for innate immune recognition of pathogen-associated molecules (*MYD88*, *LY96* and *CD14*), PCR products were electrophoresed on a 2% agarose gel containing ethidium bromide and visualized under

Figure 4 Acute response of granulosa–luteal cells to lipopolysaccharide. Acute responses of GLCs were measured following treatment with 100 ng/mL *E. coli* LPS. Cells were treated for 15–360 min before expression of *IL1B* (A), *IL6* (B), and *CXCL8* (C) was measured using real-time RT-PCR. Data are presented as mean $\pm$ S.E.M. from three to six independent experiments. ** $P \leq 0.05$ compared with untreated controls; analysis by ANOVA followed by Dunnett's pairwise *post hoc* tests.

UV illumination. Detection of the TLR family was performed using commercial primer sets from Invivogen according to the manufacturers' instructions and internal positive controls (Invivogen). Whole-blood RNA was purchased and used as an internal control for immune cell markers (Thermo Fisher Scientific).

ELISA

Progesterone and 17 β -estradiol were measured in cell-free supernatants using commercially available ELISA kits according to the manufacturer's instructions (DRG Instruments, GmbH, Germany). The limits of detection for 17 β -estradiol and progesterone were 10.7 and 0.3 pg/mL respectively; the intra-assay coefficients of variance were 1.1 and 6.4%, and the inter-assay coefficients of variance were 1.2 and 6.63% respectively.

Immunocytochemistry and fluorescence microscopy

Cells were plated onto sterile cover glass within 6-well culture plates for 24 h. On the day of labeling, cells were washed twice in warm PBS containing 0.5% BSA and subsequently blocked in 20% normal goat serum for 15 min in the incubator. Cells were again washed in warm PBS+BSA before application of a 1:10 dilution of anti-human CD45-FITC antibody (Milenyi Biotec, San Diego, CA, USA) in combination with 1 μ g/mL Hoechst (Thermo Fisher Scientific) for 30 min in the incubator. Cells were washed twice in PBS+BSA before immediate live cell image acquisition.

Cells were observed immediately using a Zeiss Axioplan 2 epifluorescence microscope and Axiovision software. Images were acquired using a Zeiss MRm digital camera. A minimum of 200 cells in five separate fields for each sample were assessed for Hoechst staining (total cells) and CD45-positive immunoreactivity (positive cells).

Statistical analysis

SPSS version 20.0 software was used for statistical analysis. Gene expression measured by real-time RT-PCR and concentrations of steroid hormones were analyzed using ANOVA, and comparisons were made with control using the Dunnett's pairwise *post hoc* test. If data were not normally distributed, it was log transformed for analysis. Data are arithmetic mean $\pm$ S.E.M., and a P value of ≤ 0.05 was assumed to be statistically significant.

Results

Granulosa–luteal cells are free of immune cells

Contamination of GLCs with immune cells was assessed to ensure that any response measured was due to target cells and not that of contaminating immune cells. Immunocytochemistry was performed to detect immunoreactivity to the immune cell marker PTPRC (CD45). The human monocyte cell line, THP-1, was 91.7% positive for PTPRC, whereas only 0.45% of GLCs displayed PTPRC immunoreactivity, suggesting a 99.5% immune cell marker-free preparation (Supplementary Fig. 1A and B, see section on supplementary data given at the end of this article). Following 40 cycles of RT-PCR, expression of *PTPRC* (CD45) was detected in blood while minimally visible in in GLC cultures

Figure 5 Pathogen-associated molecules do not change expression of *CYP19A1* or *STAR* in granulosa-luteal cells. Following a 24-h treatment of GLCs with tenfold increasing concentrations of *E. coli* LPS (A and B), FSL-1 (C and D) or Pam3CSK4 (E and F) expression of *CYP19A1* (A, C and E) and *STAR* (B, D and F) was assessed using real-time RT-PCR. Data are presented as mean $\pm$ S.E.M. from four to eight independent experiments. Compared with untreated controls; analysis by ANOVA followed by Dunnett's pairwise *post hoc* tests.

(Supplementary Fig. 1C). All samples were routinely screened for the presence of *PTPRC* at the initiation of culture to ensure an immune cell marker-free culture; any cultures shown to express *PTPRC* were excluded from analysis.

Granulosa-luteal cells express the receptors required to initiate an innate immune response

Initiation of innate immune signaling is dependent on the presence of TLR co-receptors MYD88, CD14, and LY96 (formerly MD2). Expression of *MYD88*, *LY96*, and *CD14* were all detected in GLC and blood (Fig. 1A). In addition, the expression of the granulosa cell-specific marker *CYP19A1* (aromatase) was shown to be present in GLCs and absent in blood (Fig. 1A). A PCR screen of all known human TLRs was performed to determine the capacity of GLCs to detect various pathogen-associated molecules. Expression of all ten TLRs was detected in blood, while GLCs had a distinct absence of *TLR1* and *TLR8* expression (Fig. 1B).

Quantitative PCR was used to assess the variable expression of the prototypic receptor, *TLR4*, on GLCs between 18 individual patients. Analysis showed that GLC expression of *TLR4* varied 15-fold between the highest and lowest expressing patients (Fig. 2).

Granulosa-luteal cells increase expression of inflammatory mediators in response to pathogen-associated molecules

Granulosa-luteal cells were challenged for a period of 24 h with one of the three pathogen-associated molecules: *E. coli* LPS (endotoxin, TLR4 agonist), FSL-1 (synthetic lipoprotein, TLR2/6 agonist), or Pam3CSK4 (synthetic triacylated lipopeptide, TLR1/2 agonist). Expression of the proinflammatory cytokines *IL1B* and *IL6* and the chemokine *CXCL8* was evaluated. Challenge of GLCs with LPS showed a dose-dependent increase in *IL1B*, *IL6*, and *CXCL8* expression after 24 h

(Fig. 3A, B and C; $P \leq 0.05$). Challenge with FSL-1 showed a significant increase in the expression of *IL6* and *CXCL8*, but only at higher concentrations (Fig. 3D, E and F; $P \leq 0.05$). Challenge of GLCs with Pam3CSK4 showed some numerical increase in the expression of *IL6* (Fig. 3G, H and I; not statistically significant). The human monocyte cell line, THP-1, was also challenged with pathogen-associated molecules under the same conditions as GLCs (Supplementary Fig. 2). THP-1 cells displayed very strong dose response to all three pathogen-associated molecules with increases to *IL1B*, *IL6* and *CXCL8* expression ($P \leq 0.05$).

Granulosa-luteal cells display an acute response to pathogen-associated molecules

Using LPS as the prototypical TLR4 agonist, a time-course experiment was performed to assess the acute responsiveness of GLCs. Following exposure to 100 ng/mL of LPS, GLCs displayed an increased expression of *IL1B* by 60 min, *IL6* by 90 min, and *CXCL8* by 120 min, which was maintained for 360 min (Fig. 4). Following treatment for 120 min, expression of *IL1B*, *IL6*, and *CXCL8* was higher than that observed in GLCs following 24 h (Fig. 3). The human monocyte cell line, THP-1, had a similar acute response to LPS compared with GLCs (Supplementary Fig. 3). THP-1 cells displayed increased expression of *IL1B* after 90 min and elevated *CXCL8* after only 60 min. Acute expression of *IL6* in THP-1 cells was muted. THP-1 cells had higher expression of both *IL1B* and *CXCL8* at 240 min compared with 24 h.

Effect of pathogen-associated molecules on granulosa-luteal cell expression of the steroidogenic enzymes *CYP19A1* and *STAR*

Granulosa cells of bovine have been shown to alter the expression of steroidogenic enzymes after challenge with pathogen-associated molecules (Williams *et al.* 2008, Price *et al.* 2013), whereas rats

Figure 6 Pathogen-associated molecules alter the accumulation of steroid hormone in granulosa–luteal cells. Following a 24-h treatment of GLCs with tenfold increasing concentrations of *E. coli* LPS, FSL-1, or Pam3CSK4 (PAM), accumulation of estradiol (A) or progesterone (B) was assessed in cell-free supernatants using ELISA. Data are presented as mean $\pm$ S.E.M. from six independent experiments. * $P \leq 0.08$ compared with untreated controls; analysis by nonparametric Mann–Whitney *U* test.

reduce estradiol secretion following exposure to LPS (Taylor & Terranova 1996). Interestingly, the human granulosa cell line (COV343) showed no change after challenge with pathogen-associated molecules (Price *et al.* 2012). We measured the expression of *CYP19A1* and *STAR* in GLCs following challenge with pathogen-associated molecules (Fig. 5). We did not detect any changes in the expression of either *CYP19A1* or *STAR* following any treatment.

Effect of pathogen-associated molecules on granulosa–luteal cell secretion of 17 β -estradiol and progesterone

Although no effect on *CYP19A1* or *STAR* expression was observed following exposure of GLCs to pathogen-associated molecules, we measured accumulation of 17 β -estradiol (Fig. 6A) and progesterone (Fig. 6B) in supernatants to confirm any effect on GLC function. Exposure of GLCs to higher concentrations of FSL-1

Figure 7 The TLR4 antagonist *Rhodobacter sphaeroides* LPS does not block the expression of *E. coli* LPS-induced inflammatory mediators in granulosa–luteal cells. GLCs were cultured in the presence of *E. coli* LPS alone (hatched), the TLR4 antagonist *Rhodobacter sphaeroides* LPS alone (RS-LPS; white), RS-LPS in combination with 100 ng/mL of *E. coli* LPS (black), or control culture medium (white). Following a 24-h treatment, the expression of the inflammatory mediators *IL1B* (A and E), *IL6* (B and F), and *CXCL8* (C and G) was assessed using real-time RT-PCR in THP-1 cells (A, B, C and D) or GLCs (E, F, G and H). Percent change of expression in cells exposed to RS-LPS in combination with *E. coli* LPS compared with LPS alone are presented for THP-1 (D) and GLCs (H). Data are presented as mean $\pm$ S.E.M. from three to eight independent experiments. ** $P \leq 0.05$ compared with untreated controls; analysis by ANOVA followed by Dunnett's pairwise *post hoc* tests.

reduced progesterone secretion when compared with untreated controls (2.2-fold reduction; $P \leq 0.08$). In addition, exposure of GLCs to either FSL-1 or Pam3CSK4 resulted in a numerical reduction in the secretion of both 17 β -estradiol and progesterone. Challenge of GLCs with LPS had no effect on either 17 β -estradiol or progesterone accumulation.

Figure 8 The TLR4 antagonist *Rhodobacter sphaeroides* LPS increases *CYP19A1* expression. GLCs were cultured in the presence of *E. coli* LPS alone (■), the TLR4 antagonist *Rhodobacter sphaeroides* LPS alone (RS-LPS; □), RS-LPS in combination with 100 ng/mL of *E. coli* LPS, or control culture medium). Following a 24-h treatment expression of steroidogenic factors, *STAR* (A) and *CYP19A1* (B) expressions were assessed using real-time RT-PCR. Percent change of RS-LPS alone-treated cells compared with untreated controls (C) and RS-LPS treated in combination with *E. coli* LPS compared with LPS alone (D) are presented. Data are presented as mean $\pm$ S.E.M. from three to eight independent experiments. ** $P \leq 0.05$; * $P \leq 0.08$ compared with untreated controls; analysis by nonparametric Mann-Whitney U test.

The TLR4 antagonist *Rhodobacter sphaeroides* LPS did not block expression of *E. coli* LPS-induced inflammatory mediators

Lipopolysaccharide derived from the bacterium *Rhodobacter sphaeroides* (RS-LPS) blocks pathogenic LPS/TLR4 activation by competitive binding of the TLR4 co-receptor LY96 (MD2) at 100-fold excess concentrations (Rallabhandi et al. 2012). Here, we attempted to block pathogenic LPS/TLR4 activation by adding RS-LPS in the presence of *E. coli*-derived LPS and assessing the inflammatory markers above. To confirm the validity of using RS-LPS to block TLR4 signaling, the monocyte cell THP-1 was challenged with *E. coli* LPS in combination with RS-LPS (Fig. 7A, B, C and D). Expression of LPS-induced *IL1B*, *IL6*, and *CXCL8* was reduced to vehicle-treated control when exposed to a 100-fold excess of RS-LPS in THP-1 cells. Expression of LPS-induced *IL1B*, *IL6*, and *CXCL8* was reduced to 58.2, 23.1, and 41.7% respectively of LPS alone treatment when THP-1 cells were exposed to 100-fold excess of RS-LPS (Fig. 7D). Challenge of GLCs with RS-LPS alone had no effect on the expression of the inflammatory mediators *IL1B*, *IL6*, or *CXCL8* (Fig. 7E, F, G and H). The addition of RS-LPS did not block the increased expression of *E. coli* LPS-induced inflammatory mediators, even at 100-fold excess concentrations in GLCs. Expression of LPS-induced *IL1B*, *IL6*, and *CXCL8* was 106.9, 79.9 and 40.0% respectively of LPS alone-treated GLCs when exposed to 100-fold excess of RS-LPS (Fig. 7H).

The TLR4 antagonist *Rhodobacter sphaeroides* LPS increases *CYP19A1* expression in granulosa-luteal cells

The expression of GLC *STAR* was not affected by either *E. coli* LPS or RS-LPS (Fig. 8A). As previously observed, *CYP19A1* expression was not affected in GLCs treated with *E. coli* LPS alone. However, challenge of GLCs with RS-LPS alone increased the expression of *CYP19A1* up to 14.8-fold over untreated controls (Fig. 8B). When GLCs were treated in combination with *E. coli* LPS and RS-LPS, expression of *CYP19A1* was comparable to untreated controls. Expression of RS-LPS-induced *CYP19A1* was between 549 and 1485% of untreated controls (Fig. 8C). Expression of *CYP19A1* or *STAR* in GLCs treated with *E. coli* LPS alone was comparable to GLC-treated cells in combination with RS-LPS (Fig. 8D).

Discussion

Infertility associated with the sequelae of bacterial infections of the upper reproductive tract is highly prevalent among women, estimated to represent approximately 20–30% of all infertility. Previous work has demonstrated that tubal pathologies associated with chlamydia or gonorrhea are the primary cause of infertility in these patients. However, more recent work has demonstrated that bacteria colonize the

ovarian follicle and that patients suffering PID show ovarian pathologies (Bychkov 1990, Pelzer *et al.* 2011, Pelzer *et al.* 2013). The data presented here suggest that GLCs can initiate an innate immune response to pathogen-associated molecules, potentially altering the microenvironment of the follicle. Although it is important to demonstrate the action of GLCs in initiating the immune response from a mechanistic stand point, it is also physiologically relevant as the ovarian follicle is normally replete of hematopoietic immune cells before ovulation when immune cells infiltrate the follicle to aid in the ovulatory process and control tissue damage (Brannstrom *et al.* 1994). Previous reports indicate that the cellular makeup of preovulatory human follicular fluid contains approximately 10% macrophages (Baranao *et al.* 1995), whereas in earlier follicle stages, immune cells are restricted to the ovarian theca (Brannstrom *et al.* 1993) and corpus luteum after ovulation (Petrovska *et al.* 1992). The potential of granulosa cells to alter secretion of inflammatory mediators in response to pathogen-associated molecules could alter the follicular environment or affect neighboring follicles. It is intriguing to consider the redundancies that exist between inflammatory pathways active during normal ovarian physiology and innate immune responses to pathogens. It may be that inappropriate activation of these pathways play a role in reducing ovarian function or oocyte quality following infection. A role of the chemokine IL-8 in promoting macrophages migration to preovulatory follicles has been suggested and is considered critical to facilitate the process of ovulation (Runesson *et al.* 1996). It could be presumed that inappropriate granulosa cell IL-8 secretion could recruit immune cells to the follicle before preparation for ovulation (Goto *et al.* 1997). Inflammatory mediators such as IL-1 β and IL-6 have been demonstrated to be important to oocyte developmental competence and ovulation, as such inappropriate concentration or temporal expression may have negative consequences on fertility (Machelon *et al.* 1994, Passos *et al.* 2016). The TLR4 pathway has also been demonstrated to be important in physiological ovulation in mice, again suggesting that inappropriate activation of the pathway could have negative consequences on ovarian function (Shimada *et al.* 2008).

Previous work in both bovine (Price *et al.* 2013) and rodent (Taylor & Terranova 1996) models indicates that pathogen-associated molecules have the ability to induce changes in granulosa cell steroid hormone production. Although we did not observe any changes in the expression of the steroidogenic enzymes *CYP19A1* or *STAR* in GLCs following challenge with pathogen-associated molecules, we observed a small reduction in progesterone secretion from GLCs treated with the TLR2/6 ligand FSL-1. This discrepancy from previously reported changes to steroid production is likely related to gonadotropin-induced steroid production, which was not examined in these current studies. However, LPS

does not alter FSH-induced steroid production in the human granulosa cell line COV343 (Price *et al.* 2012). Future experiments will determine the effect of pathogen-associated molecules on gonadotropin-dependent steroid hormone production in GLCs. Interestingly here, the TLR4 antagonist (RS-LPS) increased the expression of *CYP19A1* in GLCs. Currently, we have no clear explanation for this phenomenon; however, it has been shown that LPS from distinct pathogens can induce different cellular responses in other cell types (Pulendran *et al.* 2001). The mode of action for RS-LPS in this model system requires further investigation.

Although sexually transmitted infection-induced infertility is of great concern to the human population, animal models demonstrate the significant effect of upper reproductive tract infections on the ovary. Uterine infections are ubiquitous in dairy cows with approximately 40% of animals suffering significant bacterial uterine infections after parturition, many of which become infertile after the resolution of infection (reviewed in (Sheldon *et al.* 2009)). Cows that suffer these infections display disrupted ovarian function (Williams *et al.* 2008), whereas *in vitro* studies reveal that oocyte maturation is perturbed in the presence of LPS and granulosa cells initiate a significant inflammatory response to pathogen-associated molecules (Herath *et al.* 2007, Bromfield & Sheldon 2011). Similarly, in rodent models, exposure to LPS alters steroidogenesis of granulosa cells (Taylor & Terranova 1996) and significantly reduces the follicular reserve in a TLR4-dependent manner (Bromfield & Sheldon 2013). Currently, there are a limited number of studies that have investigated the effect of upper reproductive tract infections on the ovary in humans.

Studies investigating the effect of pathogen-associated molecules on fertility treatments have demonstrated a clear association between the presence of LPS in culture components, follicular fluid, or menstrual effluent to poor reproductive outcomes (Snyman & Van der Merwe 1986, Fishel *et al.* 1988, Nagata & Shirakawa 1996, Kamiyama *et al.* 2004). Although these associations have been made in clinical care, it is interesting to see in this study that patients displayed significant variability in the expression of *TLR4* in GLCs. Although this study does not have details regarding infection status of patients, which may result in variable expression of *TLR4*, it is routine for all patients undergoing fertility treatment to be free of any clinical upper reproductive tract infection. However, patient variability may be a result of subclinical infection or various single-nucleotide polymorphisms present in the human populations. Human polymorphisms of *TLR4* have been associated with the severity of inflammatory conditions such as asthma and arthritis (Zhang *et al.* 2011, Perica *et al.* 2015). Similarly, *TLR4* polymorphisms have also been associated with a decreased incidence of infections such as periodontitis (Sellers *et al.* 2015).

Although infection of the ovary itself is rare, it is proposed that systemic LPS exposure, or infection of tissues distant from the reproductive tract, can have impacts on the ovary. The presence of LPS-binding protein (LBP) in circulation is indicative of systemic LPS exposure by infection or possibly a “leaky gut wall”, which allows the translocation of LPS from the gut to circulation. In women undergoing IVF treatment, a positive correlation between the presence of LBP in circulation and follicular fluid IL-6 accumulation suggests that systemic exposure to LPS affects the follicular microenvironment. The same study revealed a negative correlation between systemic LBP and serum progesterone, similar to what was observed in this study (Tremellen *et al.* 2015). In dairy cows, bacterial infections of the mammary gland result in altered steroidogenesis, ovarian function, and oocyte quality following natural or induced disease (Lavon *et al.* 2008, 2011, Roth *et al.* 2013, Asaf *et al.* 2014).

To conclude, human GLCs from fertility patients express the molecular machinery necessary to initiate an innate immune response to pathogen-associated molecules, increasing expression of inflammatory mediators. These studies build on those previously performed in human’s investigating systemic endotoxemia and animal models of clinical disease. These data suggest that granulosa cells have the potential to respond to pathogens and consequently alter the microenvironment of the follicle, perturb oocyte quality, and potentially affect subsequently fertility.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/REP-15-0573>.

Declaration of interest

The authors have no conflict of interest to report that could be perceived as prejudicing the impartiality of the research reported.

Funding

This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

Ethical approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

References

- Asaf S, Leitner G, Furman O, Lavon Y, Kalo D, Wolfenson D & Roth Z 2014 Effects of *Escherichia coli*- and *Staphylococcus aureus*-induced mastitis in lactating cows on oocyte developmental competence. *Reproduction* **147** 33–43. (doi:10.1530/REP-13-0383)
- Baranao RI, Dain L, Palak de Fried E & Rumi LS 1995 Human granulosa cells express HLA-DR antigen and are capable of synthesizing interleukin-1. *Hormone and Metabolic Research* **27** 495–498. (doi:10.1055/s-2007-980010)
- Brannstrom M, Mayrhofer G & Robertson SA 1993 Localization of leukocyte subsets in the rat ovary during the periovulatory period. *Biology of Reproduction* **48** 277–286. (doi:10.1095/biolreprod48.2.277)
- Brannstrom M, Pascoe V, Norman RJ & McClure N 1994 Localization of leukocyte subsets in the follicle wall and in the corpus luteum throughout the human menstrual cycle. *Fertility and Sterility* **61** 488–495. (doi:10.1016/S0015-0282(16)56581-9)
- Bromfield JJ & Sheldon IM 2011 Lipopolysaccharide initiates inflammation in bovine granulosa cells via the TLR4 pathway and perturbs oocyte meiotic progression in vitro. *Endocrinology* **152** 5029–5040. (doi:10.1210/en.2011-1124)
- Bromfield JJ & Sheldon IM 2013 Lipopolysaccharide reduces the primordial follicle pool in the bovine ovarian cortex ex vivo and in the murine ovary in vivo. *Biology of Reproduction* **88** 98. (doi:10.1095/biolreprod.112.106914)
- Bychkov V 1990 Ovarian pathology in chronic pelvic inflammatory disease. *Gynecologic and Obstetric Investigation* **30** 31–33. (doi:10.1159/000293209)
- Fishel S, Jackson P, Webster J & Faratian B 1988 Endotoxins in culture medium for human in vitro fertilization. *Fertility and Sterility* **49** 108–111. (doi:10.1016/S0015-0282(16)59659-9)
- Goto J, Kanayama N, Asahina T, Okada Y, Kobayashi T & Terao T 1997 Induction of follicular growth by exogenous interleukin-8. *Human Reproduction* **12** 2729–2734. (doi:10.1093/humrep/12.12.2729)
- Herath S, Williams EJ, Lilly ST, Gilbert RO, Dobson H, Bryant CE & Sheldon IM 2007 Ovarian follicular cells have innate immune capabilities that modulate their endocrine function. *Reproduction* **134** 683–693. (doi:10.1530/REP-07-0229)
- Kamiyama S, Teruya Y, Nohara M & Kanazawa K 2004 Bacterial endotoxin in the endometrium and its clinical significance in reproduction. *Fertility and Sterility* **82** 805. (doi:10.1016/j.fertnstert.2004.06.030)
- Lavon Y, Leitner G, Goshen T, Braw-Tal R, Jacoby S & Wolfenson D 2008 Exposure to endotoxin during estrus alters the timing of ovulation and hormonal concentrations in cows. *Theriogenology* **70** 956–967. (doi:10.1016/j.theriogenology.2008.05.058)
- Lavon Y, Leitner G, Klipper E, Moallem U, Meidan R & Wolfenson D 2011 Subclinical, chronic intramammary infection lowers steroid concentrations and gene expression in bovine preovulatory follicles. *Domestic Animal Endocrinology* **40** 98–109. (doi:10.1016/j.domaniend.2010.09.004)
- Machelon V, Emilie D, Lefevre A, Nome F, Durand-Gasselin I & Testart J 1994 Interleukin-6 biosynthesis in human preovulatory follicles: some of its potential roles at ovulation. *Journal of Clinical Endocrinology and Metabolism* **79** 633–642. (doi:10.1210/jcem.79.2.7519193)
- Mitchell C & Prabhu M 2013 Pelvic inflammatory disease: current concepts in pathogenesis, diagnosis and treatment. *Infectious Disease Clinics of North America* **27** 793–809. (doi:10.1016/j.idc.2013.08.004)
- Nagata Y & Shirakawa K 1996 Setting standards for the levels of endotoxin in the embryo culture media of human in vitro fertilization and embryo transfer. *Fertility and Sterility* **65** 614–619. (doi:10.1016/S0015-0282(16)58164-3)
- Neuer A, Lam KN, Tiller FW, Kiesel L & Witkin SS 1997 Humoral immune response to membrane components of *Chlamydia trachomatis* and expression of human 60 kDa heat shock protein in follicular fluid of in-vitro fertilization patients. *Human Reproduction* **12** 925–929. (doi:10.1093/humrep/12.5.925)
- Park EK, Jung HS, Yang HI, Yoo MC, Kim C & Kim KS 2007 Optimized THP-1 differentiation is required for the detection of responses to weak stimuli. *Inflammation Research* **56** 45–50. (doi:10.1007/s00011-007-6115-5)
- Passos JR, Costa JJ, da Cunha EV, Silva AW, Ribeiro RP, de Souza GB, Barroso PA, Dau AM, Saraiva MV, Goncalves PB *et al.* 2016 Protein and

- messenger RNA expression of interleukin 1 system members in bovine ovarian follicles and effects of interleukin 1beta on primordial follicle activation and survival in vitro. *Domestic Animal Endocrinology* **54** 48–59. (doi:10.1016/j.domaniend.2015.09.002)
- Pelzer ES, Allan JA, Cunningham K, Mengersen K, Allan JM, Launchbury T, Beagley K & Knox CL 2011 Microbial colonization of follicular fluid: alterations in cytokine expression and adverse assisted reproduction technology outcomes. *Human Reproduction* **26** 1799–1812. (doi:10.1093/humrep/der108)
- Pelzer ES, Allan JA, Waterhouse MA, Ross T, Beagley KW & Knox CL 2013 Microorganisms within human follicular fluid: effects on IVF. *PLoS ONE* **8** e59062. (doi:10.1371/journal.pone.0059062)
- Perica M, Vidovic M, Lamot L, Bukovac LT, Kapitanovic S, Peric M, Barbic J & Harjacek M 2015 Single nucleotide polymorphism of toll-like receptor 4 (TLR4) is associated with juvenile spondyloarthritis in Croatian population. *Clinical Rheumatology* **34** 2079–2086. (doi:10.1007/s10067-015-2952-8)
- Petrovska M, Sedlak R, Nouza K, Presl J & Kinsky R 1992 Development and distribution of the white blood cells within various structures of the human menstrual corpus luteum examined using an image analysis system. *American Journal of Reproductive Immunology* **28** 77–80. (doi:10.1111/aji.1992.28.issue-2)
- Price JC, Cronin J & Sheldon IM 2012 Toll-like receptor expression and function in the COV434 granulosa cell line. *American Journal of Reproductive Immunology* **68** 205–217. (doi:10.1111/aji.2012.68.issue-3)
- Price JC, Bromfield JJ & Sheldon IM 2013 Pathogen-associated molecular patterns initiate inflammation and perturb the endocrine function of bovine granulosa cells from ovarian dominant follicles via TLR2 and TLR4 pathways. *Endocrinology* **154** 3377–3386. (doi:10.1210/en.2013-1102)
- Pulendran B, Kumar P, Cutler CW, Mohamadzahe M, Van Dyke T & Banchereau J 2001 Lipopolysaccharides from distinct pathogens induce different classes of immune responses in vivo. *Journal of Immunology* **167** 5067–5076. (doi:10.4049/jimmunol.167.9.5067)
- Rallabhandi P, Phillips RL, Boukhvalova MS, Pletneva LM, Shirey KA, Gioannini TL, Weiss JP, Chow JC, Hawkins LD, Vogel SN *et al.* 2012 Respiratory syncytial virus fusion protein-induced toll-like receptor 4 (TLR4) signaling is inhibited by the TLR4 antagonists Rhodobacter sphaeroides lipopolysaccharide and eritoran (E5564) and requires direct interaction with MD-2. *MBio* **3** e00218-12. (doi:10.1128/mbio.00218-12)
- Rein DB, Kassler WJ, Irwin KL & Rabiee L 2000 Direct medical cost of pelvic inflammatory disease and its sequelae: decreasing, but still substantial. *Obstetrics & Gynecology* **95** 397–402. (doi:10.1097/00006250-200003000-00016)
- Roth Z, Dvir A, Kalo D, Lavon Y, Krifucks O, Wolfenson D & Leitner G 2013 Naturally occurring mastitis disrupts developmental competence of bovine oocytes. *Journal of Dairy Science* **96** 6499–6505. (doi:10.3168/jds.2013-6903)
- Runesson E, Bostrom EK, Janson PO & Brannstrom M 1996 The human preovulatory follicle is a source of the chemotactic cytokine interleukin-8. *Molecular Human Reproduction* **2** 245–250. (doi:10.1093/molehr/2.4.245)
- Sanchotello M, Chen TY, Clinton TK, Lyles R, Moreno RF, Tilzer L, Imakawa K & Terranova PF 1992 Evidence for lipopolysaccharide binding in human granulosa-luteal cells. *Journal of Endocrinology* **135** 571–578. (doi:10.1677/joe.0.1350571)
- Sellers RM, Payne JB, Yu F, LeVan TD, Walker C & Mikuls TR 2015 TLR4 Asp299Gly polymorphism may be protective against chronic periodontitis. *Journal of Periodontal Research* **51** 203–211. (doi:10.1111/jre.12299)
- Sheldon IM, Cronin J, Goetze L, Donofrio G & Schuberth HJ 2009 Defining postpartum uterine disease and the mechanisms of infection and immunity in the female reproductive tract in cattle. *Biology of Reproduction* **81** 1025–1032. (doi:10.1095/biolreprod.109.077370)
- Shimada M, Yanai Y, Okazaki T, Noma N, Kawashima I, Mori T & Richards JS 2008 Hyaluronan fragments generated by sperm-secreted hyaluronidase stimulate cytokine/chemokine production via the TLR2 and TLR4 pathway in cumulus cells of ovulated COCs, which may enhance fertilization. *Development* **135** 2001–2011. (doi:10.1242/dev.020461)
- Snyman E & Van der Merwe JV 1986 Endotoxin-polluted medium in a human in vitro fertilization program. *Fertility and Sterility* **46** 273–276. (doi:10.1016/S0015-0282(16)49525-7)
- Taylor CC & Terranova PF 1996 Lipopolysaccharide inhibits in vitro luteinizing hormone-stimulated rat ovarian granulosa cell estradiol but not progesterone secretion. *Biology of Reproduction* **54** 1390–1396. (doi:10.1095/biolreprod54.6.1390)
- Tremellen K, Syedi N, Tan S & Pearce K 2015 Metabolic endotoxaemia – a potential novel link between ovarian inflammation and impaired progesterone production. *Gynecological Endocrinology* **31** 309–312. (doi:10.3109/09513590.2014.994602)
- Velebil P, Wingo PA, Xia Z, Wilcox LS & Peterson HB 1995 Rate of hospitalization for gynecologic disorders among reproductive-age women in the United States. *Obstetrics & Gynecology* **86** 764–769. (doi:10.1016/0029-7844(95)00252-M)
- Weiner S & Wallach EE 1974 Ovarian histology in pelvic inflammatory disease. *Obstetrics & Gynecology* **43** 431–437.
- Williams EJ, Sibley K, Miller AN, Lane EA, Fishwick J, Nash DM, Herath S, England GC, Dobson H & Sheldon IM 2008 The effect of Escherichia coli lipopolysaccharide and tumour necrosis factor alpha on ovarian function. *American Journal of Reproductive Immunology* **60** 462–473. (doi:10.1111/j.1600-0897.2008.00645.x)
- Zhang Q, Qian FH, Zhou LF, Wei GZ, Jin GF, Bai JL & Yin KS 2011 Polymorphisms in toll-like receptor 4 gene are associated with asthma severity but not susceptibility in a Chinese Han population. *Journal of Investigational Allergology & Clinical Immunology* **21** 370–377.

Received 2 December 2015

First decision 22 December 2015

Revised manuscript received 15 June 2016

Accepted 4 July 2016