The effects of radiofrequency electromagnetic radiation on sperm function

B J Houston¹, B Nixon¹, B V King², G N De Iuliis^{1,*} and R J Aitken^{1,*}

¹Priority Research Centre for Reproductive Science, School of Environmental and Life Sciences and

²School of Mathematical and Physical Sciences, University of Newcastle, Callaghan, New South Wales, Australia

Correspondence should be addressed to B J Houston; Email: brendan.houston@uon.edu.au

*(G N De Iuliis and R J Aitken contributed equally to this work).

Abstract

Mobile phone usage has become an integral part of our lives. However, the effects of the radiofrequency electromagnetic radiation (RF-EMR) emitted by these devices on biological systems and specifically the reproductive systems are currently under active debate. A fundamental hindrance to the current debate is that there is no clear mechanism of how such non-ionising radiation influences biological systems. Therefore, we explored the documented impacts of RF-EMR on the male reproductive system and considered any common observations that could provide insights on a potential mechanism. Among a total of 27 studies investigating the effects of RF-EMR on the male reproductive system, negative consequences of exposure were reported in 21. Within these 21 studies, 11 of the 15 that investigated sperm motility reported significant declines, 7 of 7 that measured the production of reactive oxygen species (ROS) documented elevated levels and 4 of 5 studies that probed for DNA damage highlighted increased damage due to RF-EMR exposure. Associated with this, RF-EMR treatment reduced the antioxidant levels in 6 of 6 studies that discussed this phenomenon, whereas consequences of RF-EMR were successfully ameliorated with the supplementation of antioxidants in all 3 studies that carried out these experiments. In light of this, we envisage a two-step mechanism whereby RF-EMR is able to induce mitochondrial dysfunction leading to elevated ROS production. A continued focus on research, which aims to shed light on the biological effects of RF-EMR will allow us to test and assess this proposed mechanism in a variety of cell types.

Reproduction (2016) **152** R263–R276

Introduction

Over the past 20 years, the use of mobile phones has increased exponentially (Gorpinchenko *et al.* 2014), with a current estimate of more than one billion users worldwide (French *et al.* 2001, Meral *et al.* 2007). In the United States, there is approximately one device in use per person, and well above more than one person in European countries such as Germany, Denmark and Italy (U.S. Census Bureau, 2012). Furthermore, the number of devices in service is rising at an estimated rate of 3% annually (ACMA 2013). Accordingly, the exposure of humans to radiofrequency electromagnetic radiation (RF-EMR) emitted from these devices has also increased substantially, with an average talk time of 30 min per day spent talking on mobile phones (CTIA 2011). The effect of this radiation on human health remains to be fully elucidated with current literature detailing an array of apparently contradictory results. Indeed, although some studies have identified pronounced deleterious effects of RF-EMR on a variety of cell types (Balode 1996, d'Ambrosio *et al.* 2002, Bilgici *et al.* 2013, Furtado-Filho

et al. 2014, Hou *et al.* 2015, Kahya *et al.* 2014, Dasdag *et al.* 2015), others have reported only very subtle or no significant effects (Marchionni *et al.* 2006, Masuda *et al.* 2006, Dasdag *et al.* 2009, Demirel *et al.* 2012, Khalil *et al.* 2014). A confounding factor in these studies involves the use of differing RF intensity, frequency, exposure length and method of administration, which discounts the possibility of direct and robust study-to-study comparisons. Such variation attempts to simulate elevated levels of exposure in certain studies and real-life mobile phone exposure in others, which is extremely hard to model given the variability that exists in each of these parameters of intensity and frequency (Lerchl 2013). For instance, the intensity of RF-EMR emitted from mobile phones varies from ~0.1–4 W/kg (Fejes *et al.* 2005, Guney *et al.* 2007, La Vignera *et al.* 2012), whereas mechanistic studies have involved intensities as high as 27.5 W/kg (De Iuliis *et al.* 2009a). Regardless of these differences, the balance of evidence supports the principle that RF-EMR has the ability to induce cellular damage (Adams *et al.* 2014). In light of this conclusion

and to work towards identifying real clinical risks, it is imperative that we develop an understanding of the mechanism(s) by which this form of radiation affects different biological systems.

Physical parameters of RF-EMR

Radiofrequency electromagnetic radiation is a form of microwave radiation. Its important properties include the frequency at which it is generated, measured in megahertz (MHz) or gigahertz (GHz), and the intensity of the waves, or the specific absorption rates (SAR), which is a measure of the rate of energy transfer from the electromagnetic field to particles in an absorber, defined at a particular point in the absorber (Durney 1986). The frequency of RF-EMR emitted by mobile phone devices is in the range of 900–1800 MHz, and the intensity of this radiation is generally restricted to a local limit of <2 W/kg and whole-body limit of 0.08 W/kg (Durney 1986, Chen 2007) to enforce safe exposure levels in humans. Meanwhile, the ability of RF-EMR itself to penetrate into the skin and body is dependent on the permittivity and conductivity of the irradiated tissue, as well as the wavelength of the radiation, which is inversely related to the wave frequency (Fig. 1). Therefore, at lower frequencies, the penetration of the RF-EMR is higher and devices operating in the 900 MHz range will irradiate the body more; approximately 25% of the body in humans compared with 20% penetration

at 1800 MHz (Durney 1986). However, it is possible that the penetration of RF-EMR into the testis may be more pronounced than other tissues, due to the fact that this organ is less protected by tissue in comparison to others. Mobile phone communications uses a variety of different frequency ranges, with the most common utilising the 880–915 MHz range for the global system for mobile communications (GSM) 900 uplink (from mobile phone to base station), 925–960 MHz for the GSM900 downlink (from base station to mobile phone), 1710–1785 MHz for the DCS1800 uplink, 1805–1880 MHz for the GSM1800 downlink, 1920–1980 MHz for the universal mobile telecommunications system (UTMS) data uplink and 2110–2170 MHz for the UTMS data downlink (Bolte & Eikelboom 2012). Of particular interest is this radiofrequency range, in which a majority of studies have used exposure frequencies of 900–1800 MHz. This in turn forms the basis of studies selected for this review.

Review focus

For the purpose of this review, we shall focus on an analysis of the effects of RF-EMR on the male reproductive system, a site that may be uniquely vulnerable to chronic EMR exposure from devices stored in the vicinity of the testes that are held in ‘standby mode’ and, more importantly, at the initiation of a call or when hands-free mode is in use. Our specific interest is to draw a consensus regarding the impact of RF-EMR on the male germline, with an emphasis on frequencies that equate to analog/digital signals (900/1800 MHz (Irmak et al. 2002)) and with specific absorption rates (SAR) of up to 4 W/kg. We imposed strict search criteria, which gives this review focus on probing a potential mechanism of action, independent of its clinical significance. To source the appropriate studies, we used the following search terms: ‘rf-emr spermatozoa’; ‘radiofrequency electromagnetic radiation spermatozoa’ and ‘cell phone radiation+spermatozoa’ in the PubMed database. Of those studies identified, we elected to review those reporting exposure at the RF range of between ~900 and 1800 MHz and those that focused on the male reproductive tract/spermatozoa. Such criteria were imposed to reflect the intensity of radiation emitted from the devices. This narrowed the list of articles to those summarised in Table 1. Largely independent of clinical significance, the unique cell biology of spermatozoa provides an ideal model in which the specific physical and chemical responses to EMR can be observed. These cells provide a sensitive model as (Aitken 2013, Aitken et al. 2014) (i) they are sensitive to damage by environmental factors including free radicals, (ii) they can be maintained for 48–72 h *in vitro* in simple, defined culture media, (iii) their motility provides a readily assessable means of monitoring adverse biological

Cell phone mode	Intensity (W/kg; 0, 10, 30 cm distance)
Standby	0.001
Talk (900 MHz)	0.011, 0.002, 0.003
Talk (1800 MHz)	0.008, 0.0009, 0.0002

Figure 1 Physical aspects of radiofrequency electromagnetic radiation. A table identifying the estimated intensity of radiation emitted from devices in talk mode of either 900 or 1800 MHz (Durney 1986, Panagopoulos et al. 2010, Liu et al. 2013a) and plot of penetration depth of this radiation in different tissue types over the MHz-GHz ranges (Gabriel et al. 1996).

Table 1 Review of studies investigating the effect of RF-EMR on the spermatozoa and male reproductive system of mice, rats and humans.

Reference	Species	Frequency (MHz)	Duration of exposure	Specific absorption rate (W/kg)	Motility	Vitality	ROS	DNA damage	Main outcomes
No effects Dasdag et al. (2003)	Sprague-Dawley rat	900	20 min per day, 4 weeks	0.52	NA	NA	NA	NA	No effects on testicular structure or sperm morphology
Imai et al. (2011)	Sprague-Dawley rat	1950	5 h per day, 5 weeks	0.4	NA	NA	NA	NA	No changes to epididymal or testis weights, increased sperm production with EMR treatment
Ozlem Nisbet et al. (2012)	Wistar rat	900/1800	2 h per day, 90 days	1.2–3/0.01–0.05 (900/1800)	–	NA	NA	NA	Increased sperm motility and morphology with EMR treatment
Sommer et al. (2009)	C57BL mouse	1966	24 h per day, 4 generations	0.08–2.34	NA	NA	NA	NA	No changes to sperm morphology, count, testis or epididymal weights
Trosic et al. (2013)	Wistar rat	915	1 h per day, 2 weeks	0.6	–	NA	NA	NA	No changes to motility, morphology or counts with EMR treatment
Tumkaya et al. (2013)	Sprague-Dawley rat	900	1 h per day, 45 days	0.48	NA	NA	NA	NA	No effects on testicular size, histology or spermatogenesis
Effects of RF-EMR Liu et al. (2013a)	Cultured mouse spermatoocyte	1800	1 min per 20 min, 24 h	0.13	NA	NA	NA	+	Increased DNA single-strand breaks with radiation intensity which was prevented with antioxidant pre-treatment
Agarwal et al. (2009)	Human spermatozoa	850	1 h	1.46	+	+	+	–	Healthy semen donors and infertility patients both experienced a loss in motility, vitality coupled with increases in ROS production. Infertility patients experienced a decreased total antioxidant status
De Iuliis et al. (2009a)	Human spermatozoa	1800	16 h	1	+	+	+	+	Dose-dependent effects for all parameters. At 1 W/kg significant decreases in motility and vitality, increases in ROS and DNA damage
Erogul et al. (2006)	Human spermatozoa	900	5 m	NA	+	NA	NA	NA	Reduced rapid and slow progressive sperm motility
Falzone et al. (2011)	Human spermatozoa	900	1 h	2	NA	NA	NA	NA	Morphological impacts: reduced acrosome and total sperm head sizes as well as zona binding
Fejes et al. (2005)	Human spermatozoa	NA	NA	NA	+	NA	NA	NA	Questionnaire for mobile phone usage, duration of mobile phone usage correlated negatively with progressive motility
Gorpinchenko et al. (2014)	Human spermatozoa	900/1800	5 h	NA	+	–	NA	+	Reduced progressive sperm motility, increased DNA fragmentation

(Continued)

Table 1 Continued.

Reference	Species	Frequency (MHz)	Duration of exposure	Specific absorption rate (W/kg)	Motility	Vitality	ROS	DNA damage	Main outcomes
Wdowiak et al. (2007)	Human spermatozoa	NA	0–2 years use of phone	NA	+	NA	NA	NA	Reduced sperm motility and increased irregular morphology
Zalata et al. (2015)	Human spermatozoa	850	60 min	NA	+	NA	NA	+	Significant reductions to sperm motility of men with asthenospermia and oligospermia, significant induction of DNA damage in sperm from healthy and sub-fertile semen profiles
Liu et al. (2015)	Sprague–Dawley rat	900	2 h per day, 50 days	0.66	NA	NA	+	NA	Decreased epididymis:body weight ratio, sperm count and total antioxidant capacity. Increased ROS concentration, apoptosis and ultrastructural neck deformations
Yan et al. (2007)	Sprague–Dawley rat	1900	6 h per day, 18 weeks	1.8	+	+	NA	NA	Significantly reduced sperm motility and vitality, abnormal sperm clumping
Aitken et al. (2005)	Swiss mouse	900	12 h per day, 7 days	0.09	–	–	NA	NA	No changes to motility, vitality, concentration or morphology with low SAR and duration. However, degradation to sperm mitochondrial genome
Al-Damegh (2012)	Wistar rat	900/1800	60 min per day, 14 days	0.9	NA	NA	+	NA	Antioxidant treatment prevented seminiferous tubule widening and reduced the lipid peroxidation onset by EMR treatment
Bin-Meferij and El-kott (2015)	Wistar rat	900	1 h per day, 8 weeks	NA	+	+	+	NA	Antioxidant treatment ameliorated a reduction in sperm motility, vitality, count, lipid peroxidation and morphological abnormalities observed with EMR exposure
Dasdag et al. (1999)	Wistar rat	900	3 min per day, 4 weeks	0.141	NA	NA	NA	NA	Thinning of seminiferous tubules, decreased progression of spermatogenesis. However, potential temperature influences
Ghanbari et al. (2013)	Wistar rat	915–950	8 h per day, 2–3 weeks	NA	+	+	+	NA	Time-dependent decreases to motility, vitality and antioxidant capacity
Kesari et al. (2011)	Wistar rat	900	2 h per day, 5 weeks	0.9	NA	NA	+	NA	Decreased glutathione peroxidase, superoxide dismutase, histone kinase expression; increased ROS, lipid peroxidation and apoptosis

Kesari and Behari (2012)	Wistar rat	900	2 h per day, 45 days	0.9	NA	NA	NA	Increased caspase activity, morphological abnormalities; decreased testosterone levels, progeny weight and number
Mailankot <i>et al.</i> (2009)	Wistar rat	900/1800	1 h per day, 4 weeks	NA	+	NA	NA	Reduced sperm motility, but not sperm count; increased MDA and decreased glutathione content of the testis and epididymis
Ozorak <i>et al.</i> (2013)	Wistar rat	900/1800	1 h per day, 4–6 weeks	0.18	NA	NA	NA	Significantly lower lipid peroxidation and total antioxidant status in the testis with 4-week EMR treatment. This change was a significant increase with EMR treatment after 6 weeks
Tas <i>et al.</i> (2014)	Wistar rat	900	3 h per day, 1 year	0.04	–	NA	NA	Increased morphological defects: tunica albuginea thinning, impaired spermatogenesis. No effects on sperm motility or concentration

NA, not mentioned or conducted in study; + negative effects documented; – no effects documented. Table arranged by model species used in study. EMR, electromagnetic radiation; MDA, malondialdehyde; ROS, reactive oxygen species; SAR, specific absorption rate.

effects and (iv) they are clinically important because DNA damage in spermatozoa has the potential to influence the health and wellbeing of the offspring. As a consequence of the information summarised in this review, we propose a mechanism for the negative effects of RF-EMR on the male germline. Given the unique susceptibility of spermatozoa to subtle oxidative insults, which may arise from RF-EMR exposure, the translation towards clinical significance, especially involving other cell types, should not be made. However, given that spermatozoa may be acutely sensitive to stressors such as RF-EMR, we propose that a clear hypothesis for a mechanism of action can be developed using this model, which can then be applied for testing in other cell types.

The impact of RF-EMR on semen quality

Mobile phone use is becoming increasingly popular worldwide, with specific population groups, including businessmen and adolescents, estimated to spend as much as half of their day in close proximity to mobile phones held in either active or standby modes (Redmayne *et al.* 2011, Roberts *et al.* 2014). Owing to the common practice of storing mobile phones in close proximity to the testes, these individuals may be unintentionally exposing their reproductive system to relatively high levels of RF-EMR. It is therefore of considerable concern that the use of mobile phones (Fejes *et al.* 2005, Yan *et al.* 2007, Agarwal *et al.* 2009, Gorpichenko *et al.* 2014, Zalata *et al.* 2015), or exposure to RF-EMR emitted by these devices (De luliis *et al.* 2009a, Al-Damegh 2012, Ghanbari *et al.* 2013), has been linked to negative impacts on semen quality. Notwithstanding considerable controversy regarding the timing and nature of such exposures (Dasdag *et al.* 2003, Imai *et al.* 2011, Tumkaya *et al.* 2013), the principle that RF-EMR can elicit a detrimental effect on sperm function is supported by a growing number of studies (Fejes *et al.* 2005, Agarwal *et al.* 2009, Mailankot *et al.* 2009, De luliis *et al.* 2009a, Liu *et al.* 2013a,b, Gorpichenko *et al.* 2014). In general, these data lend support to the notion that RF-EMR can significantly impair key aspects of sperm function including the motility and vitality of these cells and the integrity of their DNA (Table 1), suggesting a direct effect on mature spermatozoa. However, there is less compelling evidence to suggest an additional role at the level of spermatogenesis in reducing sperm counts *in vivo* (Imai *et al.* 2011, Tas *et al.* 2014). Indeed, a chronic, multigenerational study demonstrated RF-EMR to have no effects on sperm production and testicular or epididymal weight (Sommer *et al.* 2009).

Direct effects of RF-EMR on spermatozoa

In one of the earliest studies on the impact of RF-EMR on sperm quality, Wdowiak and coworkers (2007) demonstrated that males who use mobile phones exhibit

increased rates of abnormal sperm morphology and decreased motility compared with counterparts who did not use these devices. Furthermore, these effects were exacerbated with longer exposure to this form of radiation (Wdowiak *et al.* 2007). Since this report, additional studies have replicated the adverse impact of RF-EMR treatment on human sperm motility using a model waveguide device capable of emitting finely tuned electromagnetic radiation to mimic that emitted by mobile phones (Gajda *et al.* 2002, De luliis *et al.* 2009a). The waveguide approach improves control of exposure as well as replicates the use of a mobile phone held in talk mode (Agarwal *et al.* 2009).

Males experiencing subfertility, for example asthenozoospermia and oligozoospermia, appear to be particularly vulnerable to RF-EMR as highlighted by a marked decline in sperm motility after the exposure of semen samples to a mobile device for just 10 min (Zalata *et al.* 2015). Similar pronounced effects have also been documented after *in vivo* exposure of whole animals to a mobile phone operating in talk mode (Yan *et al.* 2007, Mailankot *et al.* 2009). In terms of the nature of the impaired motility, RF-EMR appears to primarily influence the capacity of spermatozoa to sustain forward progressive motility. Indeed, a study by Eroglu and coworkers (2006) confirmed that the exposure of human spermatozoa to RF-EMR compromised their ability to sustain both rapid and slow progressive motility after an alarmingly brief exposure time of only five minutes. Although other studies have required longer exposure times (hours or days) to generate significant reductions in sperm motility, impaired progressive motility (involving a decrease in the percentage of cells displaying rapid progressive motility and a corresponding increase in cells expressing slow progressive motility) appears to be a common consequence arising from RF-EMR exposure (Fejes *et al.* 2005, Gorpinchenko *et al.* 2014) and was observed in 11/15 studies, as presented in Table 1.

Nevertheless, these studies must be considered alongside others in which the presence of RF-EMR had no overt effect on either progressive (Tas *et al.* 2014) or overall sperm motility (Aitken *et al.* 2005, Imai *et al.* 2011, Trosic *et al.* 2013). A possible explanation for such inconsistencies in the effects of RF-EMR on sperm motility rests with the use of different exposure conditions. Indeed, in a majority of studies reporting negative impacts of RF-EMR on sperm motility (64%), the study design featured the use of isolated human spermatozoa that were exposed to RF-EMR via a mobile phone device. In contrast, at least half of the instances in which no effect was recorded on sperm motility, the studies involved whole-body animal exposure using a signal generator to produce the RF-EMR (Aitken *et al.* 2005, Trosic *et al.* 2013, Tas *et al.* 2014). Although these data further lend support to our proposal of spermatozoa as a sensitive model, they also highlight that *in vivo*, the body may be capable of absorbing

some of this radiation (Fig. 1), thus, diminishing the level of exposure experienced by spermatozoa within the reproductive system.

Effects of RF-EMR on spermatogenesis

In addition to the studies indicating that the RF-EMR can have detrimental effects on sperm function, there are sporadic reports that this type of radiation can also affect the testes. It has been demonstrated that a 60-minute exposure of male rats to RF-EMR daily for two weeks can cause widening of the seminiferous tubules (Al-Damegh 2012). In contrast, Dasdag and coworkers (1999) documented a thinning of seminiferous tubules in response to an intermittent mobile phone exposure of three minutes (on and off) for 2 h per day in active talk mode every day for one month. To add further difficulty to the interpretation of these data, a subsequent study by the same authors (Dasdag *et al.* 2003), reported no changes to testis structure after a similar RF-EMR exposure time of 20 min every day for one month. In addition to potential impacts on the diameter of the seminiferous tubules, chronic exposure (3 h per day for one year) of rats to RF-EMR reportedly elicited a reduction in the thickness of the tunica albuginea (Tas *et al.* 2014). Prolonged exposures (6 h daily over a 100-day period) have also been associated with patterns of sperm aggregation that were absent from unexposed rats and independent of any impact on sperm morphology (Yan *et al.* 2007). Nevertheless, abnormal sperm morphology arising from RF-EMR exposure has been documented (Wdowiak *et al.* 2007). In humans, these abnormalities have primarily been associated with the sperm head leading to a reduced capacity to engage in interactions with the oocyte (Falzone *et al.* 2011). Curiously however, Ozlem Nisbet and coworkers (2012) suggest that this form of insult appears to have no effect on the head morphology of rat spermatozoa at a frequency of 900 MHz, but instead alleviates the incidence of tail abnormalities and promotes a suite of positive functional outcomes, including increased testosterone levels and superior progressive motility. Furthermore, this group observed better formed seminiferous epithelia with 1800 MHz exposure that was not seen in 900 MHz or unexposed treatments. Moreover, another study involving exposure during pubertal development documented RF-EMR to induce no changes to the spermatogenic cycle or testicular morphology (Tumkaya *et al.* 2013).

Notwithstanding the conflicting nature of the data documented above, recent meta-analyses performed by Adams and coworkers (2014) and Liu and coworkers (2014) have concluded that RF-EMR has two major negative impacts on sperm function: significant reductions in motility and loss of viability. In line with the recent studies by Mailankot and coworkers (2009) and Trosic and coworkers (2013), this analysis confirmed that sperm concentration is not significantly

influenced by RF-EMR treatment. Although these data suggest that RF-EMR is not capable of causing major disruptions to the spermatogenic cycle, in line with Sommer and coworkers (2009), they do nonetheless highlight an effect on the functional attributes of spermatozoa. Such findings are particularly concerning given that they are attributed, at least in part, to studies involving human spermatozoa and therefore bring into question whether RF-EMR may be having any negative impact on fertility in our species. Collectively, the uncertainty surrounding the effects of RF-EMR on the male germline presents a challenge for interpretation, which is further exacerbated by the lack of any consolidated, mechanistic explanation for the effects of such low-energy radiation on biological systems.

Molecular mechanisms of RF-EMR action

Here, we focus on studies documenting the effects of RF-EMR on biology, with the purpose of identifying common pathways that may direct our understanding of how this factor influences biological systems. Furthermore, unveiling a mechanism to explain the biological stresses of RF-EMR will allow us to then rationally assess the clinical relevance of certain exposure conditions.

Generation of oxidative stress

It has previously been hypothesised that the biological effects of EMR could be attributed solely to heat stress, which is induced at the higher intensities of approximately $\geq 4\text{ W/kg}$ radiation used in some studies (Hossmann & Hermann 2003, Li *et al.* 2007). However, through the use of various 'intermittent' exposure systems (e.g. 5 min on/10 min off), it has been demonstrated that the effects of bulk heat stress are likely to be negligible at the intensities of radiation generated during typical RF-EMR exposure (Liu *et al.* 2013a). Such results have subsequently been verified in the transformed GC2 mouse spermatocyte cell line, in which it was shown that such transient exposure patterns are capable of inducing DNA fragmentation and oxidised base adduct formation (Liu *et al.* 2013b, Duan *et al.* 2015) in the absence of a significant impact on temperature.

RF-EMR treatment is known to have the capacity to induce oxidative stress, characterised by excessive generation of reactive oxygen species (ROS) that overwhelm the intrinsic cellular antioxidant capacity, in a variety of tissue types. Indeed, this phenomenon has been documented after RF-EMR treatment in whole-body and ovarian tissue models of *Drosophila* (Manta *et al.* 2014), mouse fibroblasts (Hou *et al.* 2015), cultured breast cancer cells (Kahya *et al.* 2014), rat heart tissue (Ozguner *et al.* 2005), human lens epithelial cells (Yao *et al.* 2008) and mammalian spermatozoa

(Agarwal *et al.* 2009, De Iuliis *et al.* 2009a, Kesari *et al.* 2011). We have also replicated this response using transformed male spermatogonial and spermatocyte germ cell lines, documenting an increase in ROS of mitochondrial origin (B Houston & R J Aitken 2015, unpublished observations). Furthermore, of the 27 RF-EMR exposure studies summarised in Table 1, at least 21 of these (78%) document negative effects of RF-EMR on one or more parameters of sperm function and/or testicular histology that are characteristic of responses elicited by oxidative stress, such as lipid peroxidation, impaired motility and the formation of oxidative DNA damage.

Such pronounced effects on the male germline may stem from the fact that spermatozoa are uniquely susceptible to oxidative stress. This vulnerability arises due to the highly specialised structure of the spermatozoon, featuring limited protective antioxidant capacity due to a diminutive cytoplasmic volume and, at the same time, an abundance of substrates for free radical attack including DNA, thiol-rich proteins and polyunsaturated fatty acids (PUFAs) (Aitken *et al.* 2012a). The latter are of critical importance to the spermatozoon and are required to generate the membrane fluidity needed to support both motility and the membrane-fusion events associated with fertilisation (Lenzi *et al.* 2000). Yet when peroxidised, PUFAs elicit the formation of small molecular mass, electrophilic aldehydes that perpetuate a state of oxidative stress (Aitken *et al.* 2012a) as detailed in Fig. 2.

Human spermatozoa exposed to RF-EMR exhibit significant increases in mitochondrial and cytosolic superoxide formation (Agarwal *et al.* 2009, De Iuliis *et al.* 2009a), as well as a significant reduction in sperm motility (Fejes *et al.* 2005, Gorpichenko *et al.* 2014). The causative link between excess ROS production and sperm motility loss is a well-established paradigm in sperm biology (Fig. 2). This is commonly attributed to increased lipid peroxidation and the ensuing formation of electrophilic aldehydes such as malondialdehyde, 4-hydroxynonenal (4HNE) and acrolein, which are capable of covalently binding to proteins, thus compromising their function (Jones *et al.* 1979, Koppers *et al.* 2008, 2010, Aitken *et al.* 2012a,b, Moazamian *et al.* 2015). In the case of sperm motility, these compounds appear to alkylate sperm axonemal proteins that regulate sperm motility, particularly dynein heavy chain (Baker *et al.* 2015, Moazamian *et al.* 2015). In addition, electrophiles such as 4HNE are also known to promote oxidative stress by stimulating ROS generation through the sperm mitochondria (Fig. 2). This situation arises because another group of proteins alkylated by 4HNE is the constituents of the mitochondrial electron transport chain (ETC), particularly succinic acid dehydrogenase (Aitken *et al.* 2012b). When these proteins become adducted by 4HNE, it promotes the leakage of electrons from the ETC, which are then

Figure 2 Oxidative stress cascade within the spermatozoon. ROS is formed within the cell from a variety of possible sources including mitochondrial dysfunction, plasma membrane NADPH oxidase activity, infiltrating leukocytes and environmental factors such as electromagnetic radiation. In the event these ROS outweigh the poor antioxidant capacity of the cell, or a deficiency in this protection exists, a state of oxidative stress ensues. ROS, particularly hydrogen peroxide, attack the lipid membranes which are richly bestowed with polyunsaturated fatty acids that are susceptible to oxidative attack, resulting in the formation of small, reactive aldehydes – acrolein, malondialdehyde and 4-hydroxynonenal. Although these aldehydes differ in their reactivity (Moazamian *et al.* 2015), they each target a specific subset of protein centres, typically thiol constituents, as a form of nucleophilic attack. One major consequence of this is impairment of protein function, such as key proteins involved in sperm motility. Succinate dehydrogenase, a protein complex within the mitochondria, is a predominantly vulnerable target of these electrophilic aldehydes, and alkylation of this complex results in the disruption of redox-regulated metabolism within the mitochondria, forcing electron flow to oxygen and thus forming yet more superoxide anion. Furthermore, this imbalance of ROS leads to oxidative DNA damage as hydrogen peroxide migrates to the sperm head and preferentially targets guanine residues within the sperm DNA, highlighted by significant increases in the oxidised base product 8-hydroxy-2'-deoxyguanosine.

consumed by the universal electron acceptor, oxygen, to generate superoxide anion (Aitken *et al.* 2012b). By such mechanisms, even slight increases in ROS induced by RF-EMR have the potential to become amplified through the mediation of the mitochondria. In support of this mechanism, it has been revealed that RF-EMR-induced ROS production does encourage lipid peroxidation in spermatozoa (Kesari *et al.* 2011, Al-Damegh 2012). Moreover, lipid peroxidation has also been localised within the testicular and epididymal microenvironments after RF-EMR treatment *in vivo*, and this has, in turn, been associated with a loss of sperm motility (Mailankot *et al.* 2009).

If RF-EMR is responsible for the induction of oxidative stress, we should see evidence of ROS overwhelming the sperm cell's antioxidant defences under these conditions (Gharagozloo & Aitken 2011). Indeed, intracellular concentrations of glutathione peroxidase and superoxide dismutase have been shown to be compromised in the spermatozoa of RF-EMR-exposed rats (Kesari *et al.* 2011). Furthermore, the addition of exogenous antioxidants such as vitamin C or E has been shown to significantly diminish RF-EMR-induced lipid peroxidation, while simultaneously leading to

a partial restoration of the glutathione content of the testis in RF-EMR-exposed rats (Al-Damegh 2012). As an extension of this work, both spermatozoa (Kesari *et al.* 2011) and testes (Al-Damegh 2012) respond by increasing catalase activity after exposure to EMR. This potentially represents a physiological response aimed at counteracting increases in hydrogen peroxide and other ROS formation induced by RF-EMR stress. Interestingly, it has been suggested that RF-EMR may have more pronounced effects in poor quality spermatozoa as revealed in studies where only a proportion of the sperm population was found to respond to RF-EMR treatment (De Iuliis *et al.* 2009a). If this was the case, then the increased ROS production generated in these highly vulnerable cells could reasonably be expected to impose an oxidative stress environment upon the remainder of the sperm population (Tosic & Walton 1950).

Downstream of lipid peroxidation, oxidative stress is known to culminate in oxidative damage to sperm DNA (Fig. 2). This has been characterised by elevated levels of the DNA damage marker, 8-hydroxy, 2'-deoxyguanosine (8OHdG; Aitken *et al.* 2012b,c, Aitken *et al.* 2014). Accordingly, RF-EMR exposure has been shown to elicit a significant increase in the staining intensity for this marker in human spermatozoa (De Iuliis *et al.* 2009a). RF-EMR has also been correlated with DNA strand breakage in spermatozoa (Zalata *et al.* 2015), cultured spermatogonia (B Houston & R J Aitken 2015, unpublished observations) and spermatocyte cells (Liu *et al.* 2013a). In the latter cell type, the DNA damage was successfully ameliorated by co-incubation of the cells with the antioxidant, melatonin (Liu *et al.* 2013a). Meanwhile, the observation that RF-EMR has the potential to generate sperm DNA damage is especially concerning due to the fact that these cells are capable of harbouring a considerable oxidative DNA damage load independent of any pronounced effects on motility (Aitken *et al.* 1998). These spermatozoa therefore have potential to participate in fertilisation, whereupon the oocyte would bear the responsibility for repairing the DNA before the initiation of S-phase of the first mitotic division. The fact that oocytes are relatively deficient in the first enzyme in the base excision repair pathway, OGG1 (Lord & Aitken 2015), means that any 8OHdG brought into the egg by the fertilising spermatozoon are likely to persist into the first cleavage division. As 8OHdG lesions are potentially mutagenic, these considerations may carry implications for the mutational load subsequently carried by the offspring, if the father's germline has been oxidatively damaged by RF-EMR.

The ability of RF-EMR to induce damage, which leads to negative biological outcomes is yet to reach consensus; nevertheless, biological effects of RF-EMR are more strongly demonstrated in the literature and are likely to depend on the properties of the affected macromolecule. With respect to proteins, it is expected that this form of damage could be resolved upon

turnover or degradation. However, in the case of long-lived molecules such as DNA, the impact of such damage could be far more insidious. This is particularly the case in the male germline where the integrity of the paternal genome has direct implications for future generations. Of particular concern is the potential for the damage to be acquired in post-meiotic germ cells, which have limited DNA repair mechanisms and are therefore unequipped to resolve the damage. This has been shown previously in spermatozoa, by the existence of dominant lethal mutations (Singer *et al.* 2006), which indicate the possibility of these mutations to be transferred through one generation. Given the strong paradigm for oxidative stress as a key mediator of sperm quality and that published data support the conclusion that RF-EMR can drive ROS production in the male germline, understanding how RF-EMR induces ROS is therefore of key importance.

Metabolic pathways activated by RF-EMR

It has been demonstrated that RF-EMR has the ability to stimulate signalling pathways in somatic cells, such as those associated with the extracellular signal-regulated kinase (ERK) cascade (Friedman *et al.* 2007) or heat-shock protein response (Di Carlo *et al.* 2002, Li *et al.* 2007, Valbonesi *et al.* 2014). As both of these pathways are known to be redox regulated, it is possible that RF-EMR activates these signal transduction cascades as a secondary consequence of ROS production (Christman *et al.* 1985, Polla *et al.* 1996, Nahomi *et al.* 2015). As indicated previously, the major site of intracellular ROS generation observed after RF-EMR exposure is the mitochondria.

There are several lines of evidence that point to the mitochondria being the major mediator of RF-EMR action of biological systems. Thus, in pancreatic cancer cells, it has been shown that EMR has the ability to induce extensive changes to the morphology of the mitochondria, stimulating a loss of their membrane potential and significantly increasing production of ROS (Curley *et al.* 2014). This effect is mirrored across a variety of additional somatic cell types including rat hippocampal slices where EMR evokes substantial changes to mitochondrial morphology (Zhao *et al.* 2012) and membrane potentials (Tattersall *et al.* 2001), and human peripheral blood monocytes where it induces a transient decrease in mitochondrial membrane potential that is accompanied by increased ROS production and caspase activation; the latter of which are hallmarks of an apoptotic cascade (Lu *et al.* 2012). As indicated previously, there is also very clear evidence that RF-EMR activates mitochondrial ROS generation in spermatozoa (De Iuliis *et al.* 2009a).

Although such effects of RF-EMR have been recorded at radiofrequency levels of around 900–1800 MHz, corresponding to that emitted by mobile phones

(Marchionni *et al.* 2006), contradictory stimulatory effects have in fact been observed at very low frequencies, less than 100 MHz (Marchionni *et al.* 2006, Iorio *et al.* 2011). Indeed, in marked contrast to the negative effects of RF-EMR, extremely low-frequency EMR (50 Hz) has in fact been shown to encourage sperm motility (Iorio *et al.* 2011). This effect is also believed to be a consequence of altered mitochondrial activity; however, in this instance, it appears that the EMR exposure leads to an increase in mitochondrial membrane potential (Iorio *et al.* 2011). Such a discrepancy may be explained, at least in part, by the variable degree of penetration achieved with EMR of different wavelengths (Lin 1976; Fig. 1). In this context, it is well established that the intensity of the RF-EMR decays exponentially as it penetrates the skin, whereas penetration depth varies between different tissues and organs (Fig. 1; De Iuliis *et al.* 2012, Markov & Grigoriev 2015). This radiation exposure generally depends on emitted power, but to some extent, it also depends on other parameters such as the frequency, antenna position relative to the body and the material properties of the absorbing tissue (Balzano 1999). In any case, the biophysics involved in these types of interactions is unresolved and represents a major limitation regarding RF-EMR studies (Lerchl 2013). We have also observed subtle variations in the response to RF-EMR when assessing mitochondrial function in male germ cells at different stages of maturation, with vulnerabilities to RF-EMR appearing to be dependent on the stage of development (B Houston & R J Aitken 2015, unpublished observations). This again highlights the potential difficulties with interpreting and rationalising the effects of RF-EMR on biology, given the diversity of cells that are potentially exposed by mobile phone use.

It is also probable that the variation in mitochondrial membrane potential stimulated by EMR is dependent on SAR, as extremely low-intensity radiation (2.5×10^{-5} W/kg) fails to alter mitochondrial membrane potential in human promyelotic leukaemia cells (Jin *et al.* 2012). Similarly, mitochondrial membrane potential also remains unaffected when exposed to low doses of EMR ($150\text{--}570 \mu\text{W}/\text{cm}^2$) in mouse endometrial glandular cells, but it is successfully impaired with higher intensities ($1400 \mu\text{W}/\text{cm}^2$) (Liu *et al.* 2012). In human spermatozoa, mitochondrial ROS generation was evident at SAR values above 2.8 W/kg (De Iuliis *et al.* 2009a), although there are no data linking such ROS generation to a change in mitochondrial membrane potential. Nevertheless, an increase in ROS generation has been consistently reported in studies focusing on the impacts of RF-EMR on spermatozoa (Agarwal *et al.* 2009, De Iuliis *et al.* 2009a, Kesari *et al.* 2011, Al-Damegh 2012).

It should be noted that within the electron transport chain, small concentrations of superoxide are a normal by-product of this essential redox process. However, the magnitude of ROS leakage varies between the ETC complexes, with Complex I (NADH oxidase) responsible

for a bulk of the superoxide, and with the substrate used for energy production, as observed in isolated mitochondria (Quinlan *et al.* 2013). It is also important to note that superoxide production at Complex I is much more damaging than at Complex III in spermatozoa, due to the mode of emigration of ROS from Complex I to the matrix, allowing for subsequent peroxidative damage (Koppers *et al.* 2008). Meanwhile, ROS generated at Complex III escapes to the intermembrane space, where it encounters the pool of mitochondrial antioxidant protection. The movement of electrons through the electron transport chain is a highly regulated process, partly to limit the production of deleterious amounts of ROS. Perturbation of the electron flow through this chain by RF-EMR, and the subsequent promotion of electron leakage within the mitochondria, would provide a gateway for the formation of ROS such as the superoxide anion (Martino & Castello 2011) as part of a two-step process (Fig. 3). Considering that RF-EMR specifically promotes mitochondrial ROS production (De Iuliis *et al.* 2009a, Burlaka *et al.* 2013) associated with increased expression of mitochondrial apoptotic markers (Liu *et al.* 2015) and decreased mitochondrial membrane potential (Lu *et al.* 2012), we propose that this radiation potentiates the leakage of electrons within the electron transport chain. Such electron leakage may be achieved through interference with proton transmission through the transmembrane complexes of the inner mitochondrial membrane. This is caused by the ability of modulated EMR (such as that emitted from mobile

phones) to augment the oscillation of ions, interfering with their transport through membrane proteins, thus potentially perturbing the strict membrane potentials (Panagopoulos *et al.* 2000, 2002, 2015) enforced in the specific intermembrane compartments of the mitochondria, which otherwise stabilise proton flow (Fig. 3; Perry *et al.* 2011). A consequence of reduced proton emigration is a reduced proton motive force and a subsequent reduction in ATP production (Perry *et al.* 2011). Under these conditions, when the NADH/NAD⁺ ratio is high and associated with low or compromised mitochondrial respiration, as previously shown to be induced by EMR (Sanders & Joines 1984), superoxide is formed at Complex I (Kudin *et al.* 2004, Murphy 2009). This scenario is accompanied by the ability of RF-EMR treatment to significantly impair the conformation of proteins and DNA, including key antioxidant proteins (Lu *et al.* 2012), preventing them from participating in the elimination of radicals generated during respiration. Thus, as a first step, the combined effects of RF-EMR results in an imbalance of free radical formation and antioxidant status, driving a state of oxidative stress (Fig. 3). The ROS formed through this process, modified to hydrogen peroxide via mitochondrial superoxide dismutase, would in turn have the ability to drive a lipid peroxidation cascade (Al-Damegh, 2012), resulting in the production of electrophilic aldehydes including malondialdehyde (Mailankot *et al.* 2009, Kesari *et al.* 2011) and 4HNE (Moazamian *et al.* 2015). Once formed, these potent electrophiles activate the second

Figure 3 Potential effects of RF-EMR on the mitochondrial electron transport chain. Electron flow within the transport chain usually involves transfer of electrons through Complexes I and II into the Q pool where the electrons then feed into Complex III, interact with cytochrome-C and finally Complex IV where water acts as the terminal electron acceptor. Step 1, the presence of EMR may interfere with proton flow through these complexes, reducing proton motive force and ATP production. Via such mechanisms, EMR would also increase the NADH/NAD⁺ ratio (Sanders & Joines 1984), which would, in turn, promote the leakage of electrons from NADH to oxygen, forming superoxide anion – a progenitor ROS molecule. Subsequent dismutation of superoxide to H₂O₂ allows for step 2, where an imbalance of ROS results in lipid peroxidation and the formation of electrophilic aldehydes. These nucleophilic compounds impair the electron transport chain further by binding to the complexes of the ETC, promoting additional dislocation of electron flow and generating yet more superoxide, promoting extensive lipid peroxidation, motility loss and oxidative DNA damage. Grey arrows represent proton movement, black arrows represent electron flow, dashed lines represent electron leakage and thunderbolts denote EMR. C, cytochrome-C; F, FADH₂; N, NADH; Q pool, quinone pool; GPx, glutathione peroxidase.

step of this response, inducing widespread interference within the electron transport chain by directly alkylating key proteins associated with the protein complexes of this pathway. As mentioned previously, Complex II (succinate dehydrogenase) of this chain is preferentially targeted by 4HNE (Aitken *et al.* 2012b). Modification or inhibition of Complex II prevents the oxidation of FAD in the succinate dehydrogenase-A subunit, forcing the flow of electrons to oxygen and thus resulting in elevated mitochondrial perturbation with consequential increases in superoxide formation (Zhang *et al.* 1998, Aitken *et al.* 2012b). Moreover, as mitochondria are responsible for a majority of ROS production within spermatozoa (Koppers *et al.* 2008), it is conceivable that disrupting the function of these organelles accounts for the elevated ROS production observed with RF-EMR treatment in several studies, as exemplified by De Iuliis and coworkers (2009b). An important feature of this putative mechanism is that it would account for the subtle or variable changes that RF-EMR has been recorded to induce in terms of sperm motility, owing to the fact that in species such as humans, mice and rats, the energy demands required to support motility are not exclusively dependent on oxidative phosphorylation (Williams & Ford 2001, Storey 2008). However, it should be taken into account that these cells are susceptible to a state of oxidative stress.

Conclusion

To date, contradictory studies surrounding the impacts of RF-EMR on biological systems maintain controversy over this subject. Nevertheless, research on the biological responses stimulated by RF-EMR is particularly important given our ever-increasing use of mobile phone technology. Although clinical studies are identifying possible detrimental effects of RF-EMR, it is imperative that mechanistic studies are conducted that elucidate the manner in which RF-EMR perturbs biological function, thus supplying a rational cause. A focus on the male reproductive system is justified given the potentially elevated levels of exposure this system may experience as consequences of the personal storage of mobile devices, the unique vulnerability of the highly specialised sperm cell, and the future health burden that may be created if conception proceeds with defective, DNA-damaged spermatozoa. Although this subject remains a topic of active debate, this review has considered the growing body of evidence suggesting a possible role for RF-EMR-induced damage of the male germline. In a majority of studies, this damage has been characterised by loss of sperm motility and viability as well as the induction of ROS generation and DNA damage. We have therefore given consideration to the potential mechanisms through which RF-EMR may elicit these effects on spermatozoa, which we

used as a sensitive model system. We propose a mechanistic model in which RF-EMR exposure leads to defective mitochondrial function associated with elevated levels of ROS production and culminates in a state of oxidative stress that would account the varying phenotypes observed in response to RF-EMR exposure. With further complementary data, this model will provide new impetus to the field and stimulate research that will allow us to confidently assess the reproductive hazards of mobile phone usage.

Declaration of interest

The authors declare no conflicts of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by the Australian Research Council Discovery Project scheme (grant number DP110103951) to R J A, and B V K. B H is the recipient of an Australian Postgraduate Award PhD scholarship.

References

- Adams JA, Galloway TS, Mondal D, Esteves SC & Mathews F 2014 Effect of mobile telephones on sperm quality: a systematic review and meta-analysis. *Environment International* **70** 106–112. (doi:10.1016/j.envint.2014.04.015)
- Agarwal A, Desai NR, Makker K, Varghese A, Mouradi R, Sabanegh E & Sharma R 2009 Effects of radiofrequency electromagnetic waves (RF-EMW) from cellular phones on human ejaculated semen: an in vitro pilot study. *Fertility and Sterility* **92** 1318–1325. (doi:10.1016/j.fertnstert.2008.08.022)
- Aitken RJ 2013 Human spermatozoa: revelations on the road to conception. *F1000 Prime Reports* **5** 39. (doi:10.12703/p5-39)
- Aitken RJ, Gordon E, Harkiss D, Twigg JP, Milne P, Jennings Z & Irvine DS 1998 Relative impact of oxidative stress on the functional competence and genomic integrity of human spermatozoa. *Biology of Reproduction* **59** 1037–1046. (doi:10.1095/biolreprod59.5.1037)
- Aitken RJ, Bennetts LE, Sawyer D, Wiklendt AM & King BV 2005 Impact of radio frequency electromagnetic radiation on DNA integrity in the male germline. *International Journal of Andrology* **28** 171–179. (doi:10.1111/j.1365-2605.2005.00531.x)
- Aitken RJ, De Iuliis GN, Gibb Z & Baker MA 2012a The Simmet lecture: new horizons on an old landscape – oxidative stress, DNA damage and apoptosis in the male germ line. *Reproduction in Domestic Animals* **47** (Supplement 4) 7–14. (doi:10.1111/j.1439-0531.2012.02049.x)
- Aitken RJ, Whiting S, De Iuliis GN, McClymont S, Mitchell LA & Baker MA 2012b Electrophilic aldehydes generated by sperm metabolism activate mitochondrial reactive oxygen species generation and apoptosis by targeting succinate dehydrogenase. *Journal of Biological Chemistry* **287** 33048–33060. (doi:10.1074/jbc.M112.366690)
- Aitken RJ, Gibb Z, Mitchell LA, Lambourne SR, Connaughton HS & De Iuliis GN 2012c Sperm motility is lost in vitro as a consequence of mitochondrial free radical production and the generation of electrophilic aldehydes but can be significantly rescued by the presence of nucleophilic thiols. *Biology of Reproduction* **87** 110. (doi:10.1095/biolreprod.112.102020)
- Aitken RJ, Smith TB, Jobling MS, Baker MA & De Iuliis GN 2014 Oxidative stress and male reproductive health. *Asian Journal of Andrology* **16** 31–38. (doi:10.4103/1008-682X.122203)

- Al-Damegh MA 2012 Rat testicular impairment induced by electromagnetic radiation from a conventional cellular telephone and the protective effects of the antioxidants vitamins C and E. *Clinics* **67** 785–792. (doi:10.6061/clinics/2012(07)14)
- Baker MA, Weinberg A, Hetherington L, Villaverde AI, Velkov T, Baell J & Gordon CP 2015 Defining the mechanisms by which the reactive oxygen species by-product, 4-hydroxynonenal, affects human sperm cell function. *Biology of Reproduction* **92** 108. (doi:10.1095/biolreprod.114.126680)
- Balode Z 1996 Assessment of radio-frequency electromagnetic radiation by the micronucleus test in bovine peripheral erythrocytes. *Science of the Total Environment* **180** 81–85. (doi:10.1016/0048-9697(95)04923-1)
- Balzano Q 1999 Exposure metrics for RF epidemiology: cellular phone handsets. *Radiation Protection Dosimetry* **83** 165–169. (doi:10.1093/oxfordjournals.rpd.a032654)
- Bilgili B, Akar A, Avci B & Tuncel OK 2013 Effect of 900 MHz radiofrequency radiation on oxidative stress in rat brain and serum. *Electromagnetic Biology and Medicine* **32** 20–29. (doi:10.3109/15368378.2012.699012)
- Bin-Merfij MM & El-Kott AF 2015 The radioprotective effects of *Moringa oleifera* against mobile phone electromagnetic radiation-induced infertility in rats. *International Journal of Clinical and Experimental Medicine* **8** 12487–12497.
- Bolte JF & Eikelboom T 2012 Personal radiofrequency electromagnetic field measurements in The Netherlands: exposure level and variability for everyday activities, times of day and types of area. *Environment International* **48** 133–142. (doi:10.1016/j.envint.2012.07.006)
- Burlaka A, Tsybulin O, Sidorik E, Lukin S, Polishuk V, Tsehmistrenko S & Yakymenko I 2013 Overproduction of free radical species in embryonal cells exposed to low intensity radiofrequency radiation. *Experimental Oncology* **35** 219–225.
- Chen ZN 2007 *Antennas for Portable Devices*. New York, NY, USA: Wiley.
- Christman MF, Morgan RW, Jacobson FS & Ames BN 1985 Positive control of a regulon for defenses against oxidative stress and some heat-shock proteins in *Salmonella typhimurium*. *Cell* **41** 753–762. (doi:10.1016/S0092-8674(85)80056-8)
- CTIA 2011 Semi-Annual Wireless Industry Survey (CTIA–The Wireless Association). Washington, DC, USA
- Curley SA, Palalon F, Lu X & Koshkina NV 2014 Noninvasive radiofrequency treatment effect on mitochondria in pancreatic cancer cells. *Cancer* **120** 3418–3425. (doi:10.1002/cncr.28895)
- d'Ambrosio G, Massa R, Scarfi MR & Zeni O 2002 Cytogenetic damage in human lymphocytes following GSMK phase modulated microwave exposure. *Bioelectromagnetics* **23** 7–13. (doi:10.1002/bem.93)
- Dasdag S, Ketani MA, Akdag Z, Ersay AR, Sari I, Demirtas OC & Celik MS 1999 Whole-body microwave exposure emitted by cellular phones and testicular function of rats. *Urological Research* **27** 219–223. (doi:10.1007/s002400050113)
- Dasdag S, Zulkuf Akdag M, Aksin F, Yilmaz F, Bashan M, Mutlu Dasdag M & Salih Celik M 2003 Whole body exposure of rats to microwaves emitted from a cell phone does not affect the testes. *Bioelectromagnetics* **24** 182–188. (doi:10.1002/bem.10083)
- Dasdag S, Akdag MZ, Ulukaya E, Uzunlar AK & Ocak AR 2009 Effect of mobile phone exposure on apoptotic glial cells and status of oxidative stress in rat brain. *Electromagnetic Biology and Medicine* **28** 342–354. (doi:10.3109/15368370903206556)
- Dasdag S, Akdag MZ, Erdal ME, Erdal N, Ay OI, Ay ME, Yilmaz SG, Tasdelen B & Yegin K 2015 Long term and excessive use of 900 MHz radiofrequency radiation alter microRNA expression in brain. *International Journal of Radiation Biology* **91** 306–311. (doi:10.3109/09553002.2015.997896)
- De Iuliis GN, Newey RJ, King BV & Aitken RJ 2009a Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro. *PLoS ONE* **4** e6446. (doi:10.1371/journal.pone.0006446)
- De Iuliis GN, Thomson LK, Mitchell LA, Finnie JM, Koppers AJ, Hedges A, Nixon B & Aitken RJ 2009b DNA damage in human spermatozoa is highly correlated with the efficiency of chromatin remodeling and the formation of 8-hydroxy-2'-deoxyguanosine, a marker of oxidative stress. *Biology of Reproduction* **81** 517–524. (doi:10.1095/biolreprod.109.076836)
- De Iuliis GN, King BV & Aitken RJ 2012 Electromagnetic radiation and oxidative stress in the male germ line. In *Studies on Men's Health and Fertility*, pp 119–130. Eds A Agarwal, RJ Aitken & JG Alvarez. New York, NY, USA: Humana Press.
- Demirel S, Doganay S, Turkoz Y, Dogan Z, Turan B & Firat PG 2012 Effects of third generation mobile phone-emitted electromagnetic radiation on oxidative stress parameters in eye tissue and blood of rats. *Cutaneous and Ocular Toxicology* **31** 89–94. (doi:10.3109/15569527.2012.657725)
- Di Carlo A, White N, Guo F, Garrett P & Litovitz T 2002 Chronic electromagnetic field exposure decreases HSP70 levels and lowers cytoprotection. *Journal of Cellular Biochemistry* **84** 447–454. (doi:10.1002/jcb.10036)
- Duan W, Liu C, Zhang L, He M, Xu S, Chen C, Pi H, Gao P, Zhang Y, Zhong M *et al.* 2015 Comparison of the genotoxic effects induced by 50 Hz extremely low-frequency electromagnetic fields and 1800 MHz radiofrequency electromagnetic fields in GC-2 cells. *Radiation Research* **183** 305–314. (doi:10.1667/RR13851.1)
- Durney C 1986 *Radiofrequency Radiation Dosimetry Handbook*, 4th edn. The University of Utah, Salt Lake City, UT, USA
- Erogul O, Oztas E, Yildirim I, Kir T, Aydur E, Komesli G, Irkilata HC, Irmak MK & Peker AF 2006 Effects of electromagnetic radiation from a cellular phone on human sperm motility: an in vitro study. *Archives of Medical Research* **37** 840–843. (doi:10.1016/j.arcmed.2006.05.003)
- Falzone N, Huyser C, Becker P, Leszczynski D & Franken DR 2011 The effect of pulsed 900-MHz GSM mobile phone radiation on the acrosome reaction, head morphometry and zona binding of human spermatozoa. *International Journal of Andrology* **34** 20–26. (doi:10.1111/j.1365-2605.2010.01054.x)
- Fejes I, Zavaczki Z, Szollosi J, Koloszar S, Daru J, Kovacs L & Pal A 2005 Is there a relationship between cell phone use and semen quality? *Archives of Andrology* **51** 385–393. (doi:10.1080/014850190924520)
- French PW, Penny R, Laurence JA & McKenzie DR 2001 Mobile phones, heat shock proteins and cancer. *Differentiation* **67** 93–97. (doi:10.1046/j.1432-0436.2001.670401.x)
- Friedman J, Kraus S, Hauptman Y, Schiff Y & Seger R 2007 Mechanism of short-term ERK activation by electromagnetic fields at mobile phone frequencies. *Biochemical Journal* **405** 559–568. (doi:10.1042/BJ20061653)
- Furtado-Filho OV, Borba JB, Dallegrave A, Pizzolato TM, Henriques JA, Moreira JC & Saffi J 2014 Effect of 950 MHz UHF electromagnetic radiation on biomarkers of oxidative damage, metabolism of UFA and antioxidants in the livers of young rats of different ages. *International Journal of Radiation Biology* **90** 159–168. (doi:10.3109/09553002.2013.817697)
- Gabriel S, Lau RW & Gabriel C 1996 The dielectric properties of biological tissues: III. Parametric models for the dielectric spectrum of tissues. *Physics in Medicine and Biology* **41** 2271–2293. (doi:10.1088/0031-9155/41/11/003)
- Gajda GB, McNamee JP, Thansandote A, Boonpanyarak S, Lemay E & Bellier PV 2002 Cylindrical waveguide applicator for in vitro exposure of cell culture samples to 1.9-GHz radiofrequency fields. *Bioelectromagnetics* **23** 592–598. (doi:10.1002/bem.10055)
- Ghanbari M, Mortazavi SB, Khavanin A & Khazaei M 2013 The effects of cell phone waves (900 MHz-GSM Band) on sperm parameters and total antioxidant capacity in rats. *International Journal of Fertility & Sterility* **7** 21–28.
- Gharagozloo P & Aitken RJ 2011 The role of sperm oxidative stress in male infertility and the significance of oral antioxidant therapy. *Human Reproduction* **26** 1628–1640. (doi:10.1093/humrep/der132)
- Gorpinchenko I, Nikitin O, Banyra O & Shulyak A 2014 The influence of direct mobile phone radiation on sperm quality. *Central European Journal of Urology* **67** 65–71. (doi:10.5173/cej.2014.01.art14)
- Guney M, Ozguner F, Oral B, Karahan N & Mungan T 2007 900 MHz radiofrequency-induced histopathologic changes and oxidative stress in rat endometrium: protection by vitamins E and C. *Toxicology and Industrial Health* **23** 411–420. (doi:10.1177/0748233707080906)
- Hossmann KA & Hermann DM 2003 Effects of electromagnetic radiation of mobile phones on the central nervous system. *Bioelectromagnetics* **24** 49–62. (doi:10.1002/bem.10068)
- Hou Q, Wang M, Wu S, Ma X, An G, Liu H & Xie F 2015 Oxidative changes and apoptosis induced by 1800-MHz electromagnetic radiation in NIH/3T3 cells. *Electromagnetic Biology and Medicine* **34** 85–92. (doi:10.3109/15368378.2014.900507)

- Imai N, Kawabe M, Hikage T, Nojima T, Takahashi S & Shirai T 2011 Effects on rat testis of 1.95-GHz W-CDMA for IMT-2000 cellular phones. *Systems Biology in Reproductive Medicine* **57** 204–209. (doi:10.3109/19396368.2010.544839)
- Iorio R, Delle Monache S, Bennato F, Di Bartolomeo C, Scrimaglio R, Cinque B & Colonna RC 2011 Involvement of mitochondrial activity in mediating ELF-EMF stimulatory effect on human sperm motility. *Bioelectromagnetics* **32** 15–27. (doi:10.1002/bem.20602)
- Irmak MK, Fadillioglu E, Gulec M, Erdogan H, Yagmurca M & Akyol O 2002 Effects of electromagnetic radiation from a cellular telephone on the oxidant and antioxidant levels in rabbits. *Cell Biochemistry and Function* **20** 279–283. (doi:10.1002/cbf.976)
- Jin Z, Zong C, Jiang B, Zhou Z, Tong J & Cao Y 2012 The effect of combined exposure of 900 MHz radiofrequency fields and doxorubicin in HL-60 cells. *PLoS ONE* **7** e46102. (doi:10.1371/journal.pone.0046102)
- Jones R, Mann T & Sherins R 1979 Peroxidative breakdown of phospholipids in human spermatozoa, spermicidal properties of fatty acid peroxides, and protective action of seminal plasma. *Fertility and Sterility* **31** 531–537. (doi:10.1016/S0015-0282(16)43999-3)
- Kahya MC, Naziroglu M & Cig B 2014 Selenium reduces mobile phone (900 MHz)-induced oxidative stress, mitochondrial function, and apoptosis in breast cancer cells. *Biological Trace Element Research* **160** 285–293. (doi:10.1007/s12011-014-0032-6)
- Kesari KK & Behari J 2012 Evidence for mobile phone radiation exposure effects on reproductive pattern of male rats: role of ROS. *Electromagnetic Biology and Medicine* **31** 213–222. (doi:10.3109/15368378.2012.700292)
- Kesari KK, Kumar S & Behari J 2011 Effects of radiofrequency electromagnetic wave exposure from cellular phones on the reproductive pattern in male Wistar rats. *Applied Biochemistry and Biotechnology* **164** 546–559. (doi:10.1007/s12010-010-9156-0)
- Khalil AM, Abu Khadra KM, Aljaberi AM, Gagaa MH & Issa HS 2014 Assessment of oxidant/antioxidant status in saliva of cell phone users. *Electromagnetic Biology and Medicine* **33** 92–97. (doi:10.3109/15368378.2013.783855)
- Koppers AJ, De Iuliis GN, Finnie JM, McLaughlin EA & Aitken RJ 2008 Significance of mitochondrial reactive oxygen species in the generation of oxidative stress in spermatozoa. *Journal of Clinical Endocrinology and Metabolism* **93** 3199–3207. (doi:10.1210/jc.2007-2616)
- Koppers AJ, Garg ML & Aitken RJ 2010 Stimulation of mitochondrial reactive oxygen species production by unesterified, unsaturated fatty acids in defective human spermatozoa. *Free Radical Biology and Medicine* **48** 112–119. (doi:10.1016/j.freeradbiomed.2009.10.033)
- Kudin AP, Bimpong-Buta NY, Vielhaber S, Elger CE & Kunz WS 2004 Characterization of superoxide-producing sites in isolated brain mitochondria. *Journal of Biological Chemistry* **279** 4127–4135. (doi:10.1074/jbc.M310341200)
- La Vignera S, Condorelli RA, Vicari E, D'Agata R & Calogero AE 2012 Effects of the exposure to mobile phones on male reproduction: a review of the literature. *Journal of Andrology* **33** 350–356. (doi:10.2164/jandrol.111.014373)
- Lenzi A, Gandini L, Maresca V, Rago R, Sgro P, Dondero F & Picardo M 2000 Fatty acid composition of spermatozoa and immature germ cells. *Molecular Human Reproduction* **6** 226–231. (doi:10.1093/molehr/6.3.226)
- Lerchl A 2013 Electromagnetic pollution: another risk factor for infertility, or a red herring? *Asian Journal of Andrology* **15** 201–203. (doi:10.1038/aja.2012.104)
- Li HW, Yao K, Jin HY, Sun LX, Lu DQ & Yu YB 2007 Proteomic analysis of human lens epithelial cells exposed to microwaves. *Japanese Journal of Ophthalmology* **51** 412–416. (doi:10.1007/s10384-007-0483-9)
- Lin JC 1976 Interaction of two cross-polarized electromagnetic waves with mammalian cranial structures. *IEEE Transactions on Biomedical Engineering* **23** 371–375. (doi:10.1109/TBME.1976.324646)
- Liu W, Zheng X, Qu Z, Zhang M, Zhou C, Ma L & Zhang Y 2012 Effect of 935-MHz phone-simulating electromagnetic radiation on endometrial glandular cells during mouse embryo implantation. *Journal of Huazhong University of Science and Technology* **32** 755–759. (doi:10.1007/s11596-012-1030-6)
- Liu C, Duan W, Xu S, Chen C, He M, Zhang L, Yu Z & Zhou Z 2013a Exposure to 1800 MHz radiofrequency electromagnetic radiation induces oxidative DNA base damage in a mouse spermatocyte-derived cell line. *Toxicology Letters* **218** 2–9. (doi:10.1016/j.toxlet.2013.01.003)
- Liu C, Gao P, Xu SC, Wang Y, Chen CH, He MD, Yu ZP, Zhang L & Zhou Z 2013b Mobile phone radiation induces mode-dependent DNA damage in a mouse spermatocyte-derived cell line: a protective role of melatonin. *International Journal of Radiation Biology* **89** 993–1001. (doi:10.3109/09553002.2013.811309)
- Liu K, Li Y, Zhang G, Liu J, Cao J, Ao L & Zhang S 2014 Association between mobile phone use and semen quality: a systemic review and meta-analysis. *Andrology* **2** 491–501. (doi:10.1111/j.2047-2927.2014.00205.x)
- Liu Q, Si T, Xu X, Liang F, Wang L & Pan S 2015 Electromagnetic radiation at 900 MHz induces sperm apoptosis through bcl-2, bax and caspase-3 signaling pathways in rats. *Reproductive Health* **12** 65. (doi:10.1186/s12978-015-0062-3)
- Lord T & Aitken RJ 2015 Fertilization stimulates 8-hydroxy-2'-deoxyguanosine repair and antioxidant activity to prevent methylation in the embryo. *Developmental Biology* **406** 1–13. (doi:10.1016/j.ydbio.2015.07.024)
- Lu YS, Huang BT & Huang YX 2012 Reactive oxygen species formation and apoptosis in human peripheral blood mononuclear cell induced by 900 MHz mobile phone radiation. *Oxidative Medicine and Cellular Longevity* **2012** 740280. (doi:10.1155/2012/740280)
- Mailankot M, Kunnath AP, Jayalekshmi H, Koduru B & Valsalan R 2009 Radio frequency electromagnetic radiation (RF-EMR) from GSM (0.9/1.8GHz) mobile phones induces oxidative stress and reduces sperm motility in rats. *Clinics* **64** 561–565. (doi:10.1590/S1807-59322009000600011)
- Manta AK, Stravopodis DJ, Papassideris IS & Margaritis LH 2014 Reactive oxygen species elevation and recovery in Drosophila bodies and ovaries following short-term and long-term exposure to DECT base EMF. *Electromagnetic Biology and Medicine* **33** 118–131. (doi:10.3109/15368378.2013.791991)
- Marchionni I, Paffi A, Pellegrino M, Liberti M, Apollonio F, Abeti R, Fontana F, D'Inzeo G & Mazzanti M 2006 Comparison between low-level 50 Hz and 900 MHz electromagnetic stimulation on single channel ionic currents and on firing frequency in dorsal root ganglion isolated neurons. *Biochimica et Biophysica Acta* **1758** 597–605. (doi:10.1016/j.bbame.2006.03.014)
- Markov M & Grigoriev Y 2015 Protect children from EMF. *Electromagnetic Biology and Medicine* **34** 251–256. (doi:10.3109/15368378.2015.1077339)
- Martino CF & Castello PR 2011 Modulation of hydrogen peroxide production in cellular systems by low level magnetic fields. *PLoS ONE* **6** e22753. (doi:10.1371/journal.pone.0022753)
- Masuda H, Sanchez S, Dulou PE, Haro E, Anane R, Billaudel B, Leveque P & Veyret B 2006 Effect of GSM-900 and -1800 signals on the skin of hairless rats. I: 2-hour acute exposures. *International Journal of Radiation Biology* **82** 669–674. (doi:10.1080/09553000600930079)
- Meral I, Mert H, Mert N, Deger Y, Yoruk I, Yetkin A & Keskin S 2007 Effects of 900-MHz electromagnetic field emitted from cellular phone on brain oxidative stress and some vitamin levels of guinea pigs. *Brain Research* **1169** 120–124. (doi:10.1016/j.brainres.2007.07.015)
- Moazamian R, Polhemus A, Connaughton H, Fraser B, Whiting S, Gharagozloo P & Aitken RJ 2015 Oxidative stress and human spermatozoa: diagnostic and functional significance of aldehydes generated as a result of lipid peroxidation. *Molecular Human Reproduction* **21** 502–515. (doi:10.1093/molehr/gav014)
- Murphy MP 2009 How mitochondria produce reactive oxygen species. *Biochemical Journal* **417** 1–13. (doi:10.1042/BJ20081386)
- Nahomi RB, DiMauro MA, Wang B & Nagaraj RH 2015 Identification of peptides in human Hsp20 and Hsp27 that possess molecular chaperone and anti-apoptotic activities. *Biochemical Journal* **465** 115–125. (doi:10.1042/BJ20140837)
- Ozguner F, Altinbas A, Ozaydin M, Dogan A, Vural H, Kisioglu AN, Cesur G & Yildirim NG 2005 Mobile phone-induced myocardial oxidative stress: protection by a novel antioxidant agent caffeic acid phenethyl ester. *Toxicology and Industrial Health* **21** 223–230. (doi:10.1191/0748233705th228oa)
- Ozlem Nisbet H, Nisbet C, Akar A, Cevik M & Karayigit MO 2012 Effects of exposure to electromagnetic field (1.8/0.9 GHz) on testicular function and structure in growing rats. *Research in Veterinary Science* **93** 1001–1005. (doi:10.1016/j.rvsc.2011.10.023)
- Ozorak A, Naziroglu M, Celik O, Yuksel M, Ozcelik D, Ozkaya MO, Cetin H, Kahya MC & Kose SA 2013 Wi-Fi (2.45 GHz)- and mobile

- phone (900 and 1800 MHz)-induced risks on oxidative stress and elements in kidney and testis of rats during pregnancy and the development of offspring. *Biological Trace Element Research* **156** 221–229. (doi:10.1007/s12011-013-9836-z)
- Panagopoulos DJ, Messini N, Karabarbounis A, Philippetis AL & Margaritis LH 2000 A mechanism for action of oscillating electric fields on cells. *Biochemical and Biophysical Research Communications* **272** 634–640. (doi:10.1006/bbrc.2000.2746)
- Panagopoulos DJ, Karabarbounis A & Margaritis LH 2002 Mechanism for action of electromagnetic fields on cells. *Biochemical and Biophysical Research Communications* **298** 95–102. (doi:10.1016/S0006-291X(02)02393-8)
- Panagopoulos DJ, Chavdoula ED & Margaritis LH 2010 Bioeffects of mobile telephony radiation in relation to its intensity or distance from the antenna. *International Journal of Radiation Biology* **86** 345–357. (doi:10.3109/095533000903567961)
- Panagopoulos DJ, Johansson O & Carlo GL 2015 Polarization: a key difference between man-made and natural electromagnetic fields, in regard to biological activity. *Scientific Reports* **5** 14914. (doi:10.1038/srep14914)
- Perry SW, Norman JP, Barbieri J, Brown EB & Gelbard HA 2011 Mitochondrial membrane potential probes and the proton gradient: a practical usage guide. *Biotechniques* **50** 98–115. (doi:10.2144/000113610)
- Polla BS, Kantengwa S, Francois D, Salvioli S, Franceschi C, Marsac C & Cossarizza A 1996 Mitochondria are selective targets for the protective effects of heat shock against oxidative injury. *PNAS* **93** 6458–6463. (doi:10.1073/pnas.93.13.6458)
- Quinlan CL, Perevoshchikova IV, Hey-Mogensen M, Orr AL & Brand MD 2013 Sites of reactive oxygen species generation by mitochondria oxidizing different substrates. *Redox Biology* **1** 304–312. (doi:10.1016/j.redox.2013.04.005)
- Redmayne M, Smith E & Abramson MJ 2011 Adolescent in-school cellphone habits: a census of rules, survey of their effectiveness, and fertility implications. *Reproductive Toxicology* **32** 354–359. (doi:10.1016/j.reprotox.2011.08.006)
- Roberts JA, Yaya LH & Manolis C 2014 The invisible addiction: cell-phone activities and addiction among male and female college students. *Journal of Behavioral Addictions* **3** 254–265. (doi:10.1556/JBA.3.2014.015)
- Sanders AP & Joines WT 1984 The effects of hyperthermia and hyperthermia plus microwaves on rat brain energy metabolism. *Bioelectromagnetics* **5** 63–70. (doi:10.1002/bem.2250050107)
- Sommer AM, Grote K, Reinhardt T, Streckert J, Hansen V & Lerchl A 2009 Effects of radiofrequency electromagnetic fields (UMTS) on reproduction and development of mice: a multi-generation study. *Radiation Research* **171** 89–95. (doi:10.1667/RR1460.1)
- Storey BT 2008 Mammalian sperm metabolism: oxygen and sugar, friend and foe. *International Journal of Developmental Biology* **52** 427–437. (doi:10.1387/ijdb.072522bs)
- Singer TM, Lambert IB, Williams A, Douglas GR & Yauk CL 2006 Detection of induced male germline mutation: correlations and comparisons between traditional germline mutation assays, transgenic rodent assays and expanded simple tandem repeat instability assays. *Mutation Research* **598** 164–193. (doi:10.1016/j.mrfmmm.2006.01.017)
- Tas M, Dasdag S, Akdag MZ, Cirit U, Yegin K, Seker U, Ozmen MF & Eren LB 2014 Long-term effects of 900 MHz radiofrequency radiation emitted from mobile phone on testicular tissue and epididymal semen quality. *Electromagnetic Biology and Medicine* **33** 216–222. (doi:10.3109/15368378.2013.801850)
- Tattersall JE, Scott IR, Wood SJ, Nettell JJ, Bevir MK, Wang Z, Somasiri NP & Chen X 2001 Effects of low intensity radiofrequency electromagnetic fields on electrical activity in rat hippocampal slices. *Brain Research* **904** 43–53. (doi:10.1016/S0006-8993(01)02434-9)
- Tosic J & Walton A 1950 Metabolism of spermatozoa. The formation and elimination of hydrogen peroxide by spermatozoa and effects on motility and survival. *Biochemical Journal* **47** 199–212. (doi:10.1042/bj0470199)
- Trosic I, Matausic-Pisl M, Pavicic I & Marjanovic AM 2013 Histological and cytological examination of rat reproductive tissue after short-time intermittent radiofrequency exposure. *Arhiv za Higijenu Rada i Toksikologiju* **64** 513–519.
- Tumkaya L, Kalkan Y, Bas O & Yilmaz A 2013 Mobile phone radiation during pubertal development has no effect on testicular histology in rats. *Toxicology and Industrial Health* **32** 328–336. (doi:10.1177/0748233713500820)
- Valbonesi P, Franzellitti S, Bersani F, Contini A & Fabbri E 2014 Effects of the exposure to intermittent 1.8 GHz radio frequency electromagnetic fields on HSP70 expression and MAPK signaling pathways in PC12 cells. *International Journal of Radiation Biology* **90** 382–391. (doi:10.3109/09553002.2014.892225)
- Wdowiak A, Wdowiak L & Wiktor H 2007 Evaluation of the effect of using mobile phones on male fertility. *Annals of Agricultural and Environmental Medicine* **14** 169–172.
- Williams AC & Ford WC 2001 The role of glucose in supporting motility and capacitation in human spermatozoa. *Journal of Andrology* **22** 680–695.
- Yan JG, Agresti M, Bruce T, Yan YH, Granlund A & Matloub HS 2007 Effects of cellular phone emissions on sperm motility in rats. *Fertility and Sterility* **88** 957–964. (doi:10.1016/j.fertnstert.2006.12.022)
- Yao K, Wu W, Wang K, Ni S, Ye P, Yu Y, Ye J & Sun L 2008 Electromagnetic noise inhibits radiofrequency radiation-induced DNA damage and reactive oxygen species increase in human lens epithelial cells. *Molecular Vision* **14** 964–969.
- Zalata A, El-Samanoudy AZ, Shaalan D, El-Baiomy Y & Mostafa T 2015 In vitro effect of cell phone radiation on motility, DNA fragmentation and clusterin gene expression in human sperm. *International Journal of Fertility and Sterility* **9** 129–136.
- Zhang L, Yu L & Yu CA 1998 Generation of superoxide anion by succinate-cytochrome c reductase from bovine heart mitochondria. *Journal of Biological Chemistry* **273** 33972–33976. (doi:10.1074/jbc.273.51.33972)
- Zhao L, Peng RY, Wang SM, Wang LF, Gao YB, Dong J, Li X & Su ZT 2012 Relationship between cognition function and hippocampus structure after long-term microwave exposure. *Biomedical and Environmental Sciences* **25** 182–188.

Received 9 March 2016

First decision 8 April 2016

Revised manuscript received 2 August 2016

Accepted 2 September 2016