Disorganization of the germ cell pool leads to primary ovarian insufficiency

Ikko Kawashima* and Kazuhiro Kawamura*

Department of Advanced Reproductive Medicine, St. Marianna University School of Medicine, Kawasaki City, Kanagawa, Japan

Correspondence should be addressed to K Kawamura; Email: kazuhiironanami@gmail.com

*(I Kawashima and K Kawamura contributed equally to this work)

Abstract

The mammalian ovary is an organ that controls female germ cell development, storing them and releasing mature oocytes for transporting to the oviduct. During the fetal stage, female germ cells change from a proliferative state to meiosis before forming follicles with the potential for the growth of surrounding somatic cells. Understanding of molecular and physiological bases of germ cell development in the fetal ovary contributed not only to the elucidation of genetic disorders in primary ovarian insufficiency (POI), but also to the advancement of novel treatments for patients with POI. Accumulating evidence indicates that mutations in *NOBOX*, *DAZI* and *FIGLA* genes are associated with POI. In addition, cell biology studies revealed the important roles of these genes as essential translational factors for germ cell development. Recent insights into the role of the PI3K (phosphatidylinositol 3-kinase)-AKT signaling pathway in primordial follicle activation allowed the development of a new infertility treatment, IVA (*in vitro* activation), leading to successful pregnancy/delivery in POI patients. Furthermore, elucidation of genetic dynamics underlying female germ cell development could allow regeneration of oocytes from ES (embryonic stem)/iPS (induced pluripotent stem) cells in mammals. The purpose of this review is to summarize basic findings related to female germ cell development and potential clinical implications, especially focusing on POI etiologies. We also summarize evolving new POI therapies based on IVA as well as oocyte regeneration.

Reproduction (2017) **153** R205–R213

Introduction

During the fetal stage, primordial germ cells (PGC) migrate into the fetal ovary. At 5 weeks after fertilization, the human ovary contains an estimated amount of 700–1300 germ cells (Witschi 1948). The number of germ cell reaches an estimated amount of 600,000 cells between 16 and 20 weeks of pregnancy (Barker 1963). In mouse ovaries, there are around 50 primordial germ cells at 7.5-day post coitus (dpc) (Ginsburg *et al.* 1990, MacGregor *et al.* 1995) and germ cell number peaks at 13.5 dpc, reaching about 20,000 cells (Tam & Snow 1981). For both species, variation patterns are the same, although cell number and timing are different. After increases in numbers, PGCs form colonies in the ovary. Once the PGCs form cysts, they initiate meiotic division and cell growth at 11–12 weeks post-fertilization in women (Gondos *et al.* 1986, Motta *et al.* 1997) and 12.5 dpc in mice (Tingen *et al.* 2009, Jameson *et al.* 2012). Meiotic cells were arrested in the diplotene stage of prophase I during germ cell cyst breakdown accompanied by reduction in cell numbers. Cyst breakdown is associated with the individualization of germ cells to form primordial oocytes, around 16 weeks

of pregnancy in humans and at the time of birth in mice (Grive & Freiman 2015). Then, oocyte meiosis arrests in the dictyate stage of prophase I. Individual oocyte interacts with ovarian somatic cells to form primordial follicles in the ovary. Primordial follicles remain dormant for months and years and only a small fraction of them become activated under regulation of the kit ligand (Zhang *et al.* 2014). Once primordial follicle is activated, they grow into primary, secondary and antral follicle stages, accompanied by growth and maturation of the oocyte before ovulation of the mature oocyte (Richards & Pangas 2010).

Disorganization of germ cell development and primordial follicle formation likely cause a decline in the germ cell pool, resulting in decreases of follicles in women with the most severe cases of primary ovarian insufficiency (POI). POI, also known as premature ovarian failure (POF), is defined as menopause before 40 years of age associated with elevated serum gonadotropin levels (Coulam *et al.* 1986, Simpson & Rajkovic 1999). POI patients show symptoms of estrogen deficiency and infertility due to lack of follicle growth and ovulation. These patients are resistant to traditional gonadotropin treatment, and thus, oocyte donation

is the only established treatment option. In terms of etiology, POI affects 1% of women and several different mechanisms can lead to ovarian dysfunction including genetic mutations, autoimmune diseases, iatrogenesis and other abnormalities.

Recently, a novel clinical approach, *in vitro* activation (IVA) of primordial follicle (Kawamura 2013), has been developed and succeeded in allowing POI patients to have their own genetic children. However, IVA method only activates dormant follicles and it cannot be applied to POI patients without primordial follicles. In order to treat POI patient lacking follicles, it is necessary to derive oocytes from stem cells. Remarkably, investigators attempted to derive oocytes as models for future POI therapy using murine iPS/ES cells (Hayashi *et al.* 2012, Hikabe *et al.* 2016) or human female germ stem cells (White *et al.* 2012). The present review focuses on the recent knowledge of germ cell development to facilitate future design of POI therapies.

Regulatory mechanisms underlying oocyte development and follicle formation

Germ cell proliferation and cyst formation

In human, germ cells migrate into the fetal ovary at 3 weeks after fertilization. These cells initiate proliferation and cyst formation before reaching maximum numbers in the ovary at 20 weeks of pregnancy. Germ cell growth and cyst formation are dependent upon extracellular matrixes in the basement membrane secreted by ovarian somatic cells (Heeren *et al.* 2015). Indeed, immunohistochemical studies show that dividing germ cells with Ki-67 positive signals are present in the ovary until postpartum (Stoop *et al.* 2005). Activin is known as a factor capable of inducing proliferation of these cells (Martins da Silva *et al.* 2004) (Fig. 1). As part of extracellular matrixes, collagen type IV, fibronectin and laminin have been found in germ cell cysts (Heeren *et al.* 2015).

In mice, germ cells mainly proliferate in the fetal ovary. Cell division and cyst formation of germ cells last only a few days from 7.5 dpc to 13.5 dpc (Tam & Snow 1981). Bone morphogenetic proteins (BMPs) of the transforming growth factor b (TGFb) family, including BMP2, 4 and 7, are important triggers for germ cell proliferation based on cell culture experiments (Pesce *et al.* 2002, Puglisi *et al.* 2004) (Fig. 1). In *in vivo* studies, *Bmp7* null mice showed a reduction of germ cells at 11.5 dpc (Ross *et al.* 2007). RNA-seq studies showed that pre-granulosa cells express *Bmp2* and *Bmp4* during 11.5–13.5 dpc (Jameson *et al.* 2012) (Fig. 1). These results suggest that BMPs probably induce proliferation of germ cell *in vivo*. Moreover, *in vitro* culture experiments showed that Kit Ligand (KITL) (Dolci *et al.* 1993), basic fibroblast growth factor (bFGF) (Matsui *et al.* 1992), tumor necrosis factor- α (TNF α) (Kawase *et al.* 1994) and pituitary adenylate cyclase activating peptide (PACAP) (Pesce *et al.* 1996)

Figure 1 Temporal contribution of secretory factors during human and mouse germ cell development. Primordial germ cell proliferates in the fetal ovary at 10.5 dpc in mice and at approximately 5 weeks of gestation in human. These cell proliferations are induced by BMPs in mice, but Activin A in human. FST maintains germ cell proliferation and cyst formation. Proliferating cells colonize on the basement membrane shaped by somatic cells in fetal ovary at 12.5 dpc in mice and at 11–20 weeks of gestation in human. Colonized germ cells initiate meiotic division following RA and Jagged (JAG) stimulations in mouse and RA stimulation in human. During meiosis, germ cell cyst breaks down and oogonia grow at 17.5 dpc in mice and at 12–22 weeks of gestation in human. These processes are induced by Jagged and KITL in mouse, but by unknown factors in human. Primordial follicles form from post natal day 1 (PND1) in mice and from 16 weeks of gestation in human. Activation of primordial follicles to develop primary follicles is mediated by the PI3K-Akt signaling pathway.

induce germ cell division. These reports indicate that germ cell proliferation may be regulated by multiple factors during fetal ovary development. Follistatin (FST), an antagonist of activin, was shown to maintain germ cell number and cyst (Fig. 1). Although mice lacking *Fst* have comparable number of germ cells in fetal ovary at 15.5 dpc as wild mice, germ cells decreased at 16.5 dpc and are completely depleted at birth (Yao *et al.* 2004). The treatment of mouse fetal ovaries with FST288, as activin-neutralizing isoform of FST, led to the inhibition

of cyst breakdown and germ cell growth, together with the reduction of germ cell number (Wang *et al.* 2015). These results indicate that FST controls germ cell proliferation and germ cell cyst formation (Fig. 1).

In human, the relationship between these secreted factors inducing germ cell proliferation and POI has not been reported. It is well known that patients with Turner's syndrome (complete X (45X) or partial monosomy X (46X, i)) have small numbers of germ cells at 20 weeks of gestation (Modi & Bhartiya 2003). About 90% of patients with 45X and 46X, i show primary amenorrhea (Ogata & Matsuo 1995, James *et al.* 1998) due to a decline in germ cell proliferation in the fetal ovary. Therefore, these secreted factors are likely deleted or down regulated in patients with Turner's syndrome. However, BMPs, FST and other secreted factors involved in the induction of germ cell proliferation are not located in X chromosome. Mutations of X chromosome-linked gene in fragile X mental retardation 1 (FMR1), which acts to initiate DNA replication during meiotic division (Schmucker *et al.* 1996), exhibit the POI phenotype due to the expansion of a CGG trinucleotide repeat (Pietrobono *et al.* 2005, Sullivan *et al.* 2005, Terracciano *et al.* 2005). Because FMR1 regulates mRNA trafficking from the nucleus to the cytoplasm, it is interesting to investigate changes in RNA trafficking during germ cell development.

Coordinating meiotic progression in the ovary

In female mammals, there is only one round of meiosis in germ cells during lifetime. At 11–12 weeks of gestation, germ cells in human fetal ovary initiate meiotic division, starting from the S-phase until the diplotene stage (Fig. 1). Many studies in mice revealed various intracellular factors regulating meiosis. Before cell-cycle changes, germ cells exhibit increases in RNA levels for deleted in Azoospermia-Like (*Dazl*) and DEAD-box helicase 4 (*Ddx4* (alias *Vasa*)) (Castrillon *et al.* 2000, Medrano *et al.* 2002). The DAZL protein mediates transcription of synaptonemal complex proteins (SCPs) including SCP3 and SCP1 (Reynolds *et al.* 2007). SCPs are meiosis-related proteins essential for chromosome segregation via chromosome pairing and meiotic recombination (Pittman *et al.* 1998). *Scp1* knock down in cultured fetal ovaries showed decreases in diplotene stage oogonium (Paredes *et al.* 2005), suggesting that diplotene stage progression is supported by SCP1. In contrast, studies in stimulated by retinoic acid 8 (*Stra8*) null mice showed no effect in meiotic arrest of oocytes and follicle development, but ovulated oocytes were abnormal (Doksin *et al.* 2013). Despite these studies, detailed mechanisms of these factors are not fully understood. For example, retinoic acid (RA) can induce meiosis genes expression, such as *Scp3* and *Stra8* (Oulad-Abdelghani *et al.* 1996, Menke *et al.* 2003, Tedesco *et al.* 2013) (Fig. 1). Moreover, a recent study showed that novel meiosis regulatory factors, *Jagged 1* and 2, were expressed in fetal gonads, whereas their

receptors NOTCH 1-3 were expressed in germ cells, during 11.5-14.5 dpc in mice and their inhibition led to a decline of *Stra8* gene expression and suppression of germ cell progression to the diplotene stage. These findings suggested the important roles of NOTCH receptor signaling in meiosis progression (Feng *et al.* 2014) (Fig. 1). Furthermore, meiotic progression also needs secreting factors from ovarian somatic cells based on the studies of oocyte regeneration from ES/iPS cells in mice. Further research will be required to fully understand the intraovarian mechanisms underlying meiotic division.

In contrast to accumulating knowledge on meiotic progression in mice, study in human is limited. Recent review implies the contribution of *DAZL* mutation in pathophysiology in POI (Rosario *et al.* 2016) based on a correlation between several common polymorphisms within the *DAZL* gene and onset age of POI (Tung *et al.* 2006a,b). Comprehensive studies focused on DAZL-induced meiotic cascade would open a new understanding on human meiotic progression and elucidate molecular mechanisms underlying human meiotic progression to allow the development of new methods to regenerate oocytes.

Germ cell attrition and continuing development

In human, germ cell death is observed from 8 weeks of gestation, and the number of germ cells begins to decline after a peak at 20 weeks in the fetal ovary. During this decline, the germ cell cyst breaks down, and germ cells are separated from the ovarian somatic cells followed by oogonia growth (Fig. 1). During oogonia growth, the newly formed oogonia increase their volume with the elevation of the numbers of mitochondria and ribosome from 14.5 dpc to 4 days after birth in mice (Fig. 1). The germ cell cyst breakdown is accompanied by apoptosis and autophagy resulting in rapid decreases in germ cell numbers (Rodrigues *et al.* 2009). In mice, *Jagged 1/2* and *KITL* are known to stimulate these processes (Fig. 1). Ovarian somatic cell (pre-granulosa cell)-specific deletion of *Notch2*, the receptor for *Jagged 1/2*, in mice showed the maintenance of oocyte numbers and multi-oocyte follicles at 3 weeks of age caused by defects in cyst breakdown (Xu & Gridley 2013) (Fig. 1). Using 17.5 dpc ovaries, organ culture studies indicated that *KITL* enhances germ cell cyst breakdown and oogonia growth (Jones & Pepling 2013). Although molecular mechanism underlying germ cell attrition is unknown, a recent murine study indicated that dominant oogonia receive cytoplasm, mitochondria, microtubules and centrosome from neighboring oogonia via intercellular bridges and gaps (Lei & Spradling 2016). Neighboring germ cells are connected to each other via a channel protein, *TEX14* (Greenbaum *et al.* 2009). In human, fetal ovary after 12 weeks of gestation expresses channel protein *TEX14* (Hormard *et al.* 2009), suggesting that the growth of oogonia may be induced by material transport from neighbor oogonia via the channels.

Further studies are needed to elucidate the underlying intercellular mechanisms.

No evidence of POI associated with defects in this process has been reported. However, *Fst* mutant mice with FST288 skipping exon 6 showed POI phenotypes with increased germ cell attrition, leading to decreased follicle and pups numbers (Kimura *et al.* 2010). Thus, further research on the mutation of *Fst* and factors related to oögonia growth could elucidate molecular defects in POI patients.

Primordial follicle formation

In human, primordial follicles are found in the ovary from 16 weeks of gestation (Fig. 1). After cyst breakdown, oögonia are surrounded by somatic cells, named as pre-granulosa cells. The complex of oögonia and pre-granulosa cells form 1,000,000–2,000,000 primordial follicles immediately after birth (Strauss & William 2004). Morphologically, cyst breakdown and subsequent cooperation of germ cells and ovarian somatic cells form primordial follicles. Factor in the germline alpha (*Figla*) is essential for follicle formation. The expression of *Figla* is observed at 17.5–19.5 dpc when meiosis has been completed (Liang *et al.* 1997, Soya *et al.* 2000). Of note, a study using *Figla* null mice showed the absence of primordial follicle formation at birth (Soya *et al.* 2000).

In human, *FIGLA* is expressed in fetal ovary after 15 weeks of gestation (Bayne *et al.* 2004), and thus suggesting its involvement in follicle formation. Importantly, mutation in *FIGLA* gene at exon 1 or 3 is associated with POI (Zhao *et al.* 2008, Tosh *et al.* 2015). In Chinese population, *FIGLA* mutation is detected in 4% of POI patients (4 out of 100). The onset of POI in patients with *FIGLA* mutation is relatively young (22, 27, 32 and 36 years of age) and the abnormal phenotype is observed only in ovaries (Zhao *et al.* 2008).

Follicle development

Primordial follicles keep oocytes in a dormant condition for a long time (Franchi & Mandl 1962, Weakley 1967). Studies using genetically modified mice indicate that oocyte regulates follicle survival. For example, mice lacking genes encoding oocyte transcriptional factor *Sohlh1* (spermatogenesis and oögenesis specific basic helix-loop-helix 1), *Lhx8* (LIM Homeobox 8) or *Sohlh2* showed normal primordial development, but had no follicles left at 3 weeks of age (Pangas *et al.* 2006, Choi *et al.* 2008). Oocyte-specific deletion of the homeobox gene *Nobox* in mice showed follicle-like structure lacking oocytes in the ovary at 2 weeks of age (Rajkovic *et al.* 2004). However, it is unclear how these transcriptional factors support follicular survival. On the other hand, follicular granulosa cells are also known to regulate oocyte survival. FOXL2 (Forkhead Box L2) is a

transcription factor acting in granulosa cells at primordial and primary stage. Follicles in *Foxl2* null mice showed growth arrest and all follicles were depleted at 16 weeks after birth (Schmidt *et al.* 2003).

Based on the analyses of 96 cases of POI patients, the relationship between *NOBOX* gene mutation and POI was reported (Qin *et al.* 2007), showing down regulation of follicle growth factors such as *GDF9* and *KITL* (Bouilly *et al.* 2015). Furthermore, *FOXL2* gene mutation is also found in patients with POI (Crisponi *et al.* 2001).

Activation of dormant primordial follicles to primary follicles is regulated by the PI3K-AKT pathway in the oocyte (Reddy *et al.* 2008) (Figs 1 and 2). Kit ligand binds its cognate tyrosine kinase receptor C-KIT and stimulates PI3K, leading to the conversion of the lipid second messenger PIP2 (phosphatidylinositol (4,5) bisphosphate) into PIP3 (phosphatidylinositol (3,4,5) triphosphate). Then, PIP3 activates AKT and promotes translocation of AKT to the nucleus to inhibit the activity of a transcriptional factor, FOXO3. PTEN negatively regulates this pathway by dephosphorylating PIP3 to PIP2 (Fig. 2) (Reddy *et al.* 2005).

Studies using ovarian cultures showed that KITL treatment dramatically increased the number of primary follicles (Parrot & Skinner 1999, Jones & Pepling 2013, Zhang *et al.* 2014) (Figs 1 and 2). Oocyte-specific deletion of *Pten*, a negative regulator of PI3K, in mice (GDF9-Cre mice) showed that all primordial follicles developed simultaneously, and no follicle in the ovary remained at 16 weeks of age (Reddy *et al.* 2008, Jagarlamudi *et al.*

Figure 2 The PI3K-AKT pathway regulates primordial follicle activation. FOXO3 located in oocyte nucleus prevents the activation of dormant primordial follicles. The activation of upstream receptor tyrosine kinases (RTKs) by their cognate ligands (e.g. kit ligand (KITL), etc.) stimulates PI3K activity, resulting in increases in PIP3 levels followed by AKT stimulation. Then, activated AKT translocate to the nucleus and phosphorylate FOXO3 to promote nuclear export of FOXO3, leading to the suppression of FOXO3 activity to stimulate primordial follicle activation. The activated Akt also suppresses TSC1 and TSC2 complex to increase active GTP-bound form of Ras homolog enriched in brain (Rheb) and then activates mTOR, leading to activation of dormant follicles. PTEN converts PIP3 to PIP2 thus suppressing the actions of PI3K. Deletion of the *PTEN* and *FOXO3* genes exhibit activation of primordial follicles.

2009). Furthermore, deletion of the transcription factor FOXO3A, downstream of PI3K-Akt signaling, in mice showed the same phenotype as oocyte-specific *Pten* null mice (Castrillon *et al.* 2003). Therefore, activated Foxo3a is essential for the maintenance of the oocyte pool. Also, the action of PI3K-AKT signaling induces follicle development from the primordial stage (Brenkman *et al.* 2003) (Fig. 2). Mammalian target of rapamycin (mTOR) is the catalytic subunit of two structurally distinct complexes, mTORC1 and mTORC2. *Tsc1* or *Tsc2* null mice also showed spontaneous activation of dormant primordial follicles to primary follicles (Adhikari *et al.* 2009, 2010). Interestingly, double deletion of *Tsc1* and *Pten* leads to synergistically enhanced primordial follicle activation, demonstrating the essential and cooperative roles of PI3K-AKT and mTOR signaling pathways in regulating primordial follicle dormancy (Adhikari & Liu 2010) (Fig. 2).

After development to the primary stage of follicles, oocyte secreted GDF9 that is essential for the follicle growth (Eppig 2001) (Fig. 1). *Gdf9* null mice showed defects in growth after the secondary follicle stage (Dong *et al.* 1996). Moreover, sheep with mutation of the related *BMP15* gene also showed defective follicle development after the primary stage (Galloway *et al.* 2000). Because *Bmp15* null mice are fertile (Yan *et al.* 2001), these reports indicate divergent gene functions between species. After secondary follicle stage, follicle growth and ovulation are mainly regulated by the pituitary hormone follicle stimulating hormone (FSH) and luteinizing hormone (LH) (Richard *et al.* 2002).

In human, *GDF9* and *BMP15* are expressed in the oocyte of primordial follicles. From the analyses of 127 cases of POI patients, the relationship between *GDF9* gene mutation and POI has been reported (Dixit *et al.* 2005). Furthermore, two sisters with the *BMP15* mutation are associated with the POI phenotype (Di Pasquale *et al.* 2004). *GDF9* and *BMP15* mutations are also associated with deficiency in early follicle growth leading to their infertility (Di Pasquale *et al.* 2004, Chand *et al.* 2006). Also, FSH receptor (*FSHR*) and *INHA* (Inhibin a), mutations are associated with the POI phenotype in woman. From the analyses of 7 families of POI patients, mutation of A C566T transition in exon7 of the *FSHR* gene showed reduction of ligand binding capacity and signal transduction in receptor-transfected cells (Aittomaki *et al.* 1995). Likewise, *INHA* mutation in the *INHA* sequence occurs in ~7% of POI patients (Shelling *et al.* 2000).

Development of new therapy for POI based on PI3K-Akt signaling

Primordial follicle number decreases during aging. In adult female, approximately 1000 primordial follicles are activated every month. If the number of primordial follicle in the ovary is low (less than 1000), activation

of primordial follicle is defective (Macklon *et al.* 1999), resulting in anovulation. POI shows menopause before 40 years of age with abnormally low residual follicles. Hence, it is extremely difficult to induce follicle growth in POI patients even under ovarian hyper-stimulation (Nelson 2009), and egg donation is the only established method for infertility treatment. Recently, we developed a novel method which enables POI patients to conceive their own genetic children by the activation of residual dormant follicles focusing on the PI3K-AKT-FOXO3 signaling pathway. Because of difficulty in performing gene therapy in human, we activated dormant follicles using a PTEN inhibitor and a PI3K activator based on a short-term (2 days) *in vitro* activation protocol. Previous studies showed that a PTEN inhibitor, bpV, indirectly activates the PI3K-AKT signaling pathway (Posner *et al.* 1994, Bavan *et al.* 1995, Schmid *et al.* 2004), whereas a cell-permeable phosphopeptide, 740Y-P, binds to the SH2 domain of p85 regulatory subunit of PI3K to activate its catalytic subunit (Derossi *et al.* 1998). Using both drugs, we demonstrated that short-term culture of neonatal mouse and human ovaries increased primary follicle numbers via nuclear exclusion of FOXO3 *in vitro* (Li *et al.* 2010). Also, we generated mature oocytes displaying normal epigenetic modification in imprinted genes after the transplantation of activated ovaries into the kidney capsule (Li *et al.* 2001). Based on these successes of basic findings, we named this approach as IVA (*in vitro* activation) and applied it for clinical studies to treat infertility in patients with POI. We reported two deliveries of healthy babies after IVA (Kawamura *et al.* 2012, Suzuki *et al.* 2015) (Fig. 3). After our publication of IVA, two separate centers have repeated the IVA approach and one successful delivery was reported (Zhai *et al.* 2016). Currently, the IVA treatment is expanding to more infertility centers.

Limitation of IVA treatment and future studies: the prospect of oocyte regeneration

Ovaries from around half of the POI patients contain no residual follicles under histological analyses (Kawamura *et al.* 2012, Zhai *et al.* 2016). Although we grafted the activated ovarian tissues after IVA back into these patients, no follicle growth was found (Kawamura *et al.* 2012, Suzuki *et al.* 2015). Oocyte regeneration is a potential method to treat these patients so that they could conceive their own genetic children. Recently, a group reported the success of oocyte generation from murine ES/iPS cells (Hayashi *et al.* 2012, Hikabe *et al.* 2016). Researchers induced primordial germ cell-like cells (PGCLCs) from murine ES/iPS cells and then co-cultured PGCLCs with murine fetal ovarian somatic cells to form re-aggregated ovaries. In the aggregated ovaries, PGCLCs resumed meiosis and formed follicles, leading to the production of competent oocytes for fertilization and development into healthy

Figure 3 Potential infertility treatments for POI patients to conceive their own genetic children. The IVA approach is effective to treat patients with POI who have residual follicles. However, IVA cannot apply to the POI patients without residual follicles. Oocyte donation is the only infertility therapy to treat POI patients with no residual follicles. To establish the method to conceive their own genetic children in such patients, investigators attempt to regenerate mature oocytes from iPS or oocyte stem cells.

pups (Hayashi *et al.* 2012, Hayashi & Saito 2013, Hikabe *et al.* 2016). Although this method is attractive to regenerate oocytes from POI patients using iPS cells, there are several obstacles to overcome before applying this technique to humans. In particular, it is difficult to obtain ovarian somatic cells from human fetal ovaries. Furthermore, the efficacy of the differentiation of iPS cells into mature oocytes is extremely low under published procedures (Fig. 3).

To address these issues, researchers are trying to regenerate oocytes from oocyte stem cells presumably present in ovaries. A group showed regeneration of mature oocytes in mouse and human from oocyte stem cells isolated from ovarian cortex using DDX4 antibodies through fluorescence activated cell sorting (Johnson *et al.* 2004, 2005, White *et al.* 2012) (Fig. 3). However, many labs could not confirm the presence of oocyte stem cells in mouse (Eggen *et al.* 2006, Lei & Spradling 2013) and human (Byskov *et al.* 2011). Moreover, recent work showed uncertainty in the specificity of antibodies used for the isolation of murine and human oocyte stem cells (Albertini & Gleicher 2015), and the inability of follicle formation after grafting these oocyte stem cells into mouse postnatal ovaries (Zhang *et al.* 2015). Therefore, more studies will be required to establish the method for regeneration of oocytes from oocyte stem cells.

Conclusion

Most of the knowledge accumulated on germ cell development is based on murine studies. In contrast,

knowledge on human germ cell development is limited. Although it is difficult to study human germ line cells due to ethical issues, more findings on human germ cell development could help us understand pathophysiology of POI. It could also contribute to the improvement of IVA treatment for POI patients as well as the development of new methods for oocyte regeneration from POI patients.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was supported by Grant-In-Aid for Scientific Research (16H05476, 16K11116, 15K15613 and 25462579) and funds from Takeda Science Foundation, Naito Foundation, and Mochida Memorial Foundation (to K K) and Young Research Fellowship (12J6118) (to I K) from the Japan Society for the Promotion of Science.

Acknowledgements

The authors thank Dr Aaron J W Hsueh (Stanford University School of Medicine, Stanford, CA) for critical reading and editing of the manuscript.

References

- Adhikari D & Liu K 2010 mTOR signaling in the control of activation of primordial follicles *Cell Cycle* **9** 1673–1674. (doi:10.4161/cc.9.9.11626)
- Adhikari D, Flohr G, Gorre N, Shen Y, Yang H, Lundin E, Lan Z, Gambello MJ & Liu K 2009 Disruption of Tsc2 in oocytes leads to overactivation of the entire pool of primordial follicles. *Molecular Human Reproduction* **15** 765–770. (doi:10.1093/molehr/gap092)
- Adhikari D, Zheng W, Shen Y, Gorre N, Hämäläinen T, Cooney AJ, Huhtaniemi I, Lan ZJ & Liu K 2010 Tsc/mTORC1 signaling in oocytes governs the quiescence and activation of primordial follicles. *Human Molecular Genetics* **19** 397–410. (doi:10.1093/hmg/ddp483)
- Aittomäki K1, Lucena JL, Pakarinen P, Sistonen P, Tapanainen J, Gromoll J, Kaskikari R, Sankila EM, Lehtväslaiho H, Engel AR *et al.* 1995 Mutation in the follicle-stimulating hormone receptor gene causes hereditary hypergonadotropic ovarian failure. *Cell* **82** 959–968. (doi:10.1016/0092-8674(95)90275-9)
- Albertini DF & Gleicher N 2015 A detour in the quest for oogonial stem cells: methods matter. *Nature Medicine* **21** 1126–1127. (doi:10.1038/nm.3969)
- Barker TG 1963 A quantitative and cytological study of germ cells in human ovaries. *Proceedings of the Royal Society of London: Series B, Biological Sciences* **158** 417–433. (doi:10.1098/rspb.1963.0055)
- Bayne RA, Martins da Silva SJ & Anderson RA 2004 Increased expression of the FIGLA transcription factor is associated with primordial follicle formation in the human fetal ovary. *Molecular Human Reproduction* **10** 373–381. (doi:10.1093/molehr/gah056)
- Bevan AP, Burgess JW, Yale JF, Drake PG, Lachance D, Baquiran G, Shaver A & Posner BI 1995 In vivo insulin mimetic effects of pV compounds: role for tissue targeting in determining potency. *American Journal of Physiology* **268** E60–E66.
- Bouilly J, Roucher-Boulez F, Gompel A, Bry-Gauillard H, Azibi K, Beldjord C, Dodé C, Bouligand J, Mantel AG, Hécart AC *et al.* 2015 New NOBOX mutations identified in large cohort of women with primary

- ovarian insufficiency decrease KIT-L expression. *Journal of Clinical Endocrinology and Metabolism* **100** 994–1001. (doi:10.1210/jc.2014-2761)
- Brenkman AB & Burgering BM 2003 FoxO3a eggs on fertility and aging. *Trends in Molecular Medicine* **9** 464–467. (doi:10.1016/j.molmed.2003.09.003)
- Byskov AG, Høyer PE, Yding Andersen C, Kristensen SG, Jespersen A & Møllgård K 2011 No evidence for the presence of oogonia in the human ovary after their final clearance during the first two years of life. *Human Reproduction* **26** 2129–2139. (doi:10.1093/humrep/der145)
- Castrillon DH, Quade BJ, Wang TY, Quigley C & Crum CP 2000 The human VASA gene is specifically expressed in the germ cell lineage. *PNAS* **97** 9585–9590. (doi:10.1073/pnas.160274797)
- Castrillon DH, Miao L, Kollipara R, Horner JW & DePinho RA 2003 Suppression of ovarian follicle activation in mice by the transcription factor Foxo3a. *Science* **301** 215–218. (doi:10.1126/science.1086336)
- Chand AL, Ponnampalam AP, Harris SE, Winship IM & Shelling AN 2006 Mutational analysis of BMP15 and GDF9 as candidate genes for premature ovarian failure. *Fertility and Sterility* **86** 1009–1102. (doi:10.1016/j.fertnstert.2006.02.107)
- Choi Y, Yuan D & Rajkovic A 2008 Germ cell-specific transcriptional regulator sohlh2 is essential for early mouse folliculogenesis and oocyte-specific gene expression. *Biology of Reproduction* **79** 1176–1182. (doi:10.1095/biolreprod.108.071217)
- Coulam CB, Adamson SC & Annegers JF 1986 Incidence of premature ovarian failure. *Obstetrics and Gynecology* **67** 604–606.
- Crisponi L, Deiana M, Loi A, Chiappe F, Uda M, Amati P, Bisceglia L, Zelante L, Nagaraja R, Porcu S *et al.* 2001 The putative forkhead transcription factor FOXL2 is mutated in blepharophimosis/ptosis/epicanthus in versus syndrome. *Nature Genetics* **27** 159–166. (doi:10.1038/84781)
- Derossi D, Williams EJ, Green PJ, Dunican DJ & Doherty P 1998 Stimulation of mitogenesis by a cell-permeable PI 3-kinase binding peptide. *Biochemical and Biophysical Research Communications* **251** 148–152. (doi:10.1006/bbrc.1998.9444)
- Di Pasquale E, Beck-Peccoz P & Persani L 2004 Hypergonadotropic ovarian failure associated with an inherited mutation of human bone morphogenetic protein-15 (BMP15) gene. *American Journal of Human Genetics* **75** 106–111. (doi:10.1086/422103)
- Dixit H, Rao LK, Padmalatha V, Kanakavalli M, Deenadayal M, Gupta N, Chakravarty B & Singh L 2005 Mutational screening of the coding region of growth differentiation factor 9 gene in Indian women with ovarian failure. *Menopause* **12** 749–754. (doi:10.1097/01.gme.0000184424.96437.7a)
- Dolci S, Pesce M & De Felici M 1993 Combined action of stem cell factor, leukemia inhibitory factor, and cAMP on in vitro proliferation of mouse primordial germ cells. *Molecular Reproduction and Development* **35** 134–139. (doi:10.1002/mrd.1080350206)
- Dong J, Albertini DF, Nishimori K, Kumar TR, Lu N & Matzuk MM 1996 Growth differentiation factor-9 is required during early ovarian folliculogenesis. *Nature* **383** 531–535. (doi:10.1038/383531a0)
- Eggen K, Jurga S, Gosden R, Min IM & Wagers AJ 2006 Ovulated oocytes in adult mice derive from non-circulating germ cells. *Nature* **441** 1109–1114. (doi:10.1038/nature04929)
- Eppig JJ 2001 Oocyte control of ovarian follicular development and function in mammals. *Reproduction* **122** 829–838. (doi:10.1530/rep.0.1220829)
- Franchi LL & Mandl AM 1962 The ultrastructure of oogonia and oocyte in the foetal and neonatal rat. *Proceedings of the Royal Society of London* **157** 99–129. (doi:10.1098/rspb.1962.0064)
- Galloway SM, McNatty KP, Cambridge LM, Laitinen MP, Juengel JL, Jokiranta TS, McLaren RJ, Luiro K, Dodds KG, Montgomery GW *et al.* 2000 Mutations in an oocyte-derived growth factor gene (BMP15) cause increased ovulation rate and infertility in a dosage-sensitive manner. *Nature Genetics* **25** 279–283. (doi:10.1038/77033)
- Ginsburg M, Snow MH & McLaren A 1990 Primordial germ cells in the mouse embryo during gastrulation. *Development* **110** 521–528.
- Gondos B, Westergaard L & Byskov AG 1986 Initiation of oogenesis in the human fetal ovary: ultrastructural and squash preparation study. *American Journal of Obstetrics and Gynecology* **155** 189–195. (doi:10.1016/0002-9378(86)90109-2)
- Greenbaum MP, Iwamori N, Agno JE & Matzuk MM 2009 Mouse TEX14 is required for embryonic germ cell intercellular bridges but not female fertility. *Biology of Reproduction* **80** 449–457. (doi:10.1095/biolreprod.108.070649)
- Grive KJ & Freiman RN 2015 The developmental origins of the mammalian ovarian reserve. *Development* **142** 2554–2563 (doi:10.1242/dev.125211)
- Hayashi K & Saitou M 2013 Generation of eggs from mouse embryonic stem cells and induced pluripotent stem cells. *Nature Protocols* **8** 1513–1524. (doi:10.1038/nprot.2013.090)
- Hayashi K, Ogushi S, Kurimoto K, Shimamoto S, Ohta H & Saitou M 2012 Offspring from oocytes derived from in vitro primordial germ cell-like cells in mice. *Science* **338** 971–975. (doi:10.1126/science.1226889)
- Heeren AM, van Iperen L, Klootwijk DB, de Melo Bernardo A, Roost MS, Gomes Fernandes MM, Louwe LA, Hilders CG, Helmerhorst FM, van der Westerlaken LA *et al.* 2015 Development of the follicular basement membrane during human gametogenesis and early folliculogenesis. *BMC Developmental Biology* **15** 4. (doi:10.1186/s12861-015-0054-0)
- Hikabe O, Hamazaki N, Nagamatsu G, Obata Y, Hirao Y, Hamada N, Shimamoto S, Imamura T, Nakashima K, Saitou M *et al.* 2016 Reconstitution in vitro of the entire cycle of the mouse female germ line. *Nature* **539** 299–303. (doi:10.1038/nature20104)
- Houmar B, Small C, Yang L, Nalua-Cecchini T, Cheng E, Hassold T & Griswold M 2009 Global gene expression in the human fetal testis and ovary. *Biology of Reproduction* **81** 438–443. (doi:10.1095/biolreprod.108.075747)
- Jagarlamudi K, Liu L, Adhikari D, Reddy P, Idahl A, Ottander U, Lundin E & Liu K 2009 Oocyte-specific deletion of Pten in mice reveals a stage-specific function of PTEN/PI3K signaling in oocytes in controlling follicular activation. *PLoS ONE* **4** e6186. (doi:10.1371/journal.pone.0006186)
- James RS, Coppin B, Dalton P, Dennis NR, James RS, Coppin B, Dalton P, Dennis NR, Mitchell C, Sharp AJ *et al.* 1998 A study of females with deletions of the short arm of the X chromosome. *Human Genetics* **102** 507–516.
- Jameson SA, Natarajan A, Cool J, DeFalco T, Maatouk DM, Mork L, Munger SC & Capel B 2012 Temporal transcriptional profiling of somatic and germ cells reveals biased lineage priming of sexual fate in the fetal mouse gonad. *PLoS Genetics* **8** e1002575. (doi:10.1371/journal.pgen.1002575)
- Johnson J, Canning J, Kaneko T, Pru JK & Tilly JL 2004 Germline stem cells and follicular renewal in the postnatal mammalian ovary. *Nature* **428** 145–150. (doi:10.1038/nature02316)
- Johnson J, Bagley J, Skaznik-Wikiel M, Lee HJ, Adams GB, Niikura Y, Tschudy KS, Tilly JC, Cortes ML, Forkert R *et al.* 2005 Oocyte generation in adult mammalian ovaries by putative germ cells in bone marrow and peripheral blood. *Cell* **122** 303–315. (doi:10.1016/j.cell.2005.06.031)
- Jones RL & Pepling ME 2013 KIT signaling regulates primordial follicle formation in the neonatal mouse ovary. *Developmental Biology* **382** 186–197. (doi:10.1016/j.ydbio.2013.06.030)
- Kawamura K, Cheng Y, Suzuki N, Deguchi M, Sato Y, Takae S, Ho CH, Kawamura N, Tamura M, Hashimoto S *et al.* 2013 Hippo signaling disruption and Akt stimulation of ovarian follicles for infertility treatment. *PNAS* **110** 17474–17479. (doi:10.1073/pnas.1312830111)
- Kawase E, Yamamoto H, Hashimoto K & Nakatsuji N 1994 Tumor necrosis factor-alpha (TNF-alpha) stimulates proliferation of mouse primordial germ cells in culture. *Developmental Biology* **161** 91–95. (doi:10.1006/dbio.1994.1011)
- Lei L & Spradling AC 2013 Female mice lack adult germ-line stem cells but sustain oogenesis using stable primordial follicles. *PNAS* **110** 8585–8590. (doi:10.1073/pnas.1306189110)
- Lei L & Spradling AC 2016 Mouse oocytes differentiate through organelle enrichment from sister cyst germ cells. *Science* **352** 95–99. (doi:10.1126/science.aad2156)
- Li J, Kawamura K, Cheng Y, Liu S, Klein C, Liu S, Duan EK & Hsueh AJ 2010 Activation of dormant ovarian follicles to generate mature eggs. *PNAS* **107** 10280–10284. (doi:10.1073/pnas.1001198107)
- Liang L, Soyak SM & Dean J 1997 FIGalpha, a germ cell specific transcription factor involved in the coordinate expression of the zona pellucida genes. *Development* **124** 4939–4947.
- MacGregor GR, Zambrowicz BP & Soriano P 1995 Tissue non-specific alkaline phosphatase is expressed in both embryonic and extraembryonic lineages during mouse embryogenesis but is not required for migration of primordial germ cells. *Development* **121** 1487–1496.

- Macklon NS & Fauser BC 1999 Aspects of ovarian follicle development throughout life. *Hormone Research* **52** 161–170. (Review) (doi:10.1159/000023456)
- Martins da Silva SJ, Bayne RA, Cambray N, Hartley PS, McNeilly AS & Anderson RA 2004 Expression of activin subunits and receptors in the developing human ovary: activin A promotes germ cell survival and proliferation before primordial follicle formation. *Developmental Biology* **26** 334–345. (doi:10.1016/j.ydbio.2003.10.030)
- Matsui Y, Zsebo K & Hogan BL 1992 Derivation of pluripotential embryonic stem cells from murine primordial germ cells in culture. *Cell* **70** 841–847. (doi:10.1016/0092-8674(92)90317-6)
- Menke DB, Koubova J & Page DC 2003 Sexual differentiation of germ cells in XX mouse gonads occurs in an anterior-to-posterior wave. *Developmental Biology* **262** 303–312. (doi:10.1016/S0012-1606(03)00391-9)
- Modi DN, Sane S & Bhartiya D 2003 Accelerated germ cell apoptosis in sex chromosome aneuploid fetal human gonads. *Molecular Human Reproduction* **9** 219–225. (doi:10.1093/molehr/gag031)
- Motta PM, Makabe S & Nottola SA 1997 The ultrastructure of human reproduction. I. The natural history of the female germ cell: origin, migration and differentiation inside the developing ovary. *Human Reproduction Update* **3** 281–295. (Review) (doi:10.1093/humupd/3.3.281)
- Nelson LM 2009 Clinical practice. Primary ovarian insufficiency. *New England Journal of Medicine* **360** 606–614. (doi:10.1056/NEJMc0808697)
- Noce T, Okamoto-Ito S & Tsunekawa N 2001 Vasa homolog genes in mammalian germ cell development. *Cell Structure and Function* **26** 131–136. (doi:10.1247/csf.26.131)
- Ogata T & Matsuo N 1995 Turner syndrome and female sex chromosome aberrations: deduction of the principal factors involved in the development of clinical features. *Human Genetics* **95** 607–629. (doi:10.1007/bf00209476)
- Oulad-Abdelghani M, Bouillet P, Décimo D, Gansmuller A, Heyberger S, Dollé P, Bronner S, Lutz Y & Chambon P 1996 Characterization of a premeiotic germ cell-specific cytoplasmic protein encoded by Stra8, a novel retinoic acid-responsive gene. *Journal of Cell Biology* **135** 469–477. (doi:10.1083/jcb.135.2.469)
- Pangas SA, Li X, Robertson EJ & Matzuk MM 2006 Premature luteinization and cumulus cell defects in ovarian-specific Smad4 knockout mice. *Molecular Endocrinology* **20** 1406–1422. (doi:10.1210/me.2005-0462)
- Paredes A, Garcia-Rudaz C, Kerr B, Tapia V, Dissen GA, Costa ME, Cornea A, Ojeda SR 2005 Loss of synaptonemal complex protein-1, a synaptonemal complex protein, contributes to the initiation of follicular assembly in the developing rat ovary. *Endocrinology* **146** 5267–5277. (doi:10.1210/en.2005-0965)
- Parrott JA & Skinner MK 1997 Direct actions of kit-ligand on theca cell growth and differentiation during follicle development. *Endocrinology* **138** 3819–3827. (doi:10.1210/en.138.9.3819)
- Pesce M, Canipari R, Ferri GL, Siracusa G & De Felici M 1996 Pituitary adenylate cyclase-activating polypeptide (PACAP) stimulates adenylate cyclase and promotes proliferation of mouse primordial germ cells. *Development* **122** 215–221.
- Pesce M, Klinger FG & De Felici M 2002 Derivation in culture of primordial germ cells from cells of the mouse epiblast: phenotypic induction and growth control by Bmp4 signalling. *Mechanisms of Development* **112** 15–24. (doi:10.1016/S0925-4773(01)00624-4)
- Pietrobono R, Tabolacci E, Zalfa F, Zito I, Terracciano A, Moscato U, Bagni C, Oostra B, Chiurazzi P & Neri G 2005 Molecular dissection of the events leading to inactivation of the FMR1 gene. *Human Molecular Genetics* **15** 267–277. (doi:10.1093/hmg/ddi024)
- Pittman DL, Cobb J, Schimenti KJ, Wilson LA, Cooper DM, Brignull E, Handel MA & Schimenti JC 1998 Meiotic prophase arrest with failure of chromosome synapsis in mice deficient for Dmcl1, a germline-specific RecA homolog. *Molecular Cell* **1** 697–705. (doi:10.1016/S1097-2765(00)80069-6)
- Posner BI, Faure R, Burgess JW, Bevan AP, Lachance D, Zhang-Sun G, Fantus IG, Ng JB, Hall DA, Lum BS *et al.* 1994 Peroxovanadium compounds. A new class of potent phosphotyrosine phosphatase inhibitors which are insulin mimetics. *Journal of Biological Chemistry* **269** 4596–4604.
- Puglisi R, Montanari M, Chiarella P, Stefanini M & Boitani C 2004 Regulatory role of BMP2 and BMP7 in spermatogonia and Sertoli cell proliferation in the immature mouse. *European Journal of Endocrinology* **151** 511–520. (doi:10.1530/eje.0.1510511)
- Qin Y, Choi Y, Zhao H, Simpson JL, Chen ZJ & Rajkovic A 2007 NOBOX homeobox mutation causes premature ovarian failure. *American Journal of Human Genetics* **81** 576–581. (doi:10.1086/519496)
- Rajkovic A, Pangas SA, Ballow D, Suzumori N & Matzuk MM 2004 NOBOX deficiency disrupts early folliculogenesis and oocyte-specific gene expression. *Science* **305** 1157–1159. (doi:10.1126/science.1099755)
- Reddy P, Shen L, Ren C, Boman K, Lundin E, Ottander U, Lindgren P, Liu YX, Sun QY & Liu K 2005 Activation of Akt (PKB) and suppression of FKHL1 in mouse and rat oocytes by stem cell factor during follicular activation and development. *Developmental Biology* **281** 160–170. (doi:10.1016/j.ydbio.2005.02.013)
- Reddy P, Liu L, Adhikari D, Jagarlamudi K, Rajareddy S, Shen Y, Du C, Tang W, Hämäläinen T, Peng SL *et al.* 2008 Oocyte-specific deletion of Pten causes premature activation of the primordial follicle pool. *Science* **319** 611–613. (doi:10.1126/science.1152257)
- Reynolds N, Collier B, Bingham V, Gray NK & Cooke HJ 2007 Translation of the synaptonemal complex component Sycp3 is enhanced in vivo by the germ cell specific regulator Dazl. *RNA* **13** 974–981. (doi:10.1261/rna.465507)
- Richards JS & Pangas SA 2010 The ovary: basic biology and clinical implications. *Journal of Clinical Investigation* **120** 963–972. (doi:10.1172/JCI41350)
- Richards JS, Russell DL, Ochsner S, Hsieh M, Doyle KH, Falender AE, Lo YK & Sharma SC 2002 Novel signaling pathways that control ovarian follicular development, ovulation and luteinization. *Recent Progress in Hormone Research* **57** 195–220. (doi:10.1210/rp.57.1.195)
- Rodrigues P, Limback D, McGinnis LK, Plancha CE & Albertini DF 2009 Multiple mechanisms of germ cell loss in the perinatal mouse ovary. *Reproduction* **137** 709–720. (doi:10.1530/REP-08-0203)
- Ross A, Munger S & Capel B 2007 Bmp7 regulates germ cell proliferation in mouse fetal gonads. *Sexual Development* **1** 127–137. (doi:10.1159/000100034)
- Schmucker B, Ballhausen WG & Pfeiffer RA 1996 Mosaicism of a microdeletion of 486bp involving the CGG repeat of the FMR1 gene due to misalignment of GTT tandem repeats at chi-like elements flanking both breakpoints and a full mutation. *Human Genetics* **98** 409–414. (doi:10.1007/s004390050230)
- Schmid AC, Byrne RD, Vilar R & Woscholski R 2004 Bisperoxovanadium compounds are potent PTEN inhibitors. *FEBS Letters* **566** 35–38. (doi:10.1016/j.febslet.2004.03.102)
- Shelling AN, Burton KA, Chand AL, van Ee CC, France JT, Farquhar CM & Milsom SR 2000 Inhibin: a candidate gene for premature ovarian failure. *Human Reproduction* **15** 2644–2649. (doi:10.1093/humrep/15.12.2644)
- Simpson JL & Rajkovic A 1999 Ovarian differentiation and gonadal failure. *American Journal of Medical Genetics* **89** 186–200. (doi:10.1002/(SICI)1096-8628(19991229)89:4<186::AID-AJMG3>3.0.CO;2-5)
- Soyal SM, Amleh A & Dean J 2000 FIGalpha, a germ cell-specific transcription factor required for ovarian follicle formation. *Development* **127** 4645.
- Stoop H, Honecker F, Cools M, de Krijger R, Bokemeyer C & Looijenga LH 2005 Differentiation and development of human female germ cells during prenatal gonadogenesis: an immunohistochemical study. *Human Reproduction* **20** 1466–1476. (doi:10.1093/humrep/deh800)
- Strauss JF III & Williams CJ 2004 The ovarian life cycle. In *Book Reproductive Endocrinology* 213–225.
- Sullivan AK, Marcus M, Epstein MP, Allen EG, Anido AE, Paquin JJ, Yadav-Shah M & Sherman SL 2005 Association of FMR1 repeat size with ovarian dysfunction. *Human Reproduction* **20** 402–412. (doi:10.1093/humrep/deh635)
- Suzuki N, Yoshioka N, Takae S, Sugishita Y, Tamura M, Hashimoto S, Morimoto Y & Kawamura K 2015 Successful fertility preservation following ovarian tissue vitrification in patients with primary ovarian insufficiency. *Human Reproduction* **30** 608–615. (doi:10.1093/humrep/deu353)
- Tam PPL & Snow MHL 1981 Proliferation and migration of primordial germ cells during compensatory growth in mouse embryos. *Journal of Embryology and Experimental Morphology* **64** 133–147.

- Tedesco M, Desimio MG, Klinger FG, De Felici M & Farini D** 2013 Minimal concentrations of retinoic acid induce stimulation by retinoic acid 8 and promote entry into meiosis in isolated pregonadal and gonadal mouse primordial germ cells. *Biology of Reproduction* **88** 145. (doi:10.1095/biolreprod.112.106526)
- Terracciano A, Chiurazzi P & Neri G** 2005 Fragile X syndrome. *American Journal of Medical Genetics Part C: Seminars in Medical Genetics* **15** 32–37. (doi:10.1002/ajmg.c.30062)
- Tingen C, Kim A & Woodruff TK** 2009 The primordial pool of follicles and nest breakdown in mammalian ovaries. *Molecular Human Reproduction* **15** 795–803. (doi:10.1093/molehr/gap073)
- Tosh D, Rani HS, Murty US, Deenadayal A & Grover P** 2015 Mutational analysis of the FIGLA gene in women with idiopathic premature ovarian failure. *Menopause* **22** 520–526. (doi:10.1097/GME.0000000000000340)
- Tung JY, Rosen MP, Nelson LM, Turek PJ, Witte JS, Cramer DW, Cedars MI & Reijo-Pera RA** 2006a Novel missense mutations of the Deleted-in-AZOospermia-Like (DAZL) gene in infertile women and men. *Reproductive Biology and Endocrinology* **2** 40. (doi:10.1186/1477-7827-4-40)
- Tung JY, Rosen MP, Nelson LM, Turek PJ, Witte JS, Cramer DW, Cedars MI & Pera RA** 2006b Variants in Deleted in AZOospermia-Like (DAZL) are correlated with reproductive parameters in men and women. *Human Genetics* **118** 730–740. (doi:10.1007/s00439-005-0098-5)
- Wang Z, Niu W, Wang Y, Teng Z, Wen J, Xia G & Wang C** 2015 Follistatin288 regulates germ cell cyst breakdown and primordial follicle assembly in the mouse ovary. *PLoS ONE* **15** e0129643. (doi:10.1371/journal.pone.0129643)
- Weakley BS** 1967 Light and electron microscopy of developing germ cell and follicle cells in the ovary of the golden hamster: twenty-four hours before birth to eight days post partum. *Journal of Anatomy* **101** 435–459.
- White YA, Woods DC, Takai Y, Ishihara O, Seki H & Tilly JL** 2012 Oocyte formation by mitotically active germ cells purified from ovaries of reproductive-age women. *Nature Medicine* **18** 413–421. (doi:10.1038/nm.2669)
- Witschi E** 1948 Migration of germ cells of human embryos from the yolk sac to the primitive gonadal folds. *Carnegie Institution of Washington Publication* **32** 67–80.
- Xu J & Gridley T** 2013 Notch2 is required in somatic cells for breakdown of ovarian germ-cell nests and formation of primordial follicles. *BMC Biology* **13** 13. (doi:10.1186/1741-7007-11-1)
- Yan C, Wang P, DeMayo J, DeMayo FJ, Elvin JA, Carino C, Prasad SV, Skinner SS, Dunbar BS, Dube JL et al.** 2001 Synergistic roles of bone morphogenetic protein 15 and growth differentiation factor 9 in ovarian function. *Molecular Endocrinology* **15** 854–866. (doi:10.1210/mend.15.6.0662)
- Yao HH, Matzuk MM, Jorgez CJ, Menke DB, Page DC, Swain A & Capel B** 2004 Follistatin operates downstream of Wnt4 in mammalian ovary organogenesis. *Developmental Dynamics* **230** 210–215. (doi:10.1002/dvdy.20042)
- Zhai J, Yao G, Dong F, Bu Z, Cheng Y, Sato Y, Hu L, Zhang Y, Wang J, Dai S et al.** 2016 In vitro activation of follicles and fresh tissue auto-transplantation in primary ovarian insufficiency patients. *Journal of Clinical Endocrinology and Metabolism* **101** 4405–4412. (doi:10.1210/jc.2016-1589)
- Zhang H, Risal S, Gorre N, Busayavalasa K, Li X, Shen Y, Bosbach B, Brännström M & Liu K** 2014 Somatic cells initiate primordial follicle activation and govern the development of dorm antoocytes in mice. *Current Biology* **24** 2501–2508. (doi:10.1016/j.cub.2014.09.023)
- Zhang H, Panula S, Petropoulos S, Edsgård D, Busayavalasa K, Liu L, Li X, Risal S, Shen Y, Shao J et al.** 2015 Adult human and mouse ovaries lack DDX4-expressing functional oogonial stem cells. *Nature Medicine* **21** 1116–1118. (doi:10.1038/nm.3775)
- Zhao H, Chen ZJ, Qin Y, Shi Y, Wang S, Choi Y, Simpson JL & Rajkovic A** 2008 Transcription factor FIGLA is mutated in patients with premature ovarian failure. *American Journal of Human Genetics* **82** 1342–1348. (doi:10.1016/j.ajhg.2008.04.018)

Received 8 January 2017

First decision 30 January 2017

Revised manuscript received 10 February 2017

Accepted 13 March 2017