Methylation dynamics during folliculogenesis and early embryo development in sheep

Laura Masala, Giovanni Pietro Burras, Emanuela Bellu, Federica Ariu, Luisa Bogliolo, Sergio Ledda and Daniela Bebbere

Department of Veterinary Medicine, University of Sassari, Sassari, Italy

Correspondence should be addressed to D Bebbere; Email: dbebbere@uniss.it

The journal and the authors regret an error in this article published in the May (vol 153, issue 5 pp 605–619) issue of the journal. Figures 1 and 2 were erroneously repeated although the legends were correct. Figure 2 and its legend is correct as published and the correct Figure 1 is published below with the correct legend. The journal apologises for this error.

Figure 1 Relative expression of *DNMT1*, *DNMT3A*, *DNMT3B*, *TET1*, *TET2*, *TET3*, *MBD1*, *MBD3* and *MBD4* in ovine growing oocytes (GO) at different diameter derived from pre-pubertal animals: 70–90 (small; S GO), 90–110 (medium; M GO) or 110–130 μ m (large; L GO) diameter. The relative quantification of all transcripts was performed after normalization against luciferase mRNA levels and the number of oocytes and embryos (Su *et al.* 2007, Ohsugi *et al.* 2008, Evsikov *et al.* 2009). Relative abundance values are expressed as Δ Cq (Y axis) and show the mean value $\pm$ S.E.M. of five replicates for each stage (each replicate=pool of 30 oocytes). Different letters indicate a significant difference in relative mRNA abundance (ANOVA $P < 0.05$) among the groups.