A single dose of allopregnanolone affects rat ovarian morphology and steroidogenesis

Laura Tatiana Pelegrina¹, Antonella Rosario Ramona Cáceres², Fernando Alfredo Giuliani¹, Joana Antonella Asensio¹, Fernanda Parborell³ and Myriam Raquel Laconi¹

¹Laboratorio de Fisiopatología ovárica y Neurobiología, Instituto de Medicina y Biología Experimental de Cuyo (IMBECU-CONICET), Inbiomed-UM, Mendoza, Argentina, ²Laboratorio de Fisiopatología ovárica y Neurobiología, Instituto de Medicina y Biología Experimental de Cuyo (IMBECU-CONICET), Inbiomed-UM, Universidad Juan Agustín Maza, Mendoza, Argentina, and ³Laboratorio de Fisiopatología del ovario. Instituto de Biología y Medicina Experimental (IByME-CONICET), Buenos Aires, Argentina

The authors regret an error in Figure 3 of the above titled article published in the January 2017 (vol 153 pp 75–83) issue. The value for the enzymatic activity of 3 β HSD in medial basal hypothalamus (MBH; Fig. 3A left panel) of the control has been represented as 55 mU. The correct value should have been represented as 0.5 mU. The corrected figure panel (figure 3A) is published below.

Figure 3 Spectrophotometric analysis of ALLO effect over 3 β -HSD (A and B), 3 α -HSOR (C and D) and 20 α -HSD (E and F) enzymatic activities in the medial basal hypothalamus (MBH left panel) and in the ovary (right panel) of estrous rats. Bars represent the mean $\pm$ S.E.M. ($n=6$; * $P<0.05$, ** $P<0.01$ and *** $P<0.001$).