

Steroidogenesis in Leydig cells: effects of aging and environmental factors

Yiyan Wang^{1,2}, Fenfen Chen¹, Leping Ye¹, Barry Zirkin² and Haolin Chen^{1,2}

¹The Second Affiliated Hospital and Yuying Children's Hospital of Wenzhou Medical University, Wenzhou, Zhejiang, China, ²Department of Biochemistry and Molecular Biology, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

Correspondence should be addressed to H Chen; Email: hchen13@jhu.edu

Abstract

Serum testosterone (TS) levels decrease with aging in both humans and rodents. Using the rat as a model system, it was found that age-related reductions in serum TS were not due to loss of Leydig cells, but rather to the reduced ability of the Leydig cells to produce TS in response to luteinizing hormone (LH). Detailed analyses of the steroidogenic pathway have suggested that two defects along the pathway, LH-stimulated cAMP production and cholesterol transport to and into the mitochondria, are of particular importance in age-related reductions in TS production. Although the mechanisms involved in these defects are far from certain, increasing oxidative stress appears to play a particularly important role. Interestingly, increased oxidative stress also appears to be involved in the suppressive effects of endocrine disruptors on Leydig cell TS production.

Reproduction (2017) **154** R111–R122

Introduction

Steroid hormones regulate critical phases of development and are essential for homeostasis of key physiological functions. In the male, the Leydig cells of the testis are responsible for producing testosterone (TS), and TS is essential for spermatogenesis. TS also is important for the maintenance of secondary sexual functions. A number of longitudinal studies have shown that in most men there is a slow decline in serum levels of total and bioactive TS with aging, beginning at about age 30 years, even in the absence of disease (Harman *et al.* 2001). Decline of serum TS usually is accompanied by increased or unchanging levels of LH and increased serum levels of FSH (Zwart *et al.* 1996), suggesting that the relative unresponsiveness of the Leydig cells to LH, rather than, or in addition to, defects in the hypothalamic–pituitary axis, plays a primary role in age-related reductions in serum TS (Mulligan *et al.* 2001, Keenan *et al.* 2006). This is discussed further below. TS deficiency in the adult has consequences for the general health of individuals, including increased body fat and fatigue, and decreased muscle mass, bone density, cognitive function and immune response (Huhtaniemi 2014).

In Brown Norway rats, as in aging men, serum TS levels were shown to decrease with age (Zirkin *et al.* 1993). However, unlike the many rat strains in which age-related decreases in serum TS levels are a consequence of reduced LH levels, LH levels do not decrease in Brown Norway rats. This suggests that in this strain, as in

men, there is primary hypogonadism, with age-related changes in Leydig cell steroidogenesis occurring at the gonadal level rather than secondary to hypothalamic–pituitary changes.

TS production can be affected not only by aging but also by endocrine disruptors. There now is evidence suggesting that there may be common mechanisms that explain both age-related and endocrine disruptor-induced reduced TS production. This review will emphasize research on Leydig cell steroidogenesis, focusing on the effects of aging on Leydig cell TS production. We also will discuss mechanisms in common with aging by which some endocrine disruptors also affect steroidogenesis.

Testosterone synthesis in Leydig cells

Steroidogenesis is a multi-step process that converts cholesterol into final steroid hormone products. As reviewed in Aghazadeh and coworkers (Aghazadeh *et al.* 2015), steroidogenesis consists of cholesterol mobilization from lipid droplets and/or the plasma membrane, cholesterol transport into mitochondria, pregnenolone formation in the mitochondria and subsequent conversion of pregnenolone into the final steroid products by enzymes of the smooth endoplasmic reticulum (Fig. 1). In the adult testis, Leydig cell TS production depends upon the pulsatile secretion of LH by the pituitary gland into the peripheral circulation. LH plays two essential roles in Leydig cell steroidogenesis:




Figure 1 Leydig cell steroidogenesis. LH binds its receptor on the Leydig cell membrane. LH receptor/G protein coupling results in increased cAMP and arachidonic acid (AA) production. cAMP stimulates the mobilization and transport of cholesterol to and into the mitochondria in part by activating PKA and MAPK signaling. At the same time, AA can be converted into prostaglandin by Cox2 to negatively regulate the transport of cholesterol across the mitochondrial membranes. At the inner mitochondrial membrane, cholesterol is converted to pregnenolone by CYP11A1, and pregnenolone is converted into testosterone by enzymes in the smooth endoplasmic reticulum (HSD3b, CYP17A1 and HSD17b). Aging and environmental factors may impact steroidogenesis by affecting the intracellular redox balance in part through increased ROS production. This has significant effects on cAMP formation and/or cholesterol transport into the mitochondria, and thus on steroid formation. Steroidogenesis also might be affected by ROS-induced increases in Cox2 production and redox-sensitive MAPK activation.

(1) maintenance of optimal levels of steroidogenic enzymes (trophic regulation) and (2) mobilization and transport of cholesterol into the inner mitochondrial membrane (acute regulation).

Both the trophic and acute effects of LH are mediated by signaling pathways that begin with cAMP production (Fig. 1). LH binds to and activates G protein-coupled receptors, resulting in the activation of adenylyl cyclase, increased intracellular cAMP formation and cAMP-dependent phosphorylation of proteins through protein kinase A (PKA). The acute stimulation of Leydig cells by LH results in cholesterol transfer into the mitochondria in part through the actions of steroidogenic acute regulatory protein (STAR), translocator protein (18kDa; TSPO) and other proteins of the transduceosome (Midzak *et al.* 2011). Cholesterol transport into the mitochondria, the rate-limiting step in steroid biosynthesis, is followed by the conversion of cholesterol to pregnenolone by the C27 cholesterol side-chain cleavage cytochrome P450 enzyme (CYP11A1)

located on the matrix side of the inner mitochondrial membrane. Pregnenolone then is metabolized into TS by 3β -hydroxysteroid dehydrogenase (3b-HSD; HSD3B), 17α -hydroxylase/17,20 lyase (CYP17A1) and type 3 17β -hydroxysteroid dehydrogenase (17b-HSD3, HSD17B) in the smooth endoplasmic reticulum (Payne & Hales 2004, Aghazadeh *et al.* 2015, Beattie *et al.* 2015).

Effects of aging on steroidogenesis

Aging and declining testosterone production by Leydig cells

With aging, serum TS levels decrease in the males of a number of species, including human and rodent (Harman *et al.* 2001, Mulligan *et al.* 2001, Keenan *et al.* 2006). Studies in men have shown a reduced ability of Leydig cells to produce TS in response to LH stimulation (Veldhuis *et al.* 2005, 2012, Surampudi *et al.* 2012). In many rat strains, including Sprague-Dawley, decreased

LH levels that result from hypothalamic–pituitary changes are responsible for decreased TS formation by aging Leydig cells, and thus, for decreased serum TS (Zirkin *et al.* 1993). This is referred to as secondary hypogonadism. In Brown Norway rats, however, as in men, TS levels decrease with age despite unchanging LH and increasing FSH levels (Zirkin *et al.* 1993, Chen *et al.* 1994, 1996, 2015a, Beattie *et al.* 2015). Some years ago, we showed that when Brown Norway rat testes were perfused with maximally stimulating LH, testes from old (18- to 24-month-old) rats produced significantly less TS than testes from young (3- to 6-month-old) rats (Zirkin *et al.* 1993). Consistent with this, Leydig cells isolated from aged rats produced less TS than cells from young rats in response to maximally stimulating LH (Chen *et al.* 1994). Leydig cells in the adult testis rarely turn over, and their numbers do not change or increase with age (Wang *et al.* 1993, Chen *et al.* 1994). Consequently, the reduced serum TS levels in old rats results from the relative unresponsiveness of the Leydig cells to LH (Chen *et al.* 2002).

Among the important changes in the steroidogenic pathway that have been implicated in the reduced TS production that characterizes aging Leydig cells are reductions in cAMP production and protein kinase A (PKA) activities (Chen *et al.* 2002, Liao *et al.* 1993). LH binding to LH receptors initiates a cascade of events that include receptor coupling to Gs proteins, activation of adenylyl cyclase and ultimately increased cAMP production and activation of cAMP-dependent PKA (Fig. 1). Signaling through cAMP/PKA is essential for the expression of the downstream steroidogenic proteins and enzymes of the mitochondria and smooth endoplasmic reticulum (Payne & Hales 2004). Reduced numbers of LH receptors or their reduced coupling efficiency to G-protein affects cAMP production and also affects other signaling cascades including arachidonic acid/cyclooxygenase-2 (COX-2) (Wang *et al.* 2005, Castillo *et al.* 2006) and mitogen-activated protein kinase (MAPK) (Tai & Ascoli 2011). cAMP, in response to LH, stimulates the transport of cholesterol into the mitochondria (Aghazadeh *et al.* 2015). Although the culture of old cells with LH was found to result in TS production significantly below that of LH-stimulated young cells, culture with dibutyryl cAMP (dbcAMP) for 3 days resulted in increased TS production by old cells to the level of LH-stimulated young cells (Chen *et al.* 2004a). This suggests that relative insensitivity of old Leydig cells to LH, and consequent deficient signal transduction, contributes to, or may cause, the reduced TS that characterizes aging Leydig cells.

What is the molecular mechanism by which cAMP production is reduced with aging? Stimulation of adenylyl cyclase in old Leydig cells with forskolin, which activates adenylyl cyclase directly via by-passing the ligand and G proteins, was found to result in equivalent production of cAMP by young and old cells (Chen *et al.* 2002).

Likewise, in the absence of LH, direct activation of Gs protein by cholera toxin stimulated cAMP synthesis in old cells to young levels, and inhibition of the inhibitory G protein (Gi) by pertussis toxin did not restore the ability of the aged Leydig cells to produce cAMP at high levels in response to LH (Chen *et al.* 2002). These observations suggest that G protein and adenylyl cyclase deficiencies are unlikely to cause aged cells to produce less cAMP than young cells. The number of LH-binding sites was shown to decrease significantly with age, indicating reduced plasma membrane LH receptor numbers (Chen *et al.* 2002). However, it has been known for some time that only approximately 10% of LH receptor occupancy is required to elicit a biological response, indicative of reserve or 'spare' receptors (Hsueh *et al.* 1977). In fact, Leydig cells from young, LH-suppressed rats, which have even lower numbers of LH receptors than aged cells, nonetheless were found to have an unaltered ability to synthesize cAMP in response to LH (Chen *et al.* 2002). These results, taken together, lead to the conclusion that LH receptor-G protein coupling deficiency is likely to be responsible, at least in part, for reduced cAMP production by aged Leydig cells. It also has been suggested that increased cAMP degradation may play a role (Sokanovic *et al.* 2014). In rodents, Leydig cell cAMP is metabolized by phosphodiesterases (PDE4a, PDE4b, and PDE2a) and found to be increased in aging Leydig cells (Sokanovic *et al.* 2014). However, the extent to which these increases might reduce cAMP levels in aged cells is unclear.

In addition to its effects on cAMP production, LH stimulation also results in arachidonic acid release in Leydig cells (Fig. 1) (Wang *et al.* 2005, Castillo *et al.* 2006). The released arachidonic acid is metabolized by cellular lipoxygenases, epoxygenases or cyclooxygenases (COX). The metabolites have been shown to be capable of modulating steroidogenesis in part by affecting STAR protein expression (Wang *et al.* 2005). In MA-10 Leydig cells and rat primary Leydig cells, inhibition of COX-2 activity resulted in significantly increased TS production, suggesting that COX-2 can negatively affect steroidogenesis (Wang *et al.* 2005, Chen *et al.* 2007). In aging Leydig cells, COX-2 message and protein levels have been shown to be elevated relative to their levels in cells from young rats (Wang *et al.* 2005, Chen *et al.* 2007), suggesting the possible involvement of COX-2 in age-related reductions in TS production (Wang *et al.* 2005, Chen *et al.* 2007). This is supported by the observation that long-term treatment of aged rats with COX-2 antagonist can partially reverse reduction in serum TS levels (Wang *et al.* 2005). These findings suggest the possibility that COX2 may also be involved in the age-related decline in TS production.

Members of the MAPK signaling family, including ERK and p38, also have been implicated in age-related reductions in steroidogenesis (Fig. 1) (Abidi *et al.* 2008a, Sokanovic *et al.* 2014). LH has been shown to

increase ERK1/2 phosphorylation (Tai & Ascoli 2011). Inhibition of ERK1/2 phosphorylation can significantly block the effect of LH stimulation, suggesting that ERK1/2 in part mediates LH function. Age-related modulation of the p38 MAPK signaling pathway has been shown in adrenal and Leydig cells to be associated with reductions in steroidogenesis elicited by prooxidants (Abidi *et al.* 2008a,b), and reductions in steroidogenesis can be partially prevented by inhibition of p38 (Chen *et al.* 2010). In a recent study, it was found that the oxidant-sensitive p38 MAPK can inhibit cAMP-induced steroidogenesis in part by repressing STAR (Zaidi *et al.* 2014), suggesting that age-related Leydig cell dysfunction may be related to changes in both MAPK and cAMP signaling (Sokanovic *et al.* 2014).

Cholesterol synthesis, mobilization and mitochondrial transport

In addition to the reductions in LH-stimulated cAMP in aged Brown Norway rat Leydig cells, there are reductions in STAR, TSPO, the mitochondrial enzyme CYP11A1 and downstream steroidogenic enzymes of the smooth endoplasmic reticulum (Luo *et al.* 1996, 2001, 2005, Culty *et al.* 2002). Although the mitochondrial and smooth endoplasmic reticulum steroidogenic enzymes are reduced in aging Leydig cells, their levels nonetheless are sufficient to support high levels of steroid production if enough cholesterol is translocated into the mitochondria and thus is available to CYP11A1 of the inner mitochondrial membrane (Culty *et al.* 2002). This suggests that defects in cholesterol import into the mitochondria might underlie the differential steroidogenic abilities of young vs old Leydig cells and may ultimately be responsible for the reduced TS formation in response to LH that characterizes the old cells. Cholesterol translocation involves mobilization of cholesterol and its transport to the outer and then the inner mitochondrial membrane where CYP11A1 resides (Aghazadeh *et al.* 2015, Venugopal *et al.* 2016).

The precise location of the intracellular cholesterol utilized in steroid formation is uncertain. Studies have shown that cholesterol utilized in steroidogenic cells derives in part from lipoprotein in the circulation that is imported by membrane-bound scavenger receptor class B type I (SR-B1) lipoprotein receptor (Azhar & Reaven 2002) or is synthesized *de novo* within the cells from acetyl-CoA. Cholesterol may be stored in cytoplasmic lipid droplets in the form of cholesteryl esters. A recent study reported that the plasma membrane is a major source of cholesterol for steroidogenesis (Venugopal *et al.* 2016). In response to hormonal stimulation, cholesteryl esters are converted into free cholesterol by cholesterol esterases and then transferred to the mitochondrial outer membrane (Shen *et al.* 2003). Important components in this process, including SR-B1, carboxyesterase (ES-10) and

hormone-sensitive lipase (HSL) are downregulated in aged Leydig cells (Chen *et al.* 2004b), suggesting that cholesterol import, synthesis and mobilization are all affected by aging. This is consistent with the observations that cholesterol synthesis and mobilization are reduced in Leydig cells of aged as compared to young rats (Liao *et al.* 1993).

Cholesterol translocation into mitochondria is mediated by cAMP through the proteins of the transduceosome (Papadopoulos *et al.* 2015). Translocator protein (TSPO), one such protein, comprises 2% of the outer mitochondrial membrane proteins of young Leydig cells. TSPO is reduced significantly in old Leydig cells (Culty *et al.* 2002). Incubating old Leydig cells with LH plus a specific TSPO drug ligand resulted in the restoration of TS production to the high levels of LH-stimulated young cells (Chung *et al.* 2013). In complementary *in vivo* studies, administering TSPO ligand to aged rats restored serum TS to the level of young rats. (Chung *et al.* 2013). A domain in the C-terminus of TSPO was characterized as a cholesterol recognition/interaction amino acid consensus (CRAC). These studies, among those conducted over many years, strongly implicate TSPO as a mediator of cholesterol translocation and thus steroidogenesis (Papadopoulos *et al.* 2015). However, recent studies of the effects of the genetic deletion of *Tspo* in mice have reported that TSPO knockout has little or no effect on steroidogenesis, thus calling into question the role of TSPO in cholesterol translocation (Tu *et al.* 2014). The latter results are disputed, however (Fan *et al.* 2015, Papadopoulos *et al.* 2015), and yet to be confirmed.

Several additional cytosolic proteins, including PKA, acyl-CoA binding domain-containing protein 3 (aka PAP7) and STAR may be involved in the import of cholesterol to the inner mitochondrial membrane (Papadopoulos *et al.* 2007, Miller 2013). STAR, for one, plays a significant role in steroidogenesis (Kallen *et al.* 1998, Clark 2016). When Leydig cells are stimulated by LH, STAR levels increase rapidly (Clark *et al.* 1994). In rodents, deletion of the *STAR* gene, or knockdown of the STAR protein, has been shown to suppress steroidogenesis (Lin *et al.* 1995). Impairment in the synthesis of adrenal and gonadal steroids has been shown to be associated with congenital lipoid adrenal hyperplasia in the human, a disease characterized in part by mutations of the *STAR* gene (Bose *et al.* 1997). Although the importance of STAR in mitochondrial cholesterol transfer is clear, the mechanism by which it functions remains uncertain. Early *in vitro* studies showed that STAR contains a domain (StAR-related lipid-transfer domain, or START) that binds cholesterol and may be involved in cholesterol transfer (Ponting & Aravind 1999) and thus in steroid production (Arakane *et al.* 1996). *In vivo*, the expression of a full-length STAR transgene in *STAR*-knockout mice restored adrenal and gonadal steroidogenesis (Sasaki *et al.* 2008).

These observations, and the localization of STAR at the outer mitochondrial membrane, suggest a critical role for STAR in cholesterol translocation (Clark 2016, Stocco *et al.* 2017). Moreover, recent reports of functional interaction between TSPO and STAR have led to the proposal that such interaction may be an integral part of cholesterol translocation to and into the mitochondria in steroidogenic cells (Liu *et al.* 2006, Miller 2013, Papadopoulos *et al.* 2015).

There are other potential players in steroid formation that are part of the transducesome. These include the 14-3-3 proteins. It has been shown that 14-3-3 γ is increased fourfold upon hCG stimulation and that it binds to STAR and acts as a negative regulator of steroid formation (Midzak *et al.* 2015). Each of TSPO, STAR, 14-3-3 γ and perhaps other transducesome proteins might serve as targets to regulate and thus enhance TS formation by old Leydig cells in a more physiological fashion than providing TS exogenously.

Mechanism(s) responsible for age-related changes in testosterone formation

Extrinsic factors

Cellular aging can result from changes in factors within and/or outside the target cells. Intrinsic factors, including reactive oxygen species (ROS), are discussed below. Extrinsic factors might include hormones, growth factors, oxidants and antioxidants produced by local cells or transported to the target cells via the circulation. Extrinsic factors have been shown to contribute to the aging of cells of the immune system (Badowski *et al.* 2014), skin (Lephart 2016) and muscle (Cannon 1998), among others, and to affect Leydig cells as well. For example, after the elimination of Leydig cells from the testes of young (3 months old) and aged (18 months old) Brown Norway rats by administering ethane dimethanesulfonate (EDS), a new generation of Leydig cells was restored to the aged testes by 10 weeks that produced as much TS as cells restored to the young testes (Chen *et al.* 1996). However, TS production by the newly formed Leydig in the testes of old rats was reduced to 'old' levels 20 weeks later. This was in striking contrast to the new cells formed in young testes, the TS production of which did not change in that period of time (Chen *et al.* 2015a). These observations suggested that factors outside the newly formed cells themselves may play important roles in their aging.

Leydig cell steroidogenic function has been shown to be affected by factors produced locally, and also at a distance via the circulatory system. Within the testis, Sertoli cell and macrophage products have been shown to have stimulatory effects on Leydig cell steroidogenesis (Haider 2004). Negative effects also have been shown. For example, hydrogen peroxide produced by macrophages may negatively impact

neighboring Leydig cells over time (Hales 2002). Factors produced by the pituitary (gonadotropins), liver (IGF-1), thyroid (T3), pancreas β -cells (insulin), bone marrow (osteocalcin) and immune system reach the Leydig cells via the circulatory system. It remains to be determined how changes in these extrinsic factors affect the steroidogenic function of aging Leydig cells.

Intrinsic factors: oxidants and antioxidant molecules

Imbalance between prooxidants and antioxidants often occurs as cells age (Rebrin *et al.* 2003). Such imbalance can result in an altered redox state and an accumulation of oxidative damage to intracellular macromolecules, thus contributing to age-related functional deficits (Finkel & Holbrook 2000). Leydig cells produce ROS from several sources, including the mitochondrial electron transport chain and mitochondrial and microsomal cytochrome P450 enzyme reactions (Hanukoglu 2006). Recent studies indicate ROS production might be in response to LH, incubating Leydig cells with LH resulted in increased ROS levels as well as in DNA damage (Beattie *et al.* 2013). Leydig cells from aged rats produce significantly more reactive oxygen than cells from young rats, and this occurs despite reduced mitochondrial volume (Chen *et al.* 2001). Aging of Leydig cells also is accompanied by the reduced expressions of key enzymatic and non-enzymatic antioxidants, including Cu-Zn-SOD, Mn-SOD, glutathione peroxidase (GPX-1), microsomal glutathione S-transferase (MGST1), glutathione S-transferase (GSTM2) and glutathione (GSH), leading to increased oxidative stress and oxidative damage (e.g. lipid peroxidation) (Chen *et al.* 2001, Cao *et al.* 2004). Age-related decreases in Leydig cell antioxidant activities, gene expression and protein levels are consistent with the hypothesis that the loss of steroidogenic function that accompanies Leydig cell aging may result in part from an altered antioxidant defense system (Cao *et al.* 2004, Luo *et al.* 2006).

Experimental studies designed to go beyond correlation to establish cause-effect relationships have been reported. The long-term administration of the antioxidant vitamin E delayed age-related decreases in steroidogenesis, while long-term vitamin E deficiency had the opposite effect (Abidi *et al.* 2004, Chen *et al.* 2005). GSH is among the most significant antioxidants reduced in aging Leydig cells (Cao *et al.* 2004, Luo *et al.* 2006). To examine whether loss of GSH would affect Leydig cell TS production *in vivo*, young and old rats were treated with buthionine sulfoximine (BSO) for a week to reduce Leydig cell intracellular GSH levels. Significantly decreased TS production by both young and old Leydig cells was seen following experimental reduction of GSH, suggesting that such reductions might lead to alterations in the redox environment of Leydig cells and ultimately to decreases in TS production

(Chen *et al.* 2008). Depletion of GSH *in vitro* also reduced Leydig cell steroidogenic function, while the antioxidants vitamin E, N-tert-butyl- α -phenylnitron and Trolox suppressed the effect of loss of GSH (Chen *et al.* 2008). A critical role of GSH in regulation of Leydig cell steroidogenesis was also suggested by a recent study reporting that the expression of γ -glutamyl transferase 5 (GGT5), an enzyme involved in GSH metabolism, was negatively correlated with Leydig cell steroidogenesis (Li *et al.* 2016).

Nrf2/Keap1 signaling is an important mechanism by which cells respond to oxidative stress (Cho *et al.* 2005, Li & Kong 2009). Normally, Nrf2 is bound to Keap1 in the cytoplasm and undergoes ubiquitination-dependent proteasomal degradation. With exposure to oxidative stress, Nrf2 dissociates from Keap1. When separated from Keap1, Nrf2 binds to the antioxidant response element (ARE) in the promoter of target genes, and thus stimulates the transcription of numerous antioxidant molecules and phase two enzymes (Cho *et al.* 2005, Li & Kong 2009). With aging, GSH and antioxidant enzymes typically decline despite elevated levels of oxidative stress. This is true of many cell types, including Leydig cells (Rebrin *et al.* 2003, Cao *et al.* 2004, Luo *et al.* 2006), and leads to a prooxidant state in aging cells. We reasoned that if Nrf2 and thus the oxidant/antioxidant environment plays a role in age-related reductions in Leydig cell TS production, the experimental knockout of Nrf2 should result in increased oxidative stress and therefore, over time, in decreased TS production. Using this genetic approach, we found that at 3 months, there was no significant difference between wild-type and knockout mice in either serum TS concentration or in the ability of Leydig cells to produce TS. However, by middle age (8 months), at which time cells from wild-type mice had not yet lost steroidogenic function, TS production by Leydig cells from the knockout mice was reduced significantly. By 24 months, both the wild-type and knockout mice had reduced serum TS and Leydig cell TS production, but with more extensive reductions in the knockouts (Chen *et al.* 2015b). These observations indicate that without the antioxidant master regulator Nrf2, the age-related loss of Leydig cell steroidogenic function accelerated. Moreover, the antioxidant capacity of the testis was significantly reduced in the knockout as compared to the wild-type mice at each age. With reduced expression of numerous antioxidant molecules and therefore reduced total antioxidant capacity, an increasingly prooxidant environment was seen that was reminiscent of aging, accompanied by reduced TS production by Leydig cells and reduced serum TS levels. The specific targets of an altered redox environment, and thus, the mechanism(s) by which a prooxidant environment affects Leydig cell steroidogenesis, remain unclear.

Effects of environmental factors on steroidogenesis

There is growing realization that many of the man-made chemicals released into the atmosphere have significant public health consequences, including perturbation of the endocrine system. Endocrine-disrupting chemicals (EDCs) are defined as 'substances in our environment, food and consumer products that interfere with hormone biosynthesis, metabolism or action resulting in a deviation from normal homeostatic control or reproduction' (Diamanti-Kandarakis *et al.* 2009). In this section, we will briefly discuss the effects of EDC on Leydig cell steroidogenesis, and the striking similarity in mechanisms by which these effects and those that result in age-related reduction in TS production are elicited.

Effects of EDCs on cAMP production and mitochondrial cholesterol metabolism

As indicated previously, LH-stimulated cAMP production and mitochondrial cholesterol transport and metabolism have been shown to be critically involved in age-related reductions in TS formation (Fig. 1). These same elements of the steroidogenic pathway are affected by a number of EDCs, leading to reduced TS (summarized in Table 1). Using mouse Leydig tumor cells as a model system, 2,2',4,4'-tetrabromodiphenyl ether (BDE-47), a polybrominated diphenyl ether, was found to elicit decreases in hCG-induced cAMP levels and in the synthesis of CYP11A1 (Han *et al.* 2012). Exposure of rat primary Leydig cells to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) also was found to inhibit steroidogenesis through its effects on cAMP and CYP11A1 (Johnson *et al.* 1992, Fukuzawa *et al.* 2004). Permethrin, an insecticide, disrupted TS formation in mice by decreasing the protein and mRNA levels of STAR and CYP11A1, also indicative of an effect on cholesterol transport and metabolism (Zhang *et al.* 2007). Atrazine, a widely used herbicide, has similar effects; exposure of rats to atrazine during postnatal days 23–30 caused reduced cAMP levels in Leydig cells and decreased cholesterol transport (Pogrmic *et al.* 2009). Exposure of Leydig cells to triclosan, a chemical widely used in antimicrobial preparations, resulted in significantly decreased activity of adenyl cyclase, synthesis of cAMP and transcriptional and translational activity of P450_{scc}, 3 β -HSD, 17 β -HSD and STAR (Kumar *et al.* 2008). Lindane, an insecticide that also is used to treat lice and scabies, also was found to affect cAMP, STAR and steroidogenesis (Ronco *et al.* 2001, Saradha *et al.* 2008). Phthalates are plasticizers that are highly abundant, exhibit antiandrogenic properties and have different effects depending upon when in the lifecycle they are administered. Thus, the exposure of males to phthalates *in utero* can affect both Leydig cell development and adult steroidogenic function (Foster *et al.* 2001, Martinez-Arguelles *et al.* 2013). MEHP was found to inhibit steroidogenesis in

Table 1 Environmental factors affecting LHR-cAMP and mitochondrial cholesterol cascades.

Toxin	Animal cell	Age	Dose (/kgBW) duration	Effects on (LC or Testis)	Mechanisms sensitive targets	Reference
BDE-47	MA10		1 µM; 24 h	P4 production↓	cAMP↓; Cyp11a1↓	Han (2012)
TCDD	Mice	Adult	100 µg; once	Testicular TS↓	LHR↓; Cyp11a1↓	Fukuzawa (2004)
Permethrin	Mice	Adult	35 mg; 6 week	Serum TS↓ Testicular TS↓	TSPO↓; STAR↓; Cyp11a1↓	Zhang (2007)
Atrazine	Rat	Pnd23-50	200 mg; 4 week	TS+DHT productions↓	cAMP↓; LHR↓; SR-B1↓ SF-1↓; STAR↓; Cyp17a1↓; Hsd17b3↓	Pogrmic (2009)
Triclosan	Rat LCs	Adult	0.01 µM; 2 h	TS production↓	cAMP↓; STAR↓; Cyp17a1↓ Cyp11a1↓; Hsd17b3↓; Hsd3b1↓ Preventable by forskolin	Kumar (2008)
Lindane	Rat LCs	Adult	10 µg/mL; 3 h	TS production↓	cAMP↓	Ronco (2001)
PFDaA	mLTC-1/rat LCs		10 µM; 24 h	P4 production ↓ TS production↓	STAR↓	Shi (2010)
Salinomycin	Mice	Adult	3 mg; 4 week	Testicular TS↓ Testis W↓ Vesicles W↓ Epididymis W↓	STAR↓; Cyp11a1↓	Ojo (2013)
Quinalphos	Mice	Adult	7.5 mg; 7 day	Serum TS↓ Testis W↓ Vesicles W↓ Epididymis W↓ Prostrate W↓	STAR↓; Cyp11a1↓	Kokilavani (2014)
MEHP	MA-10		200 µM; 24 h	P4 production↓	cAMP↓; STAR↓	Zhou (2013)
Cobalt	MA10		100 µM; 24 h	P4 production↓	Cyp11a1	Kumar (2014)
Dimethoate	Rat	Adult	15 mg; 5 week	Serum TS↓ LH↑ FSH↑	STAR↓ Hsd17b3↓; Hsd3b1↓	Astiz (2009)
Dimethoate	Rat LCs		1 ppm; 24 h	TS production↓	STAR↓ Hsd17b3↓; Hsd3b1↓	Astiz (2012)

BDE-47, 2,2',4,4'-Tetrabromodiphenyl Ether; Cyp11a1, Cholesterol side-chain cleavage; Cyp17a1, Cytochrome P450 17a1 (Steroid 17 α -hydroxylase/17,20 lyase); DEHP, Diethylhexyl Phthalate; DHT, Dihydrotestosterone; Hsd17b3, Hydroxysteroid 17-Beta Dehydrogenase 3; LHR, Luteinizing Hormone/choriogonadotropin Receptor; MEHP, Monoethylhexyl phthalic acid; P4, Progesterone; PFDaA, Perfluorododecanoic acid; Pnd, Postnatal day; SF-1, Steroidogenic Factor 1 (NR5A1); SR-B1, Scavenger Receptor class B member 1; STAR, Steroidogenic Acute Regulatory protein; TS, Testosterone; TCDD, 2,3,7,8-Tetrachlorodibenzodioxin; TSPO, Translocator Protein.

MA-10 Leydig cells by interfering with LH-stimulated cAMP production and mitochondrial cholesterol transport/metabolism (Zhou *et al.* 2013).

Reduced testosterone formation in response to environmental factors: mechanisms

Although EDCs capable of affecting steroidogenesis are diverse in nature, an altered redox environment that leads to increased oxidative stress and thus reduced steroidogenesis are commonly seen as consequences of exposures, as they are in aging (summarized in Table 2, Mathur & D'Cruz 2011). For example, exposure of mLTC-1 cells to perfluorododecanoic acid (PFDaA) was shown to result in increased levels of ROS and hydrogen peroxide and in the inhibition of STAR expression and steroidogenesis (Shi *et al.* 2010). Exposing purified rat Leydig cells to the polychlorinated biphenyl Aroclor 1254 resulted in significant decline in the activities of enzymatic and non-enzymatic antioxidant enzymes, increase in the levels of ROS and decrease in the mRNA levels of steroidogenic enzymes (Murugesan *et al.* 2008). These changes were partially prevented by the

antioxidants vitamin C and E (Murugesan *et al.* 2005). Exposure of mice to salinomycin resulted in decreased testicular TS in association with depletion of SOD and GSH and increased lipid peroxidation (Ojo *et al.* 2013). Exposure of rats to chlorpyrifos resulted in reduced TS levels, increased oxidative stress and decreased antioxidant enzymes (Sai *et al.* 2014). The testicular antioxidant defense system and lipid peroxidation were also significantly affected by quinalphos exposure (Debnath & Mandal 2000, Kokilavani *et al.* 2014).

Phthalates are another group of environmental contaminants that may affect steroidogenesis by increasing oxidative stress (Martinez-Arguelles *et al.* 2013). One of the well-established consequences of phthalate exposure is activation of peroxisome proliferator-activated receptor (PPAR)- α and - γ nuclear receptors (Hurst & Waxman 2003). Activation of PPARs not only affected lipid metabolism but also increased oxidative stress in the cells (O'Brien *et al.* 2005, Zhang *et al.* 2016). Exposure of rats to DEHP during prepuberty and puberty significantly decreased the GSH/GSSG ratio and increased TBARS levels in the testis, suggesting that phthalate is capable of

Table 2 Environmental factors affecting redox environment of Leydig cells.

Toxin	Animal cell	Age	Dose (/kgBW); duration	Effects on (LC or Testis)	Mechanisms sensitive targets	Reference
Lindane	Rat	Adult	5 mg; 12 h	StAR↓ Hsd17b3↓; Hsd3b1↓	H ₂ O ₂ ↑	Saradha (2008)
PFD ₀ A	mLTC-1/rat LCs		10 μM; 24 h	P4 production ↓ TS production↓	StAR↓; H ₂ O ₂ ↑ Preventable by MnTMPyP	Shi (2010)
Aroclor1254	rat LCs	Adult	10 ⁻⁸ M; 6 h	TS production↓ Cyp11a1↓ Hsd17b3↓ Hsd3b1↓	H ₂ O ₂ ↑; LPO↑ Hydroxyl radical↑; SOD↓ CAT↓; GPx↓; γ-GT↓; GR↓ GST↓; Vitamin C and E↓	Murugesan (2008)
Salinomycin	Mice	Adult	3 mg; 4 week	Testicular TS↓ StAR↓ Cyp11a1↓	LPO↑ GSH↓; CAT↓; LDH↓; SOD↓	Ojo (2013)
Quinalphos	Rat	Adult	250 μg; 3 day	Serum TS↓ Testis W↓ Germ cell loss↑	LPO↑; SOD↑; GPx ↑ CAT↓; GSH↓	Debnath (2000)
Quinalphos	Mice	Adult	7.5 mg; 7 day	Serum TS↓ StAR↓; Cyp11a1↓ Hsd17b3↓ Hsd3b1↓	SOD↓; CAT↓; GPx↓ Vitamin C↓; LDH↑ Preventable by antioxidant <i>Cissus quadrangularis</i>	Kokilavani (2014)
DEHP	Rat	Pnd21	1000 mg; 10 day	Serum TS↓	Serum LH↓; Serum FSH↓ Preventable by selenium diet;	Erkekoglu (2011)
DEHP	Rat	Pnd21	1000 mg; 10 day	GSH/GSSG↓ TBARS↑	Cu,Zn-SOD↓; CAT↑ GSH/GSSG↓; GPx4↓ Preventable by selenium diet	Erkekoglu (2014)
MEHP/ DEHP	MA-10		3 mM; 24 h	ROS↑ DNA damage↓	GPx1↓; TrxR↓; GST↓ GSH↓; p53↑ Selenium dependent	Erkekoglu (2010)
MEHP	MA-10		1 μM; 48 h	P4 production↓ TS production↓	ROS↑; Cyp1a1 network↑	Fan (2010)
MEHP	MA-10		200 μM; 24 h	P4 production↓ cAMP↓; StAR↓	ROS↑ GSH depended Vitamin E protective	Zhou (2013)
Mercury	Rat	Adult	50 ppm; 90 day	Serum TS↓ Sperm count↓	LPO↑; TBARS↑ SOD↓; CAT↓	Boujbiha (2009)
Lead	Rat	Adult	25 μg; 15 day	Serum TS↓ Hsd17b3↓ Hsd3b1↓	LPO↑; GSH↓; GST↓; SOD↓ CAT↓; GPx↓; TBARS↑ Preventable by Vitamin C	Pandya (2012)
Cadmium	Rat	Adult	25 μg; 15 day	Serum TS↓ Hsd17b3↓ Hsd3b1↓	LPO↑; GSH↓; GST↓; SOD↓ CAT↓; GPx↓; TBARS↑ Preventable by Vitamin C	Pandya (2012)
Cobalt	MA10		100 μM; 24 h	P4 production↓ Cyp11a1↓	ROS↑ HIF-1a activity↑ Preventable by PDTC	Kumar (2014)
Arsenite	Mice	Adult	11.5 ppm; 36 day	Serum TS↓ Hsd17b3↓ Hsd3b1↓	GSH↓ Preventable by Vitamin C	Chang (2007)
BPA	R2C		0.1 nM; 24 h	TS production↓ Aromatase↑	COX2↑; PGE2↑ EP2↑; EP4↑; CREB↑ Preventable by inhibitions of PKA/ Akt/ ERK/ JNK/ p38	Kim (2010)
Dimethoate	Rat	Adult	15 mg; 5 week	Serum TS↓ StAR↓ LH↑	LPO↑; PGE2↑; PGF2α↑ COX2↑; α-tocopherol↓ Preventable by TROLOX or rofecoxib	Astiz (2009)
Dimethoate	Rat LCs		1 ppm; 24 h	TS production↓ StAR↓	LPO↑; COX2↑ Preventable by PUFA	Astiz (2012)

BPA, Bisphenol A; CAT, Catalase; COX2, Cyclooxygenase-2; CREB, cAMP Response Element Binding protein; EP2, Prostaglandin E2 receptor 2; EP4, Prostaglandin E2 receptor 4; GPx, Glutathione Peroxidase; GR, Glutathione Reductase; GST, Glutathione-S-Transferase; γ-GT, γ-Glutamyl Transpeptidase; HIF-1a, Hypoxia Inducible Factor 1, alpha subunit; LDH, Lactate Dehydrogenase; LPO, Lipid peroxidation; MnTMPyP, Manganese (III) tetrakis(1-methyl-4-pyridyl)porphyrin pentachloride; PGE2, Prostaglandin E2; PUFA, Polyunsaturated Fatty Acids; Ppm, Parts per million; SOD, Superoxide Dismutase; TBARS, Thiobarbituric Acid Reactive Substances; TrxR, Thioredoxin Reductase.

increasing oxidative stress and affecting the redox environment *in vivo* (Erkekoglu *et al.* 2014). In MA-10 cells, MEHP treatment resulted in increased ROS levels, while depletion of GSH by BSO pretreatment greatly exacerbated the MEHP induced-ROS production and resulted in reduced progesterone production (Zhou *et al.* 2013). These results suggest that a likely mechanism by which MEHP acts is through increased oxidative stress (Zhao *et al.* 2012).

There are metals in the environment that also have been shown to affect Leydig cell steroidogenesis, with suggestions that they may do so by increasing oxidative stress. For example, mercury affects Leydig cell steroidogenesis and has effects on the antioxidant defense system (Boujbiha *et al.* 2009). Vitamin E was shown to protect against mercury-induced toxicity in mice (Rao & Sharma 2001) and sodium selenite and/or vitamin E to result in reduced lipid peroxidation, increased superoxide dismutase, catalase and glutathione peroxidase activities and reduced histopathological lesions (Kalender *et al.* 2013). Lead and cadmium, either alone or in combination, were found to disrupt the testicular steroidogenesis and antioxidant defense mechanisms, and the administration antioxidant agents were found to reduce metal-induced oxidative stress and to provide protection against lead and cadmium toxicity (Liu *et al.* 2009, 2013, Ayinde *et al.* 2012, Pandya *et al.* 2012). Exposure of MA-10 Leydig cells to cobalt chloride resulted in an increase in the production of ROS and a decrease in progesterone production (Kumar *et al.* 2014). The adverse effects of arsenite on the male reproductive system also may be mediated by oxidative stress; the exposure of mice to arsenite reduced testicular GSH levels and increased protein carbonyl content, accompanied by decreases in testicular steroidogenic enzyme activities, and changes induced by arsenite were partially prevented by the antioxidant ascorbic acid (Chang *et al.* 2007).

In addition to modifying intracellular oxidative stress, some environmental compounds also have been shown to affect COX-2, arachidonate metabolism and MAPK signaling molecules. For example, bisphenol A (BPA) induced a decrease in TS production in rat Leydig R2C cells, associated with increased COX-2 and MAPK signaling (Kim *et al.* 2010). Dimethoate, a widely used organophosphate, was reported to reduce Leydig cell steroidogenic function in association with decreased arachidonate and increased COX-2 (Astiz *et al.* 2009, 2012).

Summary and conclusions

Leydig cell TS production decreases with aging and exposure to environmental contaminants. Although the exact mechanisms responsible for changes in steroidogenesis remain uncertain, there appear to be

common causative features. Both aging and a number of EDC exposures have effects on LH-stimulated cAMP production and cholesterol transport to and metabolism within in mitochondria. Increased oxidative stress appears to be responsible for at least some of the changes in the steroidogenic pathway. COX-2 and MAPK also may be involved. In addition to intrinsic mechanisms, aging and environmental factors also may affect Leydig cell steroidogenesis through endocrine, autocrine and paracrine mechanisms. In light of increased oxidative stress that accompanies aging, it would be of particular interest to determine whether EDC exposures affect aged cells differently than young cells. As yet, this has received little attention.

Declaration of interest

The authors declare no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by NIH grant R37 AG21092 (B R Z), National Natural Science Foundation of China grants NSFC81471411 (H C) and NSFC81102150 (L Y), and Natural Science Foundation of Zhejiang Province grant Y17H040047 (H C)

References

- Abidi P, Leers-Sucheta S & Azhar S 2004 Suppression of steroidogenesis and activator protein-1 transcription factor activity in rat adrenals by vitamin E deficiency-induced chronic oxidative stress. *Journal of Nutritional Biochemistry* **15** 210–219. (doi:10.1016/j.jnutbio.2003.11.007)
- Abidi P, Leers-Sucheta S, Cortez Y, Han J & Azhar S 2008a Evidence that age-related changes in p38 MAP kinase contribute to the decreased steroid production by the adrenocortical cells from old rats. *Aging Cell* **7** 168–178. (doi:10.1111/j.1474-9726.2007.00364.x)
- Abidi P, Zhang H, Zaidi SM, Shen WJ, Leers-Sucheta S, Cortez Y, Han J & Azhar S 2008b Oxidative stress-induced inhibition of adrenal steroidogenesis requires participation of p38 mitogen-activated protein kinase signaling pathway. *Journal of Endocrinology* **198** 193–207. (doi:10.1677/JOE-07-0570)
- Aghazadeh Y, Zirkin BR & Papadopoulos V 2015 Pharmacological regulation of the cholesterol transport machinery in steroidogenic cells of the testis. *Vitamins and Hormones* **98** 189–227. (doi:10.1016/bs.vh.2014.12.006)
- Arakane F, Sugawara T, Nishino H, Liu Z, Holt JA, Pain D, Stocco DM, Miller WL & Strauss JF 3rd 1996 Steroidogenic acute regulatory protein (StAR) retains activity in the absence of its mitochondrial import sequence: implications for the mechanism of StAR action. *PNAS* **93** 13731–13736. (doi:10.1073/pnas.93.24.13731)
- Astiz M, Hurtado de Catalfo GE, de Alaniz MJ & Marra CA 2009 Involvement of lipids in dimethoate-induced inhibition of testosterone biosynthesis in rat interstitial cells. *Lipids* **44** 703–718. (doi:10.1007/s11745-009-3323-5)
- Astiz M, Hurtado de Catalfo G, de Alaniz MJ & Marra CA 2012 Exogenous arachidonate restores the dimethoate-induced inhibition of steroidogenesis in rat interstitial cells. *Lipids* **47** 557–569. (doi:10.1007/s11745-012-3669-y)

- Ayinde OC, Ogunnowo S & Ogedegbe RA 2012 Influence of vitamin C and vitamin E on testicular zinc content and testicular toxicity in lead exposed albino rats. *BMC Pharmacology and Toxicology* **13** 17–24. (doi:10.1186/2050-6511-13-17)
- Azhar S & Reaven E 2002 Scavenger receptor class BI and selective cholesteryl ester uptake: partners in the regulation of steroidogenesis. *Molecular and Cellular Endocrinology* **195** 1–26. (doi:10.1016/S0303-7207(02)00222-8)
- Badowski M, Shultz CL, Eason Y, Ahmad N & Harris DT 2014 The influence of intrinsic and extrinsic factors on immune system aging. *Immunobiology* **219** 482–485. (doi:10.1016/j.imbio.2014.02.008)
- Beattie MC, Chen H, Fan J, Papadopoulos V, Miller P & Zirkin BR 2013 Aging and luteinizing hormone effects on reactive oxygen species production and DNA damage in rat Leydig cells. *Biology of Reproduction* **88** 100–106. (doi:10.1095/biolreprod.112.107052)
- Beattie MC, Adekola L, Papadopoulos V, Chen H & Zirkin BR 2015 Leydig cell aging and hypogonadism. *Experimental Gerontology* **68** 87–91. (doi:10.1016/j.exger.2015.02.014)
- Bose HS, Pescovitz OH & Miller WL 1997 Spontaneous feminization in a 46,XX female patient with congenital lipoid adrenal hyperplasia due to a homozygous frameshift mutation in the steroidogenic acute regulatory protein. *Journal of Endocrinology and Metabolism* **82** 1511–1515. (doi:10.1210/jcem.82.5.3962)
- Boujbiha MA, Hamden K, Guerzazi F, Bouslama A, Omezzine A, Kammoun A & El Feki A 2009 Testicular toxicity in mercuric chloride treated rats: association with oxidative stress. *Reproductive Toxicology* **28** 81–89. (doi:10.1016/j.reprotox.2009.03.011)
- Cannon JG 1998 Intrinsic and extrinsic factors in muscle aging. *Annals of the New York Academy Science* **854** 72–77. (doi:10.1111/j.1749-6632.1998.tb09893.x)
- Cao L, Leers-Sucheta S & Azhar S 2004 Aging alters the functional expression of enzymatic and non-enzymatic anti-oxidant defense systems in testicular rat Leydig cells. *Journals of the Steroid Biochemistry and Molecular Biology* **88** 61–67. (doi:10.1016/j.jsmb.2003.10.007)
- Castillo AF, Maciel FC, Castilla R, Duarte A, Maloberti P, Paz C & Podesta EJ 2006 cAMP increases mitochondrial cholesterol transport through the induction of arachidonic acid release inside this organelle in Leydig cells. *FEBS Journal* **273** 5011–5021. (doi:10.1111/j.1742-4658.2006.05496.x)
- Chang SI, Jin B, Youn P, Park C, Park JD & Ryu DY 2007 Arsenic-induced toxicity and the protective role of ascorbic acid in mouse testis. *Toxicology and Applied Pharmacology* **218** 196–203. (doi:10.1016/j.taap.2006.11.009)
- Chen H, Hardy MP, Huhtaniemi I & Zirkin BR 1994 Age-related decreased Leydig cell testosterone production in the brown Norway rat. *Journal of Andrology* **15** 551–557. (doi:10.1002/j.1939-4640.1994.tb00498.x)
- Chen H, Huhtaniemi I & Zirkin BR 1996 Depletion and repopulation of Leydig cells in the testes of aging brown Norway rats. *Endocrinology* **137** 3447–3452. (doi:10.1210/endo.137.8.8754773)
- Chen H, Cangelo D, Benson S, Folmer J, Zhu H, Trush MA & Zirkin BR 2001 Age-related increase in mitochondrial superoxide generation in the testosterone-producing cells of Brown Norway rat testes: relationship to reduced steroidogenic function? *Experimental Gerontology* **36** 1361–1373. (doi:10.1016/S0531-5565(01)00118-8)
- Chen H, Hardy MP & Zirkin BR 2002 Age-related decreases in Leydig cell testosterone production are not restored by exposure to LH in vitro. *Endocrinology* **143** 1637–1642. (doi:10.1210/endo.143.5.8802)
- Chen H, Liu J, Luo L & Zirkin BR 2004a Dibutyl cyclic adenosine monophosphate restores the ability of aged Leydig cells to produce testosterone at the high levels characteristic of young cells. *Endocrinology* **145** 4441–4446. (doi:10.1210/en.2004-0639)
- Chen H, Irizarry RA, Luo L & Zirkin BR 2004b Leydig cell gene expression: effects of age and caloric restriction. *Experimental Gerontology* **39** 31–43. (doi:10.1016/j.exger.2003.09.021)
- Chen H, Liu J, Luo L, Baig MU, Kim JM & Zirkin BR 2005 Vitamin E, aging and Leydig cell steroidogenesis. *Experimental Gerontology* **40** 728–736. (doi:10.1016/j.exger.2005.06.004)
- Chen H, Luo L, Liu J & Zirkin BR 2007 Cyclooxygenases in rat Leydig cells: effects of luteinizing hormone and aging. *Endocrinology* **148** 735–742. (doi:10.1210/en.2006-0925)
- Chen H, Pechenino AS, Liu J, Beattie MC, Brown TR & Zirkin BR 2008 Effect of glutathione depletion on Leydig cell steroidogenesis in young and old brown Norway rats. *Endocrinology* **149** 2612–2619. (doi:10.1210/en.2007-1245)
- Chen H, Zhou L, Lin CY, Beattie MC, Liu J & Zirkin BR 2010 Effect of glutathione redox state on Leydig cell susceptibility to acute oxidative stress. *Molecular and Cellular Endocrinology* **323** 147–154. (doi:10.1016/j.mce.2010.02.034)
- Chen H, Guo J, Ge R, Lian Q, Papadopoulos V & Zirkin BR 2015a Steroidogenic fate of the Leydig cells that repopulate the testes of young and aged Brown Norway rats after elimination of the preexisting Leydig cells. *Experimental Gerontology* **72** 8–15. (doi:10.1016/j.exger.2015.08.014)
- Chen H, Jin S, Guo J, Kombairaju P, Biswal S & Zirkin BR 2015b Knockout of the transcription factor Nrf2: effects on testosterone production by aging mouse Leydig cells. *Molecular and Cellular Endocrinology* **409** 113–120. (doi:10.1016/j.mce.2015.03.013)
- Cho HY, Reddy SP, Debiase A, Yamamoto M & Kleeberger SR 2005 Gene expression profiling of NRF2-mediated protection against oxidative injury. *Free Radical Biology and Medicine* **38** 325–343. (doi:10.1016/j.freeradbiomed.2004.10.013)
- Chung JY, Chen H, Midzak A, Burnett AL, Papadopoulos V & Zirkin BR 2013 Drug ligand-induced activation of translocator protein (TSPO) stimulates steroid production by aged brown Norway rat Leydig cells. *Endocrinology* **154** 2156–2165. (doi:10.1210/en.2012-2226)
- Clark BJ 2016 ACTH action on StAR biology. *Frontiers in Neuroscience* **10** 547. (doi:10.3389/fnins.2016.00547)
- Clark BJ, Wells J, King SR & Stocco DM 1994 The purification, cloning, and expression of a novel luteinizing hormone-induced mitochondrial protein in MA-10 mouse Leydig tumor cells characterization of the steroidogenic acute regulatory protein (StAR). *Journal of Biological Chemistry* **269** 28314–28322.
- Culty M, Luo L, Yao ZX, Chen H, Papadopoulos V & Zirkin BR 2002 Cholesterol transport, peripheral benzodiazepine receptor, and steroidogenesis in aging Leydig cells. *Journal of Andrology* **23** 439–447. (doi:10.1002/j.1939-4640.2002.tb02251.x)
- Debnath D & Mandal TK 2000 Study of quinalphos (an environmental oestrogenic insecticide) formulation (Ekalux 25 E.C.)-induced damage of the testicular tissues and antioxidant defence systems in Sprague-Dawley albino rats. *Journal of Applied Toxicology* **20** 197–204. (doi:10.1002/(SICI)1099-1263(200005/06)20:3<197::AID-JAT634>3.0.CO;2-7)
- Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM, Zoeller RT & Gore AC 2009 Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocrine Reviews* **30** 293–342. (doi:10.1210/er.2009-0002)
- Erkekoglu P, Rachidi W, Yuzugullu OG, Giray B, Favier A, Ozturk M & Hincal F 2010 Evaluation of cytotoxicity and oxidative DNA damaging effects of di(2-ethylhexyl)-phthalate (DEHP) and mono(2-ethylhexyl)-phthalate (MEHP) on MA-10 Leydig cells and protection by selenium. *Toxicology of Applied Pharmacology* **248** 52–62. (doi:10.1016/j.taap.2010.07.016)
- Erkekoglu P, Zeybek ND, Giray B, Asan E, Arnaud J & Hincal F 2011 Reproductive toxicity of di(2-ethylhexyl) phthalate in selenium-supplemented and selenium-deficient rats. *Drug and Chemical Toxicology* **34** 379–389. (doi:10.3109/01480545.2010.547499)
- Erkekoglu P, Giray B, Rachidi W, Hinginger-Favier I, Roussel AM, Favier A & Hincal F 2014 Effects of di(2-ethylhexyl)phthalate on testicular oxidant/antioxidant status in selenium-deficient and selenium-supplemented rats. *Environmental Toxicology* **29** 98–107. (doi:10.1002/tox.20776)
- Fan J, Traore K, Li W, Amri H, Huang H, Wu C, Chen H, Zirkin B & Papadopoulos V 2010 Molecular mechanisms mediating the effect of mono-(2-ethylhexyl) phthalate on hormone-stimulated steroidogenesis in MA-10 mouse tumor Leydig cells. *Endocrinology* **151** 3348–3362. (doi:10.1210/en.2010-0010)
- Fan J, Campioli E, Midzak A, Culty M & Papadopoulos V 2015 Conditional steroidogenic cell-targeted deletion of TSPO unveils a crucial role in viability and hormone-dependent steroid formation. *PNAS* **112** 7261–7266. (doi:10.1073/pnas.1502670112)
- Finkel T & Holbrook NJ 2000 Oxidants, oxidative stress and the biology of ageing. *Nature* **408** 239–247. (doi:10.1038/35041687)
- Foster PM, Mylchreest E, Gaido KW & Sar M 2001 Effects of phthalate esters on the developing reproductive tract of male rats. *Human Reproduction Update* **7** 231–235. (doi:10.1093/humupd/7.3.231)

- Fukuzawa NH, Ohsako S, Wu Q, Sakaue M, Fujii-Kuriyama Y, Baba T & Tohyama C 2004 Testicular cytochrome P450_{sc} and LHR as possible targets of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the mouse. *Molecular and Cellular Endocrinology* **221** 87–96. (doi:10.1016/j.mce.2004.02.005)
- Haider SG 2004 Cell biology of Leydig cells in the testis. *International Review of Cytology* **233** 181–241. (doi:10.1016/s0074-7696(04)33005-6)
- Hales DB 2002 Testicular macrophage modulation of Leydig cell steroidogenesis. *Journal of Reproductive Immunology* **57** 3–18. (doi:10.1016/S0165-0378(02)00020-7)
- Han X, Tang R, Chen X, Xu B, Qin Y, Wu W, Hu Y, Xu B, Song L, Xia Y et al. 2012 2,2',4,4'-Tetrabromodiphenyl ether (BDE-47) decreases progesterone synthesis through cAMP-PKA pathway and P450_{sc} downregulation in mouse Leydig tumor cells. *Toxicology* **302** 44–50. (doi:10.1016/j.tox.2012.07.010)
- Hanukoglu I 2006 Antioxidant protective mechanisms against reactive oxygen species (ROS) generated by mitochondrial P450 systems in steroidogenic cells. *Drug Metabolism Reviews* **38** 171–196. (doi:10.1080/03602530600570040)
- Harman SM, Metter EJ, Tobin JD, Pearson J, Blackman MR & Baltimore Longitudinal Study of A 2001 Longitudinal effects of aging on serum total and free testosterone levels in healthy men. Baltimore Longitudinal Study of Aging. *Journal of Clinical Endocrinology and Metabolism* **86** 724–731. (doi:10.1210/jcem.86.2.7219)
- Hsueh AJ, Dufau ML & Catt KJ 1977 Gonadotropin-induced regulation of luteinizing hormone receptors and desensitization of testicular 3':5'-cyclic AMP and testosterone responses. *PNAS* **74** 592–595. (doi:10.1073/pnas.74.2.592)
- Huhtaniemi IT 2014 Andropause – lessons from the European male ageing study. *Annales d'Endocrinologie* **75** 128–131. (doi:10.1016/j.ando.2014.03.005)
- Hurst CH & Waxman DJ 2003 Activation of PPAR_α and PPAR_γ by environmental phthalate monoesters. *Toxicology Science* **74** 297–308. (doi:10.1093/toxsci/kfg145)
- Johnson L, Dickerson R, Safe SH, Nyberg CL, Lewis RP & Welsh TH Jr 1992 Reduced Leydig cell volume and function in adult rats exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin without a significant effect on spermatogenesis. *Toxicology* **76** 103–118. (doi:10.1016/0300-483X(92)90158-B)
- Kalender S, Uzun FG, Demir F, Uzunhisarcikli M & Aslanturk A 2013 Mercuric chloride-induced testicular toxicity in rats and the protective role of sodium selenite and vitamin E. *Food and Chemical Toxicology* **55** 456–462. (doi:10.1016/j.fct.2013.01.024)
- Kallen CB, Arakane F, Christenson LK, Watari H, Devoto L & Strauss JF 3rd 1998 Unveiling the mechanism of action and regulation of the steroidogenic acute regulatory protein. *Molecular and Cellular Endocrinology* **145** 39–45. (doi:10.1016/S0303-7207(98)00167-1)
- Keenan DM, Takahashi PY, Liu PY, Roebuck PD, Nehra AX, Iranmanesh A & Veldhuis JD 2006 An ensemble model of the male gonadal axis: illustrative application in aging men. *Endocrinology* **147** 2817–2828. (doi:10.1210/en.2005-1356)
- Kim JY, Han EH, Kim HG, Oh KN, Kim SK, Lee KY & Jeong HG 2010 Bisphenol A-induced aromatase activation is mediated by cyclooxygenase-2 up-regulation in rat testicular Leydig cells. *Toxicology Letter* **193** 200–208. (doi:10.1016/j.toxlet.2010.01.011)
- Kokilavani P, Suriyakalaa U, Elumalai P, Abirami B, Ramachandran R, Sankarganesh A & Achiraman S 2014 Antioxidant mediated ameliorative steroidogenesis by *Commelina benghalensis* L and *Cissus quadrangularis* L. against quinalphos induced male reproductive toxicity. *Pesticide Biochemistry and Physiology* **109** 18–33. (doi:10.1016/j.pestbp.2014.01.002)
- Kumar V, Balomajumder C & Roy P 2008 Disruption of LH-induced testosterone biosynthesis in testicular Leydig cells by triclosan: probable mechanism of action. *Toxicology* **250** 124–131. (doi:10.1016/j.tox.2008.06.012)
- Kumar A, Rani L & Dhole B 2014 Role of oxygen in the regulation of Leydig tumor derived MA-10 cell steroid production: the effect of cobalt chloride. *Systems Biology in Reproductive Medicine* **60** 112–118. (doi:10.3109/19396368.2013.861034)
- Lephart ED 2016 Skin aging and oxidative stress: equal's anti-aging effects via biochemical and molecular mechanisms. *Ageing Research Reviews* **31** 36–54. (doi:10.1016/j.arr.2016.08.001)
- Li W & Kong AN 2009 Molecular mechanisms of Nrf2-mediated antioxidant response. *Molecular Carcinogenesis* **48** 91–104. (doi:10.1002/mc.20465)
- Li W, Wu ZQ, Zhang S, Cao R, Zhao J, Sun ZJ & Zou W 2016 Augmented expression of gamma-glutamyl transferase 5 (GGT5) impairs testicular steroidogenesis by deregulating local oxidative stress. *Cell and Tissue Research* **366** 467–481. (doi:10.1007/s00441-016-2458-y)
- Liao C, Reaven E & Azhar S 1993 Age-related decline in the steroidogenic capacity of isolated rat Leydig cells: a defect in cholesterol mobilization and processing. *Journal of Steroid Biochemistry and Molecular Biology* **46** 39–47. (doi:10.1016/0960-0760(93)90207-D)
- Lin D, Sugawara T, Strauss JF 3rd, Clark BJ, Stocco DM, Saenger P, Rogol A & Miller WL 1995 Role of steroidogenic acute regulatory protein in adrenal and gonadal steroidogenesis. *Science* **267** 1828–1831. (doi:10.1126/science.7892608)
- Liu J, Rone MB & Papadopoulos V 2006 Protein-protein interactions mediate mitochondrial cholesterol transport and steroid biosynthesis. *Journal of Biological Chemistry* **281** 38879–38893. (doi:10.1074/jbc.M608820200)
- Liu J, Qu W & Kadiiska MB 2009 Role of oxidative stress in cadmium toxicity and carcinogenesis. *Toxicology and Applied Pharmacology* **238** 209–214. (doi:10.1016/j.taap.2009.01.029)
- Liu Q, Gu JH, Yuan Y, Liu XZ, Wang YJ, Wang HD, Liu ZP, Wang ZY & Bian JC 2013 Effect of cadmium on rat Leydig cell testosterone production and DNA integrity in vitro. *Biomedical and Environmental Science* **26** 769–773. (doi:10.3967/0895-3988.2013.09.009)
- Luo L, Chen H & Zirkin BR 1996 Are Leydig cell steroidogenic enzymes differentially regulated with aging? *Journal of Andrology* **17** 509–515. (doi:10.1002/j.1939-4640.1996.tb01827.x)
- Luo L, Chen H & Zirkin BR 2001 Leydig cell aging: steroidogenic acute regulatory protein (StAR) and cholesterol side-chain cleavage enzyme. *Journal of Andrology* **22** 149–156. (doi:10.1002/j.1939-4640.2001.tb02165.x)
- Luo L, Chen H & Zirkin BR 2005 Temporal relationships among testosterone production, steroidogenic acute regulatory protein (StAR), and P450 side-chain cleavage enzyme (P450_{sc}) during Leydig cell aging. *Journal of Andrology* **26** 25–31. (doi:10.1002/j.1939-4640.2005.tb02868.x)
- Luo L, Chen H, Trush MA, Show MD, Anway MD & Zirkin BR 2006 Aging and the brown Norway rat Leydig cell antioxidant defense system. *Journal of Andrology* **27** 240–247. (doi:10.2164/jandrol.05075)
- Martinez-Arguelles DB, Campioli E, Culty M, Zirkin BR & Papadopoulos V 2013 Fetal origin of endocrine dysfunction in the adult: the phthalate model. *Journal of Steroids Biochemistry and Molecular Biology* **137** 5–17. (doi:10.1016/j.jsbmb.2013.01.007)
- Mathur PP & D'Cruz SC 2011 The effect of environmental contaminants on testicular function. *Asian Journal of Andrology* **13** 585–591. (doi:10.1038/aja.2011.40)
- Midzak A, Akula N, Lecanu L & Papadopoulos V 2011 Novel androstenetriol interacts with the mitochondrial translocator protein and controls steroidogenesis. *Journal of Biological Chemistry* **286** 9875–9887. (doi:10.1074/jbc.M110.203216)
- Midzak A, Zirkin B & Papadopoulos V 2015 Translocator protein: pharmacology and steroidogenesis. *Biochemical Society Transactions* **43** 572–578. (doi:10.1042/BST20150061)
- Miller WL 2013 Steroid hormone synthesis in mitochondria. *Molecular and Cellular Endocrinology* **379** 62–73. (doi:10.1016/j.mce.2013.04.014)
- Mulligan T, Iranmanesh A & Veldhuis JD 2001 Pulsatile iv infusion of recombinant human LH in leuprolide-suppressed men unmasks impoverished Leydig-cell secretory responsiveness to midphysiological LH drive in the aging male. *Journal of Clinical Endocrinology and Metabolism* **86** 5547–5553. (doi:10.1210/jcem.86.11.8004)
- Murugesan P, Muthusamy T, Balasubramanian K & Arunakaran J 2005 Studies on the protective role of vitamin C and E against polychlorinated biphenyl (Aroclor 1254) – induced oxidative damage in Leydig cells. *Free Radical Research* **39** 1259–1272. (doi:10.1080/10715760500308154)
- Murugesan P, Muthusamy T, Balasubramanian K & Arunakaran J 2008 Polychlorinated biphenyl (Aroclor 1254) inhibits testosterone biosynthesis and antioxidant enzymes in cultured rat Leydig cells. *Reproductive Toxicology* **25** 447–454. (doi:10.1016/j.reprotox.2008.04.003)
- O'Brien ML, Spear BT & Glauert HP 2005 Role of oxidative stress in peroxisome proliferator-mediated carcinogenesis. *Critical Reviews in Toxicology* **35** 61–88. (doi:10.1080/10408440590905957)

- Ojo OO, Bhaduria S & Rath SK 2013 Dose-dependent adverse effects of salinomycin on male reproductive organs and fertility in mice. *PLoS ONE* **8** e69086. (doi:10.1371/journal.pone.0069086)
- Pandya C, Pillai P, Nampoothiri LP, Bhatt N, Gupta S & Gupta S 2012 Effect of lead and cadmium co-exposure on testicular steroid metabolism and antioxidant system of adult male rats. *Andrologia* **44** 813–822. (doi:10.1111/j.1439-0272.2010.01137.x)
- Papadopoulos V, Liu J & Culty M 2007 Is there a mitochondrial signaling complex facilitating cholesterol import? *Molecular and Cellular Endocrinology* **265–266** 59–64. (doi:10.1016/j.mce.2006.12.004)
- Papadopoulos V, Aghazadeh Y, Fan J, Campioli E, Zirkin B & Midzak A 2015 Translocator protein-mediated pharmacology of cholesterol transport and steroidogenesis. *Molecular and Cellular Endocrinology* **408** 90–98. (doi:10.1016/j.mce.2015.03.014)
- Payne AH & Hales DB 2004 Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine Reviews* **25** 947–970. (doi:10.1210/er.2003-0030)
- Ponting CP & Aravind L 1999 START: a lipid-binding domain in StAR, HD-ZIP and signalling proteins. *Trends in Biochemical Science* **24** 130–132. (doi:10.1016/S0968-0004(99)01362-6)
- Pogrmic K, Fa S, Dakic V, Kaisarevic S & Kovacevic R 2009 Atrazine oral exposure of peripubertal male rats downregulates steroidogenesis gene expression in Leydig cells. *Toxicology Science* **111** 189–197. (doi:10.1093/toxsci/kfp135)
- Rao MV & Sharma PSN 2001 Protective effect of vitamin E against mercuric chloride reproductive toxicity in male mice. *Reproductive Toxicology* **15** 705–712. (doi:10.1016/S0890-6238(01)00183-6)
- Rebrin I, Kamzalov S & Sohal RS 2003 Effects of age and caloric restriction on glutathione redox state in mice. *Free Radical Biology and Medicine* **35** 626–635. (doi:10.1016/S0891-5849(03)00388-5)
- Ronco AM, Valdes K, Marcus D & Llanos M 2001 The mechanism for lindane-induced inhibition of steroidogenesis in cultured rat Leydig cells. *Toxicology* **159** 99–106. (doi:10.1016/S0300-483X(00)00414-5)
- Sai L, Li X, Liu Y, Guo Q, Xie L, Yu G, Bo C, Zhang Z & Li L 2014 Effects of chlorpyrifos on reproductive toxicology of male rats. *Environmental Toxicology* **29** 1083–1088. (doi:10.1002/tox.21838)
- Saradha B, Vaithinathan S & Mathur PP 2008 Single exposure to low dose of lindane causes transient decrease in testicular steroidogenesis in adult male Wistar rats. *Toxicology* **244** 190–197. (doi:10.1016/j.tox.2007.11.011)
- Sasaki G, Ishii T, Jeyasuria P, Jo Y, Bahat A, Orly J, Hasegawa T & Parker KL 2008 Complex role of the mitochondrial targeting signal in the function of steroidogenic acute regulatory protein revealed by bacterial artificial chromosome transgenesis in vivo. *Molecular Endocrinology* **22** 951–964. (doi:10.1210/me.2007-0493)
- Shen WJ, Patel S, Natu V, Hong R, Wang J, Azhar S & Kraemer FB 2003 Interaction of hormone-sensitive lipase with steroidogenic acute regulatory protein: facilitation of cholesterol transfer in adrenal. *Journal of Biological Chemistry* **278** 43870–43876. (doi:10.1074/jbc.M303934200)
- Shi Z, Feng Y, Wang J, Zhang H, Ding L & Dai J 2010 Perfluorododecanoic acid-induced steroidogenic inhibition is associated with steroidogenic acute regulatory protein and reactive oxygen species in cAMP-stimulated Leydig cells. *Toxicology Science* **114** 285–294. (doi:10.1093/toxsci/kfq014)
- Sokanovic SJ, Janjic MM, Stojkov NJ, Baburski AZ, Bjelic MM, Andric SA & Kostic TS 2014 Age related changes of cAMP and MAPK signaling in Leydig cells of Wistar rats. *Experimental Gerontology* **58** 19–29. (doi:10.1016/j.exger.2014.07.004)
- Stocco DM, Zhao AH, Tu LN, Morohaku K & Selvaraj V 2017 A brief history of the search for the protein(s) involved in the acute regulation of steroidogenesis. *Molecular and Cellular Endocrinology* **441** 7–16. (doi:10.1016/j.mce.2016.07.036)
- Surampudi PN, Wang C & Swerdloff R 2012 Hypogonadism in the aging male diagnosis, potential benefits, and risks of testosterone replacement therapy. *International Journal of Endocrinology* **2012** 625434. (doi:10.1155/2012/625434)
- Tai P & Ascoli M 2011 Reactive oxygen species (ROS) play a critical role in the cAMP-induced activation of Ras and the phosphorylation of ERK1/2 in Leydig cells. *Molecular Endocrinology* **25** 885–893. (doi:10.1210/me.2010-0489)
- Tu LN, Morohaku K, Manna PR, Pelton SH, Butler WR, Stocco DM & Selvaraj V 2014 Peripheral benzodiazepine receptor/translocator protein global knock-out mice are viable with no effects on steroid hormone biosynthesis. *Journal of Biology Chemistry* **289** 27444–27454. (doi:10.1074/jbc.M114.578286)
- Veldhuis JD, Veldhuis NJ, Keenan DM & Iranmanesh A 2005 Age diminishes the testicular steroidogenic response to repeated intravenous pulses of recombinant human LH during acute GnRH-receptor blockade in healthy men. *American Journal of Physiology Endocrinology and Metabolism* **288** E775–E781. (doi:10.1152/ajpendo.00410.2004)
- Veldhuis JD, Liu PY, Keenan DM & Takahashi PY 2012 Older men exhibit reduced efficacy of and heightened potency downregulation by intravenous pulses of recombinant human LH: a study in 92 healthy men. *American Journal of Physiology Endocrinology and Metabolism* **302** E117–E122. (doi:10.1152/ajpendo.00450.2011)
- Venugopal S, Martinez-Arguelles DB, Chebbi S, Hullin-Matsuda F, Kobayashi T & Papadopoulos V 2016 Plasma membrane origin of the steroidogenic pool of cholesterol used in hormone-induced acute steroid formation in Leydig cells. *Journal of Biology Chemistry* **291** 26109–26125. (doi:10.1074/jbc.M116.740928)
- Wang C, Leung A & Sinha-Hikim AP 1993 Reproductive aging in the male brown-Norway rat: a model for the human. *Endocrinology* **133** 2773–2781. (doi:10.1210/endo.133.6.8243304)
- Wang X, Shen CL, Dyson MT, Eimerl S, Orly J, Hutson JC & Stocco DM 2005 Cytochrome P-450₁₇ regulation of the age-related decline in testosterone biosynthesis. *Endocrinology* **146** 4202–4208. (doi:10.1210/en.2005-0298)
- Zaidi SK, Shen WJ, Bittner S, Bittner A, McLean MP, Han JH, Davis RJ, Kraemer FB & Azhar S 2014 p38 MAPK regulates steroidogenesis through transcriptional repression of STAR gene. *Journal of Molecular Endocrinology* **53** 1–16. (doi:10.1530/JME-13-0287)
- Zhang SY, Ito Y, Yamanoshita O, Yanagiba Y, Kobayashi M, Taya K, Li C, Okamura A, Miyata M, Ueyama J *et al.* 2007 Permethrin may disrupt testosterone biosynthesis via mitochondrial membrane damage of Leydig cells in adult male mouse. *Endocrinology* **148** 3941–3949. (doi:10.1210/en.2006-1497)
- Zhang W, Shen XY, Zhang WW, Chen H, Xu WP & Wei W 2017 The effects of Di 2-Ethyl Hexyl Phthalate (DEHP) on cellular lipid accumulation in HepG2 cells and its potential mechanisms in the molecular level. *Toxicology Mechanisms and Methods* **27** 245–252. (doi:10.1080/15376516.2016.1273427)
- Zhao Y, Ao H, Chen L, Sottas CM, Ge RS, Li L & Zhang Y 2012 Mono-(2-ethylhexyl) phthalate affects the steroidogenesis in rat Leydig cells through provoking ROS perturbation. *Toxicology In Vitro* **26** 950–955. (doi:10.1016/j.tiv.2012.04.003)
- Zhou L, Beattie MC, Lin CY, Liu J, Traore K, Papadopoulos V, Zirkin BR & Chen H 2013 Oxidative stress and phthalate-induced down-regulation of steroidogenesis in MA-10 Leydig cells. *Reproductive Toxicology* **42** 95–101. (doi:10.1016/j.reprotox.2013.07.025)
- Zirkin BR, Santulli R, Strandberg JD, Wright WW & Ewing LL 1993 Testicular steroidogenesis in the aging brown Norway rat. *Journal of Andrology* **14** 118–123. (doi:10.1002/j.1939-4640.1993.tb01663.x)
- Zwart AD, Urban RJ, Odell WD & Veldhuis JD 1996 Contrasts in the gonadotropin-releasing hormone dose-response relationships for luteinizing hormone, follicle-stimulating hormone and alpha-subunit release in young vs older men: appraisal with high-specificity immunoradiometric assay and deconvolution analysis. *European Journal of Endocrinology* **135** 399–406. (doi:10.1530/eje.0.1350399)

Received 2 February 2017

First decision 28 February 2017

Revised manuscript received 14 July 2017

Accepted 26 July 2017