Update on mammalian sperm capacitation: how much does the horse differ from other species?

Bart Leemans¹, Tom A E Stout^{2,3}, Catharina De Schauwer¹, Sonia Heras¹, Hilde Nelis¹, Maarten Hoogewijs¹, Ann Van Soom¹ and Bart M Gadella^{3,4}

¹Department of Reproduction, Obstetrics and Herd Health, Faculty of Veterinary Medicine, Ghent University, Ghent, Belgium, ²Department of Equine Sciences, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands, ³Department of Farm Animal Health, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands and ⁴Department of Biochemistry and Cell Biology, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands

Correspondence should be addressed to B Leemans; Email: baleeman.leemans@ugent.be

Abstract

In contrast to various other mammalian species, conventional *in vitro* fertilization (IVF) with horse gametes is not reliably successful. In particular, stallion spermatozoa fails to penetrate the zona pellucida, most likely due to incomplete activation of stallion spermatozoa (capacitation) under *in vitro* conditions. In other mammalian species, specific capacitation triggers have been described; unfortunately, none of these is able to induce full capacitation in stallion spermatozoa. Nevertheless, knowledge of capacitation pathways and their molecular triggers might improve our understanding of capacitation-related events observed in stallion sperm. When sperm cells are exposed to appropriate capacitation triggers, several molecular and biochemical changes should be induced in the sperm plasma membrane and cytoplasm. At the level of the sperm plasma membrane, (1) an increase in membrane fluidity, (2) cholesterol depletion and (3) lipid raft aggregation should occur consecutively; the cytoplasmic changes consist of protein tyrosine phosphorylation and elevated pH, cAMP and Ca²⁺ concentrations. These capacitation-related events enable the switch from progressive to hyperactivated motility of the sperm cells, and the induction of the acrosome reaction. These final capacitation triggers are indispensable for sperm cells to migrate through the viscous oviductal environment, penetrate the cumulus cells and zona pellucida and, finally, fuse with the oolemma. This review will focus on molecular aspects of sperm capacitation and known triggers in various mammalian species. Similarities and differences with the horse will be highlighted to improve our understanding of equine sperm capacitation/fertilizing events.

Reproduction (2019) **157** R181–R197

Introduction

Conventional *in vitro* fertilization (IVF), that is co-incubation of mature oocytes with capacitated sperm in a petri dish, is a standard procedure for producing embryos *in vitro* in a range of species, including man (Stephoe & Edwards 1978), the cow (Perry 2013), the pig and various laboratory animals (Galli *et al.* 2003, Betteridge 2006). By contrast, conventional IVF has to date been very poorly successful in the horse. Equine IVF is hampered by the inability of stallion spermatozoa to penetrate the zona pellucida *in vitro* (Tremoleda *et al.* 2003). In theory, this failure could be attributed to deficiencies in either *in vitro* prepared sperm and/or *in vitro* matured oocytes. Given that *in vitro* matured oocytes placed in the oviduct can be fertilized *in vivo* (Hinrichs *et al.* 2002), while *in vivo* matured oocytes harvested from pre-ovulatory follicles cannot be fertilized *in vitro* (Palmer *et al.* 1991), the inability of sperm cells to penetrate the zona pellucida *ex vivo* is most likely

the result of deficient capacitation, that is activation of spermatozoa (Tremoleda *et al.* 2003). Although current equine IVF conditions do support the binding of stallion sperm to the oocyte's zona pellucida, this does not result in subsequent initiation of the acrosome reaction mandatory to penetrate the oocyte (Tremoleda *et al.* 2003). In contrast to sperm cells from other mammalian species, stallion sperm incubated under typical IVF conditions, that is in medium containing HCO₃⁻, Ca²⁺ and albumin, also fail to display other capacitation-related characteristics, such as hyperactivated motility, increased plasma membrane fluidity and protein tyrosine phosphorylation (Tremoleda *et al.* 2003) (Fig. 1).

Capacitation is the maturation process that sperm cells undergo after ejaculation, during their passage through the female genital tract (Suarez 2002). These biochemical changes enable the sperm to bind to and penetrate the zona pellucida and subsequently fuse with the oocyte (Fig. 1) (Yanagimachi 1994). The sperm

Figure 1 Pathways involved in sperm capacitation, with emphasis on the horse. When spermatozoa undergo capacitation, the following changes occur: (1) increase in membrane fluidity, (2) cholesterol depletion, (3) aggregation of lipid raft receptors, (4) protein tyrosine phosphorylation, (5) hyperactivated motility and (6) the acrosome reaction. PTK, protein tyrosine kinase; ROS, reactive oxygen species; ZP3, ZP protein 3; ZP3-receptor, ZP protein 3-receptor; +, activation; –, inhibition.

plasma membrane fuses with the outer acrosomal membrane, resulting in the exocytotic event known as the acrosome reaction that involves the release of lytic enzymes which locally dissolve the zona pellucida. The sperm cells also acquire hyperactivated motility in order to ‘drill’ through the zona pellucida (Yanagimachi 1994). Equids are one of the exceptions, in which capacitation does not appear to progress as described above. Using an oviduct epithelial explant model, we identified physiological triggering of three important capacitation parameters, namely increased intracellular pH (Leemans *et al.* 2014), protein tyrosine phosphorylation in the sperm tail (Leemans *et al.* 2014, 2015b) and hyperactivated motility (Leemans *et al.* 2015b). However, this approach was not sufficient to induce the acrosome reaction and fertilization during co-incubation with oocytes. Hence, our understanding of sperm activation in the oviduct, the ability of these spermatozoa to achieve fertilizing competence and, in particular, the intracellular pathways involved in this process needs to improve. This review focuses on what is currently known about capacitation pathways in sperm cells of non-equine species, the provision of fuel to support the highly energy-dependent capacitation events and how much these events differ to capacitation-related events observed in stallion spermatozoa.

Energy metabolism in stallion spermatozoa to support energy-demanding, fertility-related processes such as capacitation

Sperm capacitation and other fertility-related pathways require large quantities of energy. To fuel these crucial capacitation events, individual sperm cells must generate relatively large amounts of ATP. In various mammalian species including man and the mouse, sperm motility and the maintenance of membrane integrity rely predominantly on ATP generated by non-aerobic glucose metabolism via glycolysis (Mukai & Okuno 2004, Storey 2008, du Plessis *et al.* 2015), as demonstrated by the fact that mitochondrial oxidative chain uncouplers did not affect sperm motility or membrane integrity in sperm suspended in glucose-containing media. By contrast, the proportion of membrane intact and motile stallion sperm, and the velocity of that motion, in glucose-enriched media were all significantly reduced after exposure to mitochondrial uncouplers (Gibb *et al.* 2014, Davila *et al.* 2016). Inhibiting complex IV or ATP synthase, two proteins on the inner mitochondrial membrane critical to electron flow during oxidative phosphorylation, had similar effects on stallion sperm (Davila *et al.* 2016). Moreover, subsequent uncoupling of mitochondria or inhibition of mitochondrial respiration led to a massive

drop in cytoplasmic ATP concentrations in stallion spermatozoa. Overall, this indicates that maintenance of stallion sperm membrane integrity and motility, two important requirements for fertility, are highly dependent on aerobic production of ATP (Gibb *et al.* 2014, Plaza Davila *et al.* 2015). As a consequence of the requirement for stallion sperm to generate mitochondrial ATP aerobically, increased cytoplasmic concentrations of reactive oxygen species (ROS), such as superoxide anions and their reduced form, hydrogen peroxide, are observed during stallion sperm incubation and activation. In turn, this cytoplasmic ROS accumulation leads to increased production of metabolic by-products, and the combination of accumulating ROS and cytotoxic lipid aldehydes (acrolein, 4-hydroxynonenal and malondialdehyde) ultimately leads to massive lipid peroxidation and DNA damage and thereby accelerated cell death (Gibb *et al.* 2016). To counteract the adverse effect of endogenous ROS production, and enable ROS homeostasis (Jones 2006) or defense against oxidative stress, stallion seminal plasma contains relatively high concentrations of antioxidants such as catalase (Ortega Ferrusola *et al.* 2009), superoxide dismutase (Ball *et al.* 2000, Ortega Ferrusola *et al.* 2009), glutathione (Hemachand & Shaha 2003, Jones 2006) and ergothioneine (Mann 1975). Moreover, mitochondrial aldehyde dehydrogenase 2 has been identified in stallion sperm cells and proposed to serve as a detoxifier of aldehydes generated either endogenously or exogenously (Gibb *et al.* 2016). However, ROS defense mechanisms will eventually become exhausted and are not thus completely protective against sperm cell degeneration.

In this aspect, an interesting observation has been made with respect to stallion sperm oxidative stress and fertility. Spermatozoa from matings that did result in conception (and therefore considered 'more fertile') had lower percentages of membrane intact and higher percentages of cells displaying ROS-induced damage than spermatozoa from matings that did not result in conception (Gibb *et al.* 2014). It has been proposed that this can be explained by the fact that fertile stallions produce metabolically more active sperm cells that exhibit higher levels of oxidative phosphorylation. This will result in higher concentrations of ROS, leading to increased lipid peroxidation and, in turn, impaired sperm motility and membrane integrity in more fertile stallions. This has led to the suggestion that spermatozoa from highly fertile stallions 'live fast and die young'. It should also be borne in mind that a modest degree of cytoplasmic ROS production plays an important physiological role in sperm capacitation. These aspects are discussed below.

How to trigger capacitation *in vitro*?

In vivo, capacitation is initiated when a sperm cell is exposed to the environment within the female genital

tract, that is the uterus and oviduct, close to the time of ovulation (reviewed by Leemans *et al.* (2016a)); capacitation events can be mimicked *in vitro* by incubating sperm cells in medium containing HCO_3^- , Ca^{2+} and albumin, after performing density gradient centrifugation (e.g. Percoll) to separate the sperm cells from the seminal plasma. HCO_3^- , Ca^{2+} and albumin are three capacitation factors known to induce the sperm changes required for the acquisition of fertilizing potential across many species (mice: Visconti *et al.* 1995a,b; hamster: Visconti *et al.* 1999b; man: Osheroff *et al.* 1999; pig: Flesch & Gadella 2000). In addition, species-specific factors have been identified. In cattle, for example, heparin-like molecules such as glycosaminoglycans are an essential capacitation trigger (Parrish *et al.* 1988). Unfortunately, the species-specific capacitation triggers required to establish a repeatable, effective equine IVF system have yet to be identified. In various studies of stallion sperm capacitation, 'capacitating medium' has included the three general capacitation triggers (HCO_3^- , Ca^{2+} and albumin) even though it is known that full capacitation is not accomplished; non-capacitating medium, lacking these three triggers, is used as control medium (McPartlin *et al.* 2008, Leemans *et al.* 2016b). In Table 1, an overview of molecules frequently used as capacitation triggers in mammals, including the horse, is shown.

Capacitation events

Although capacitation was described for the first time in 1951 (Chang 1951, Austin 1952), the process is still not completely understood. It is known that capacitation involves a series of changes including (1) the removal of seminal plasma and surface-adhered decapacitation factors from the sperm plasma membrane, (2) reorganization of the sperm plasma membrane and (3) activation of intracellular pathways (Fig. 1).

After the removal of seminal plasma, a rapid increase in membrane fluidity (within 10 min) is induced by an increasing intracellular HCO_3^- concentration and the activation of second messenger systems, including soluble adenylyl cyclase and a rise in intracellular Ca^{2+} (pig: Flesch & Gadella 2000, Gadella *et al.* 2008; horse: Rath *et al.* 2003). The activation of soluble adenylyl cyclase and the concomitant production of cAMP results in depletion of cholesterol from the sperm plasma membrane via a cholesterol acceptor such as albumin (>1 h), which is followed by a slower series of functional membrane changes in which lipid ordered microdomains aggregate at the apical ridge of the sperm head (>1 h) (pig: van Gestel *et al.* 2005a). These microdomains contain functional zona pellucida-binding protein complexes (pig: van Gestel *et al.* 2007) and the soluble *N*-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) proteins, which play an important role in the induction of the acrosome

Table 1 Overview of capacitation triggers and their *in vitro* capacitation effect in different mammalian species.

Capacitating trigger	Capacitation effect	Species	Reference
Ca ²⁺	Membrane fluidity	Mouse	Visconti <i>et al.</i> (1995a,b)
	Protein tyrosine phosphorylation	Man	Osheroff <i>et al.</i> (1999)
	Hyperactivated motility	Pig	Flesch & Gadella (2000)
	Acrosome reaction	Cow	Byrd (1981), Breininger <i>et al.</i> (2010)
		Horse	McPartlin <i>et al.</i> (2008)
HCO ₃ ⁻	Membrane fluidity	Mouse	Visconti <i>et al.</i> (1995a,b)
	Protein tyrosine phosphorylation	Hamster	Visconti <i>et al.</i> (1999b)
	Hyperactivated motility	Man	Osheroff <i>et al.</i> (1999)
	Acrosome reaction	Pig	Flesch & Gadella (2000)
		Cow	Breininger <i>et al.</i> (2010)
		Horse	Rathi <i>et al.</i> (2003), McPartlin <i>et al.</i> (2008)
		Mouse	Visconti <i>et al.</i> (1995a,b)
Albumin	Cholesterol depletion	Pig	Flesch & Gadella (2000)
		Cow	Byrd (1981)
		Horse	McPartlin <i>et al.</i> (2008)
		Mouse	Visconti <i>et al.</i> (1999a)
Methyl β -cyclodextrin	Cholesterol extraction	Pig	van Gestel <i>et al.</i> (2005b)
		Horse	Bromfield <i>et al.</i> (2014)
		Cow	Parrish <i>et al.</i> (1988), Gualtieri <i>et al.</i> (2005), Breininger <i>et al.</i> (2010)
Heparin	Membrane fluidity		
	Hyperactivated motility		
	Protein tyrosine phosphorylation		
	Acrosome reaction		
Progesterone	Acrosome reaction	Horse	Cheng <i>et al.</i> (1998a,b)
	Hyperactivated motility	Man	Lishko <i>et al.</i> (2011)
Ca ²⁺ ionophore A23187	Acrosome reaction	Mouse	Tateno <i>et al.</i> (2013)
		Man	Bielfeld <i>et al.</i> (1994), Liu <i>et al.</i> (2011)
		Pig	
		Cow	Birck <i>et al.</i> (2009)
		Horse	Fraser <i>et al.</i> (1995)
		Mice	Balao da Silva <i>et al.</i> (2013), Tateno <i>et al.</i> (2013)
		Horse	Graham (1996)
Lysophospha-tidylcholine c-AMP and caffeine	Hyperactivated motility		
	Acrosome reaction		
	Protein tyrosine phosphorylation		
ROS	Hyperactivated motility		
	Protein tyrosine phosphorylation		
Alkaline medium pH	Protein tyrosine phosphorylation		
	Hyperactivated motility		
Procaine	Hyperactivated motility		

reaction (pig: Tsai *et al.* 2010, 2012). Simultaneously, the production of cAMP enables the activation of protein kinase A which is required to phosphorylate the tyrosine residues on sperm proteins (Rathi *et al.* 2003, Ijiri *et al.* 2012, Signorelli *et al.* 2012). In various species, this cAMP-dependent protein tyrosine phosphorylation, especially in the sperm tail, has been associated with the acquisition of hyperactivated sperm motility and is considered a marker for some essential elements of the capacitation process (mouse: Visconti *et al.* 1995a,b; pig: Flesch *et al.* 1999, Harayama *et al.* 2012).

Plasma membrane changes

Spermatozoa have a highly polarized morphology with a heterogenic surface differentiated into at least four surface membrane domains: the apical ridge and the pre-equatorial, equatorial and post-equatorial surface

areas, respectively. Each of these regions plays a specific role in fertilization: (1) the apical ridge facilitates zona pellucida-binding (rabbit: O'Rand & Fisher 1987), (2) the apical ridge, together with the pre-equatorial surface area, is involved in the acrosome reaction while (3) the equatorial surface area initiates binding to the oolemma and the subsequent fertilization fusion. These events occur in the region of the sperm head where the sperm plasma membrane, including the lipid rafts (microdomains), covers the acrosome and plays an essential role in the dynamics of sperm capacitation (Gadella *et al.* 2008).

Cholesterol depletion

A redistribution of phospholipids in the surface of the sperm head is an essential step in the capacitation process. HCO₃⁻/soluble adenylyl cyclase/cAMP mediate

an increase in membrane fluidity, detected as enhanced merocyanine 540 binding (Gadella & Harrison 2000, 2002), that enables the redistribution of cholesterol from the equatorial area to the apical part of the sperm plasma membrane, while seminolipids move retrogradely (pig: Gadella *et al.* 1994, 1995). Both molecules are considered to be plasma membrane stabilizers, which prevent membrane fusion prior to capacitation. The HCO_3^- -mediated repacking of the sperm surface lipids is also essential in enabling the efflux of cholesterol which further increases plasma membrane fluidity by allowing the aggregation of lipid rafts, another essential step in the reorganization of the plasma membrane.

In vitro, the extraction of cholesterol from the sperm plasma membrane can be achieved by the oxidation of membrane sterols (Boerke *et al.* 2013), or active cholesterol transporters that deliver free cholesterol to the hydrophobic pocket of albumin (Flesch *et al.* 2001). Brouwers *et al.* (bull: 2011) showed that the production of ROS is an essential step in oxysterol formation. ROS production at low levels during sperm capacitation was first demonstrated in 1993, and indicates that sperm capacitation should be considered an oxidative process dependent on the active generation of ROS (man: de Lamirande & Gagnon 1993). This process may also help to regulate protein tyrosine phosphorylation, another important capacitation marker, by increasing cAMP production (Aitken *et al.* 2004, Aitken 2011) and suppressing tyrosine phosphatase activity (Takakura *et al.* 1999). Since oxysterols are more hydrophilic than intact cholesterol, they can more easily move through the plasma membrane to bind to sterol acceptor molecules like albumin. Indeed, the presence of bovine serum albumin in capacitating medium is essential because of its unique ability to scavenge hydrophilic oxidation products and facilitate cholesterol extraction (Boerke *et al.* 2008). An alternative macromolecule not of animal origin (important in terms of risk of disease transmission) is methyl β -cyclodextrin. Whereas albumin extracts only 20% of the cholesterol from the non-raft sperm plasma membrane fraction, methyl β -cyclodextrin extracts 50% of the cholesterol from the entire sperm plasma membrane, including the lipid rafts (pig: van Gestel *et al.* 2005b). However, methyl β -cyclodextrin might have a deteriorative effect on sperm cells, even in low concentrations. A deteriorative effect of methyl β -cyclodextrin on mouse oocytes has also been reported (Buschiazzi *et al.* 2013). In stallion sperm, however, standard capacitating conditions, that is inclusion of Ca^{2+} , HCO_3^- and bovine serum albumin (individual or in combination), does not facilitate cholesterol removal from the sperm plasma membrane. Indeed, while HCO_3^- induces an increase in ROS, this is abolished by the addition of Ca^{2+} or albumin (Macias Garcia *et al.* 2015). These observations suggest a species-specific difference in the regulation of plasma membrane changes during stallion sperm capacitation.

Aggregation of lipid rafts

As demonstrated in pig spermatozoa, increased membrane fluidity and cholesterol depletion are followed by a redistribution of the laterally segregated molecules, that is lipid ordered microdomains aggregate at the apical ridge area of the sperm head (Boerke *et al.* 2008). The proteins and lipids which make up these microdomains also show a capacitation-dependent change to include higher proportions of cholesterol, sphingomyelin, gangliosides and phospholipids with saturated long-chain acyl chains and lipid-modified proteins such as GPI-anchored proteins (Simons & Toomre 2000, van Gestel *et al.* 2005a). Moreover, caveolin-1 and flotillin-1 are lipid raft-specific markers (van Gestel *et al.* 2005a) that accumulate in these microdomains along with functional zona pellucida-binding protein complexes. In man, angiotensin-converting enzyme and protein disulfide isomerase A6 were identified in the microdomains as proteins able to interact with heat shock protein A2, which plays a pivotal role in the remodeling of the sperm surface during capacitation and the recognition of human oocytes (Bromfield *et al.* 2016). In pigs, it has been shown that isoforms of AQN-3 (spermadhesin), P47 (porcine homologue of SED-1), fertilin β and peroxiredoxin 5 are key proteins regulating the primary binding between capacitated spermatozoa and the zona pellucida (van Gestel *et al.* 2007). However, it is not known whether these proteins reside in microdomains.

Hyperactivated motility

General sperm motility

A specific cytoplasmic pH is crucial to allow sperm motility. For de-membranated sperm, maximal percentages of motile sperm were recorded between pH 7.0 and 8.1 for bull (Ho *et al.* 2002), pH 7.8 for human (Giroux-Widemann *et al.* 1991) and between pH 7.5 and 8.0 for ram (Ishijima & Witman 1987) spermatozoa. For comparison, the maximum percentage of motile stallion spermatozoa was observed in medium at pH 7.0 (Loux *et al.* 2014). An appropriate cytoplasmic pH enhances symmetrical flagellar, and therefore, progressive sperm motility, which requires the activation of dynein ATPases on phosphorylated dynein molecules and their interaction with Ca^{2+} which, in turn, results in the sliding of the adjacent outer axonemal doublet microtubules (Shingyoji *et al.* 1977). When the doublets slide along one another, the sliding force is translated into a bend in the sperm tail (Tash 1989). The normal flagellar waveform requires an asynchronous phosphorylation and dephosphorylation of the dynein arms along the complete axonemal length (Wargo & Smith 2003). In mammals, sperm motility is mostly initiated and maintained by ATP, Ca^{2+} and HCO_3^- -driven cAMP-dependent phosphorylation of flagellar proteins

(Tash & Means 1983, White & Aitken 1989, Ho *et al.* 2002). Like HCO_3^- , Ca^{2+} also directly regulates soluble adenylyl cyclase, which results in cAMP generation and activation of protein kinase A (Fig. 1) (Liguori *et al.* 2004, Hess *et al.* 2005). An important downstream target of cAMP in the sperm flagellum is serine/threonine kinase protein kinase A (Visconti *et al.* 1997). The phosphorylation of serine/threonine activates protein kinase A which results in the downstream phosphorylation of tyrosine kinases whose targets are primarily located in the sperm tail (man: Leclerc *et al.* 1996; mouse: Si & Olds-Clarke 2000). On the other hand, dynein dephosphorylation is evoked by the calmodulin-dependent protein serine/threonine phosphatase, calcineurin, which is necessary to balance the cAMP-driven serine/threonine kinases in the sperm tail. The resulting net phosphorylation represents the sperm motility status well (Tash & Bracho 1994). If serine/threonine phosphatase activity is dominant, spermatozoa are mainly immotile whereas serine/threonine kinase activity correlates with increased motility (man and Rhesus monkey: Smith *et al.* 1996; bull: Vijayaraghavan *et al.* 1996).

Relationship between Ca^{2+} , pH and hyperactivated sperm motility?

Sperm cells must acquire hyperactivated motility to (1) detach from their oviduct epithelial cell binding and leave the oviductal reservoir (Suarez & Pacey 2006), (2) migrate through the viscous lumen of the oviduct (Suarez & Dai 1992) and (3) penetrate the cumulus matrix and zona pellucida of the mature oocyte in order to fuse with the oolemma (Stauss *et al.* 1995). In many species, hyperactivated motility is characterized by a highly asymmetrical and high-amplitude flagellar beating pattern that gives rise to a whip-like motion of the sperm tail which, *in vitro*, results in circular, figure of eight or zigzag swimming trajectories (Yanagimachi 1994, Ishijima *et al.* 2006). The onset and maintenance of hyperactivated motility is associated with an influx of Ca^{2+} into the cytosol of the sperm tail (Suarez *et al.* 1993, Suarez 2008). Ho *et al.* (2002) showed in de-membranated bull sperm that a symmetrical, progressive sperm movement was maintained when the intracellular Ca^{2+} concentration was ~50 nM. When hyperactivated motility was initiated, the intracellular Ca^{2+} concentration increased to 400 nM. By contrast, Loux *et al.* (2014) reported that the degree of hyperactivated motility was inversely related to the cytoplasmic Ca^{2+} concentration in stallion spermatozoa. Using a similar de-membranated sperm model, they demonstrated that an increasing Ca^{2+} concentration did not induce hyperactivated motility in stallion spermatozoa at any pH. Moreover, cytoplasmic alkalization induced both a hyperactivated motility response and cytoplasmic Ca^{2+} rise in intact sperm cells (Loux *et al.* 2013). Loux *et al.* (2013) observed an intracellular pH increase from 7.1

to 7.3–7.4 in hyperactivating conditions. A pH effect on the motility of de-membranated stallion spermatozoa was not apparent. However, an extremely low Ca^{2+} concentration (27 pM) was sufficient to maintain motility in de-membranated stallion sperm at pH 7.4 (Loux *et al.* 2014). Moreover, procaine which is considered a potent inducer of hyperactivated motility in intact stallion spermatozoa, acts independently to extracellular Ca^{2+} influx (Loux *et al.* 2013). Overall, a species-specific relationship between Ca^{2+} sensitivity and hyperactivated motility is apparent in the horse.

In vitro, Ca^{2+} ionophores such as A23187 or ionomycin can induce and maintain hyperactivated motility in mouse spermatozoa for several hours (Suarez *et al.* 1987, Tatenio *et al.* 2013). Besides the reliable induction of the acrosome reaction in stallion spermatozoa, Ca^{2+} ionophore A23187 exposure results in a complete loss of sperm motility and membrane integrity within 1 h (Christensen *et al.* 1996, Rathie *et al.* 2001). It is possible that these adverse effects of Ca^{2+} ionophore on sperm motility and membrane integrity are related to the fact that stallion sperm, in contrast to other mammals, are dependent on mitochondrial ATP (as discussed above). In this respect, excessive mitochondrial Ca^{2+} concentration can lead to mitochondrial failure and initiate cell death (Contreras *et al.* 2010). Moreover, Ca^{2+} ionophores uncouple oxidative phosphorylation by inhibiting mitochondrial ATPase activity (Humes & Weinberg 1980, Krumschnabel *et al.* 1999). As mentioned above, other inhibitors that uncouple oxidative phosphorylation adversely affect stallion sperm motility and membrane integrity (Gibb *et al.* 2014, Davila *et al.* 2016). This is because production of mitochondrial ATP to support stallion sperm motility and membrane integrity is severely compromised soon after exposure. This is a cause for concern with respect to a capacitation induction protocol, because sperm viability and motility would need to be conserved over a period of hours during *in vitro* fertilization. Other pharmacological agents such as caffeine (Ho & Suarez 2001b), thimerosal (Ho & Suarez 2001b, Marquez & Suarez 2004) and thapsigargin (Ho & Suarez 2001b, Ho & Suarez 2003) also trigger an intracellular Ca^{2+} rise and initiate asymmetrical flagellar beating in mouse spermatozoa. The effect of these molecules on the cytoplasmic Ca^{2+} concentration and hyperactivated motility have yet to be studied in the horse.

In mammals such as man, cattle and mice, CATSPER channels present on the principal piece of the sperm tail's need be activated to induce hyperactivated motility. There are four CATSPER genes coding for proteins that are structurally similar to subunits of conventional voltage-gated cation channels. When a mouse single CATSPER gene knock-out was created, males were infertile because the spermatozoa were not able to achieve hyperactivated motility (Carlson *et al.* 2005, Qi *et al.* 2007). The general trigger of these CATSPER

channels is alkaline depolarization evoked by a change in the oviductal ionic environment, which results in an elevated pH of the oviduct fluid (Carlson *et al.* 2003, Navarro *et al.* 2008). Contact between spermatozoa and the alkaline oviductal environment increases the intracellular pH and activates the CATSPER channels. In stallion sperm, CATSPER1 mRNA has been identified, and CATSPER1 protein was localized to the principal piece of the sperm tail. However, analysis of the equine CATSPER1 protein revealed species-specific differences in the structure in the pH sensor region (Loux *et al.* 2013). This indicated that, despite the presence of a CATSPER channel in stallion sperm, the relationship between hyperactivated motility and Ca^{2+} influx is likely to be weak.

A clear increase in oviduct pH and, as a consequence, in sperm cytoplasmic pH likely to activate the CATSPER channels, was observed in *Rhesus* monkeys approaching ovulation (7.1–7.3 to 7.5–7.8) (Maas *et al.* 1977). In the mouse, a Na^+ -dependent $\text{Cl}^-/\text{HCO}_3^-$ exchange controls the intracellular pH of the sperm (Zeng *et al.* 1996), while in man, an outflow of H^+ is initiated by activating a voltage-gated proton channel (Lishko *et al.* 2010). To maximize the subsequent Ca^{2+} entry through the CATSPER channels, both a pH-sensitive efflux of K^+ by KSPER (Navarro *et al.* 2007) and an activation of Cl^- channels by closing Na^+ channels might support capacitation-related sperm plasma membrane hyperpolarization (mouse: Hernandez-Gonzalez *et al.* 2007). Hyperpolarization also plays a central role in the acrosome reaction. In human spermatozoa, it has been demonstrated that CATSPER channels can be alternatively activated by progesterone and, to a lesser extent, prostaglandins (Lishko *et al.* 2011). Other Ca^{2+} metabolism-related pathways identified in the mammalian sperm tail, include the following: (1) a transient receptor potential Ca^{2+} channel, which may affect sperm motility and re-sequestration of Ca^{2+} into sperm stores (Castellano *et al.* 2003), (2) cyclic-nucleotide-gated Ca^{2+} channels (Wiesner *et al.* 1998) and (3) voltage-gated Ca^{2+} channels (Benoff 1998, Darszon *et al.* 2006). However, the role of the various Ca^{2+} regulatory pathways in the physiological activation of hyperactivated motility is unclear.

Besides the extracellular Ca^{2+} influx through the CATSPER channels, there are also intracellular Ca^{2+} stores, in the so-called redundant nuclear envelope (RNE) located at the base of the sperm tail, which are able to provide Ca^{2+} to the sperm cytoplasm (Ho & Suarez 2001b, 2003). Inositol 1,4,5-triphosphate (IP_3)-gated channels on the RNE membranes trigger Ca^{2+} release to the sperm cytoplasm whereas calreticulin, a Ca^{2+} binding protein, sequesters Ca^{2+} in the RNE (bull: Ho & Suarez 2001b, 2003; man: Naaby-Hansen *et al.* 2001). In the mouse, it has been shown that ryanodine receptors on the RNE membrane also play a role in intracellular Ca^{2+} release (Trevino *et al.* 1998).

As such, both Ca^{2+} sources contribute to initiation and maintenance of high Ca^{2+} concentrations during sperm hyperactivation (Ho & Suarez 2001b, 2003). More specifically, the Ca^{2+} -induced Ca^{2+} release from the RNE amplifies the intracellular Ca^{2+} rise resulting from CATSPER activation (Costello *et al.* 2009, Olson *et al.* 2010). Furthermore, this activation of CATSPER and IP_3 -gated channels is clearly involved in induction of the acrosome reaction (Quill *et al.* 2003, Ren *et al.* 2001). So far, a role of intracellular Ca^{2+} stores in the induction of hyperactivated motility in stallion spermatozoa is purely theoretical.

Hypermotility is induced by HCO_3^- and Ca^{2+} -driven pathways

So far, it has not been elucidated how sperm cells modify the beat of their tail, at the molecular level. It is known that the Ca^{2+} rise and the subsequent onset of hyperactivated motility changes the sliding of the microtubules (Lindemann & Lesich 2010). Furthermore, it has been demonstrated in primates (Mahony & Gwathmey 1999), pigs (Harayama *et al.* 2012) and rodents (Si & Okuno 1999) that hyperactivated motility is associated with a highly increased cAMP-dependent tyrosine phosphorylation of the flagellar proteins. The generator of tyrosine phosphorylation events, protein kinase A, is connected to the fibrous sheath of the sperm tail by A-kinase-anchoring proteins, and it has been suggested that these proteins play an important role in hyperactivated sperm motility (Hamster: Si 1999). Moreover, A-kinase-anchoring proteins could attach protein kinase A to specific subcellular regions in close proximity to motility-related targets in the axoneme (Carrera *et al.* 1994, Mandal *et al.* 1999). A-kinase-anchoring protein 3 and A-kinase-anchoring protein 4 play a central role in activating tyrosine kinases to cause extensive tyrosine phosphorylation of sperm tail proteins (hamster: Si 1999; man: Ficarro *et al.* 2003). This altered tail protein tyrosine phosphorylation status is required for the acquisition of hyperactivated sperm motility (hamster: Si 1999; man: Leclerc *et al.* 1996). Changes in A-kinase-anchoring protein-mediated protein targeting might also play an essential role in this process.

Independent of protein kinase A, Ca^{2+} is involved in pathways regulating sperm motility. Calmodulin, located in the principal piece of the sperm tail (Schlingmann *et al.* 2007), is a Ca^{2+} -binding protein essential to the Ca^{2+} -dependent modulation of mammalian sperm motility (Ho & Suarez 2001a, Ignatz & Suarez 2005). It has been suggested that this pathway progresses in parallel with cAMP/protein kinase A activity, although the two act independently (Litvin *et al.* 2003, Schlingmann *et al.* 2007). Binding of Ca^{2+} to calmodulin activates Ca^{2+} /calmodulin-dependent kinases (Marin-Briggiler *et al.* 2005) which phosphorylate a specific axonemal protein, resulting in hyperactivated motility

(Suarez 2008). Calmodulin kinases present in the flagellum of bull (Ignatz & Suarez 2005) and human (Marin-Briggiler *et al.* 2005) sperm were associated with hyperactivated motility. On the other hand, phosphatase activities are also directly involved in axonemal function because they reverse the latter effect by regulating dynein ATPase activities (Tash *et al.* 1988, Carrera *et al.* 1996) (Fig. 2).

Is hyperactivated motility linked to protein tyrosine phosphorylation during stallion sperm capacitation?

As indicated above, in various mammalian species it has been shown that hyperactivated motility is associated with a marked increase in cAMP/protein kinase A and Ca^{2+} /calmodulin-dependent kinase activity that induces tyrosine phosphorylation of the flagellar proteins that, in turn, alter the sliding of the microtubules within the sperm axoneme (mouse: Visconti *et al.* 1995a,b; pig: Flesch *et al.* 1999, Harayama *et al.* 2012). Surprisingly, hyperactivated motility and protein tyrosine phosphorylation can also occur independently (mouse: Olds-Clarke 1989; cow: Marquez & Suarez 2004). This latter observation certainly seems to be true for stallion sperm. For example, type 10 sAC induced protein tyrosine phosphorylation in stallion sperm, without initiating hyperactivated motility (McPartlin *et al.* 2011). An increased PKA activity and protein tyrosine phosphorylation without hyperactivation was also observed after incubating stallion spermatozoa *in vitro* with membrane-soluble cAMP analogues or a phosphodiesterase inhibitor (caffeine) (Pommer *et al.* 2003). ROS (Baumber *et al.* 2003) and modified-Whitten's capacitating medium with higher alkalinity

(pH=7.8–8.0) (Gonzalez-Fernandez *et al.* 2012, 2013) also induce protein tyrosine phosphorylation in the absence of, or only a mild, hyperactivated motility response. Similarly, binding of stallion sperm to late pre-ovulatory stage oviduct epithelium induces protein tyrosine phosphorylation without hyperactivated motility. This protein tyrosine phosphorylation was regulated primarily by a high pH within oviductal secretory cells (Leemans *et al.* 2014). When stallion sperm are incubated in capacitating conditions at pH 7.4, a Ca^{2+} -mediated inhibitory effect on protein tyrosine phosphorylation occurs due to the formation of a Ca^{2+} /calmodulin complex that supports sperm phosphatase activity (Gonzalez-Fernandez *et al.* 2012). Under alkaline-capacitating conditions (pH=7.8–8.0) however, Ca^{2+} /calmodulin-dependent kinases dominate and play a downstream role in protein kinase A-dependent protein tyrosine phosphorylation. As such, Ca^{2+} -induced protein tyrosine kinases activate protein tyrosine phosphorylation in stallion spermatozoa downstream of protein kinase A (Gonzalez-Fernandez *et al.* 2013).

Interestingly, after cryopreservation a premature induction of protein tyrosine phosphorylation is observed, commonly referred to as 'cryocapacitation' (Bailey *et al.* 2000, Watson 2000). The increased osmolality associated with cryopreservation may induce a ROS-dependent increase in protein tyrosine phosphorylation in stallion sperm (Aitken *et al.* 1983, Burnaugh *et al.* 2010). After cryopreservation, sperm cells display an increased intracellular Ca^{2+} concentration, increased ROS generation and a reduced antioxidant capacity. Recently, Ortega-Ferrusola *et al.* (2017) examined the effect of cryopreservation on both apoptosis and

Figure 2 Schematic representation of signaling pathways involved in the regulation of mammalian sperm (hyper)motility. Progressive motility is regulated by low-level activity of the AC/cAMP/PKA pathway, while hyperactivated motility is induced when the latter pathway is activated to a much higher level, in combination with the activation of the calmodulin kinase (CaMK) pathway. Image modified from Turner (2006).

premature capacitation-like changes simultaneously, to investigate the reduced longevity of frozen/thawed stallion sperm. In general, cryopreservation induced an increase in caspase 3 activity, translocation of phosphatidylserine from the inner to outer layer of the plasma membrane lipid bilayer, a reduction in mitochondrial membrane potential and an increased cytoplasmic Na^+ concentration, all of which indicate the occurrence of an apoptotic-like process in frozen/thawed stallion sperm cells. Additionally, a small subpopulation of cryopreserved sperm showed capacitation-like characteristics after thawing, such as increased membrane fluidity and protein tyrosine phosphorylation. To avoid the negative effects of cooling and cryopreservation on stallion sperm, Gibb *et al.* (2015) recently described a promising approach to store stallion spermatozoa at room temperature. Taking into account that stallion spermatozoa rely heavily on mitochondrial ATP production which generates ROS (as discussed previously), combined supplementation of sperm storage medium with 10 mM pyruvate and 50 mM L-carnitine was found to counteract the adverse effects of ROS production during room temperature storage. Much improved preservation of sperm total and progressive motility after 72 h incubation and significantly higher acetyl-L-carnitine concentrations were observed using this antioxidant-enriched sperm storage medium.

In contrast to other mammalian species, no intracellular Ca^{2+} rise and activation of hyperactivated motility is observed when stallion sperm are exposed to progesterone or prostaglandin E_1 (Loux *et al.* 2013). Other hyperactivation triggers for stallion sperm have been described, but do not support protein tyrosine phosphorylation in the sperm tail. For example, three parameters of progressive motility (straight line velocity, straightness and linearity) were decreased when stallion sperm was incubated in 10% follicular fluid, indicating the acquisition of hyperactivated motility (Lange-Consiglio & Cremonesi 2012). Similarly, Leemans *et al.* (2015b) demonstrated the induction of hyperactivated motility when stallion spermatozoa were exposed to an alkaline follicular fluid fraction, while procaine also effectively induced hyperactivated motility in stallion spermatozoa; however, both did so in the absence of protein tyrosine phosphorylation (McPartlin *et al.* 2009, Leemans *et al.* 2015a).

Acrosome reaction

The Ca^{2+} -dependent release of the acrosomal contents is an essential step in mammalian fertilization because it facilitates penetration through the acellular glycoprotein coat of the oocyte. After traversing the zona pellucida, the sperm cell will fuse with the oolemma of the mature oocyte (Yanagimachi 1994). *In vivo*, the acrosome reaction takes place in the female genital tract, at the site of fertilization. The acrosome reaction is a multipoint

membrane fusion event between the sperm plasma membrane and the outer acrosomal membrane (Vigil 1989, Tsai *et al.* 2010), which results in the generation of mixed vesicles containing both plasma membrane and outer acrosomal membrane material. The remaining unfused acrosomal membranes, that is the equatorial area of the outer acrosome and the sperm plasma membrane connected to the inner acrosomal membrane covering the apical part of the nucleus, subsequently function as the sperm plasma membrane surface (Kim & Gerton 2003, Vjugina & Evans 2008). This re-designed outer sperm membrane contains a hairpin structure that is able to bind to the oolemma to enable gamete fusion and oocyte activation (Yanagimachi 1994). In boar sperm, it has been demonstrated that soluble N-ethylmaleimide- SNARE protein interactions play a fundamental role in the fusion of the sperm plasma membrane and the outer acrosomal membrane, because a *trans* ternary-soluble N-ethylmaleimide-sensitive factor attachment protein receptor protein complex is formed during capacitation. Key factors involved in this process are syntaxin 1B and VAMP 3 from the plasma membrane, and SNAP 23 from the outer acrosomal membrane (Tsai *et al.* 2010). Additionally, the entry of Ca^{2+} (*in vitro* by use of Ca^{2+} ionophores; *in vivo* after ZP binding) is required to facilitate the conversion to *cis*-soluble N-ethylmaleimide-SNARE complexes which will result in acrosomal exocytosis and penetration of the zona pellucida (Roggero *et al.* 2007).

In mice, one of the main molecules triggering the acrosome reaction in capacitated spermatozoa is zona pellucida glycoprotein 3 (ZP3) present on mature oocytes (Bleil & Wassarman 1990, Yanagimachi 1994, Arnoult *et al.* 1999). More recently, it has been shown that only contact with the intercellular matrix of the cumulus cell complex was able to induce the acrosome reaction and allow mouse spermatozoa to pass through the zona pellucida (Inoue *et al.* 2011, Jin *et al.* 2011). In many other mammals, including the horse, it has been demonstrated that capacitated, acrosome-intact spermatozoa initiate zona pellucida binding (Yanagimachi 1994), although stallion spermatozoa showed a low incidence of acrosome reaction after 1 h of *in vitro* binding to the zona pellucida (Ellington *et al.* 1993, Cheng *et al.* 1996, Meyers *et al.* 1996). Progesterone, present in follicular fluid or cumulus cell secretions, is another factor that might be responsible for inducing the acrosome reaction in horse spermatozoa (Saaranen *et al.* 1993, Cheng *et al.* 1998a, Lange-Consiglio & Cremonesi 2012) (Fig. 1). Interestingly, the progesterone-induced acrosome reaction did not proceed in a protein kinase A, but rather in a protein kinase C and protein tyrosine kinase-dependent manner (Rathi *et al.* 2003), which is very similar to the zona pellucida-mediated induction of the acrosome reaction. Indeed, Breitbart and Naor (1999) showed that zona pellucida protein 3 activates a sperm protein tyrosine

kinase coupled to phospholipase C (PLC), which in turn stimulates protein kinase C by generating diacylglycerol (DAG) from phosphatidylinositol-bisphosphate (PIP₂) (Fig. 1). How the generation of diacylglycerol eventually results in the onset of the acrosome reaction is not yet known. By contrast, when stallion spermatozoa are incubated *in vitro* in HCO₃⁻-enriched conditions, the acrosome reaction is mainly supported by a protein kinase A instead of a protein tyrosine kinase and protein kinase C-dependent pathway (Rathi *et al.* 2003) (Fig. 1), indicating that progesterone and HCO₃⁻ induce the acrosome reaction in different ways. *In vitro*, the acrosome reaction can also be reliably induced by non-physiological triggers, such as Ca²⁺ ionophore in combination with HCO₃⁻ (Rathi *et al.* 2001).

In mice and pigs, it has been demonstrated that several neurotransmitter receptor/ion channels are present in sperm cell membranes. At least two of these channels, the glycine receptor/Cl⁻ channel and a nicotinic acetylcholine receptor, are important for the ZP-initiated acrosome reaction, but not for the progesterone-initiated acrosome reaction (mouse: Meizel 2004; pig: Melendrez & Meizel 1995). In general, the glycine receptor/Cl⁻ channel is involved in inhibiting neurotransmission in the central nervous system, where glycine inhibits transmission by increasing Cl⁻ influx through the glycine receptor/Cl⁻ channel to induce hyperpolarization (Betz & Becker 1988). However, in some neurons, neutrophils, and glycerol kinase pseudogene 3 cells, glycine appears to cause depolarization by activating the glycine receptor/Cl⁻ channel (presumably stimulating Cl⁻ efflux), resulting in an increased cytoplasmic Ca²⁺ concentration (Boehm *et al.* 1997, Tapia *et al.* 1997, Weaver *et al.* 1998). The latter has also been demonstrated in sperm cells which came into contact with zona pellucida proteins. The depolarizing effect was evident as Ca²⁺ influx through voltage-gated Ca²⁺ channels (Arnoult *et al.* 1996, Florman *et al.* 1998). This event initiated the onset of the acrosome reaction. In the horse, the potential role of a glycine receptor/Cl⁻ channel and/or a nicotinic acetylcholine receptor in the onset of the acrosome reaction has not been studied. McPartlin *et al.* (2011) showed that the cAMP-driven activation of guanine-nucleotide exchange factors (RAPGEF3/RAPGEF4) induced a sperm membrane depolarization in capacitated stallion spermatozoa. Depolarization-dependent Ca²⁺ influx subsequently initiated acrosomal exocytosis. However, the activation of these factors did not play any role in the activation of protein kinase A or in protein tyrosine phosphorylation (Fig. 1).

Equine IVF: an update

Worldwide, only two IVF foals have been born, both more than 25 years ago in 1990–1991 (Palmer *et al.* 1991, Bézard *et al.* 1992). The success of IVF in these cases was attributed to *in vitro* co-incubation of equine

oocytes with calcium ionophore-treated stallion spermatozoa. Unfortunately, neither this nor other equine IVF protocols have proven to be reproducible. Indeed, subsequent attempts have yielded extremely poor equine IVF results, with reported cleavage rates varying from 0 to 33% (Zhang *et al.* 1990, Choi *et al.* 1994, Li *et al.* 1995, Dell'aquila *et al.* 1996, 1997a,b, Alm *et al.* 2001, Hinrichs *et al.* 2002, Tremoleda *et al.* 2003, Roasa *et al.* 2007, Mugnier *et al.* 2009a,b, Lange-Consiglio & Cremonesi 2012). A detailed overview of the published equine IVF studies was produced by Leemans *et al.* (2016a).

During the last decade, the utility of procaine in equine IVF media has been a topic of discussion. As mentioned previously, procaine is a potent inducer of hyperactivated motility in stallion spermatozoa (McPartlin *et al.* 2009, Leemans *et al.* 2015a). Indeed, procaine-induced hyperactivation of stallion spermatozoa can even be induced in non-capacitating medium in the absence of external Ca²⁺ (Loux *et al.* 2013), with CATSPER channels apparently not participating in procaine-induced hyperactivation (Loux *et al.* 2013). Initial reports further suggested that procaine promoted equine *in vitro* fertilization by inducing hyperactivated motility while other requisite sperm capacitation events, such

Figure 3 Methods to distinguish fertilization from parthenogenesis. (A) Previously the presence of 2 pronuclei (2 PN), evaluated using nuclear stains such as propidium iodide, was considered adequate to indicate fertilization of equine oocytes. However, this method cannot exclude parthenogenesis. Therefore, the additional presence of the sperm tail (ST) and the second extruded polar body (PB) shown by (B) MitoTracker Green FM pre-labeling combined with Hoechst post-fixation (Leemans *et al.* 2015a; published with permission from Oxford University Press (permission number 4450310125499)); (C) lacmoid post-fixation, should be demonstrated; or (D) paternal (pPN) and maternal (mPN) pronuclei in equine zygotes should be distinguished by different histone 3 methylation (H3K9me3) patterns.

as protein tyrosine phosphorylation, were proposed to have been triggered by other medium constituents or a high pH (McPartlin *et al.* 2009, Ambruosi *et al.* 2013). Cleavage rates exceeding 60% of co-incubated oocytes were reported. However, none of these cleaved oocytes developed to the blastocyst stage. Leemans *et al.* (2015a) subsequently demonstrated that, although procaine did indeed trigger sperm hyperactivation, procaine-induced oocyte cytoplasmic cleavage also occurred in the absence of sperm. Moreover, gamete fusion induced oocyte activation, indicated by cytoplasmic Ca^{2+} oscillations, was not observed in the presence of procaine. It transpired that procaine acted further downstream in the oocyte-activation process, and only induced cytokinesis-like events via pH-dependent depolymerization of F-actin, whereas nuclear activation was not triggered. Moreover, the procaine concentration used to induce sperm hyperactivated motility and oocyte cytokinesis was DNA toxic (Leemans *et al.* 2015a). Rather than true fertilization, procaine-induced DNA fragmentation with degeneration of the cleaving blastomeres occurring at the 8–16 cell stage, that is stage of genome activation in horse embryos (Brinsko *et al.* 1995).

As a result of the initial misinterpretation, more stringent evaluation methods are clearly needed to confirm fertilization in horse oocytes. For various mammals, the presence of two pronuclei after 20–24 h gamete co-incubation, as visualized by nuclear chromatin stains such as Hoechst or propidium iodide, is considered valid proof of fertilization because of the low incidence of parthenogenesis (around 5%) in current farm animal IVF systems (Beek *et al.* 2012, Heras *et al.* 2014, 2015). However, the presence of two pronuclei is not sufficient to distinguish normal fertilization from parthenogenesis in the horse. Using techniques such as pre-labeling sperm with MitoTracker Green FM/Hoechst, and lacmoid post-fixation, Leemans *et al.* (2015a) were able to differentiate fertilization (two polar bodies within the perivitelline space, two pronuclei and a sperm tail within the ooplasm) from parthenogenesis (one polar body, one or two pronuclei and no sperm tail) (Fig. 3). Recently, a staining technique to differentiate the paternal from the maternal pronucleus in equine zygotes produced by intracytoplasmic sperm injection (ICSI) was established (Heras *et al.* 2015), based on different histone 3 methylation (H3K9me3) patterns. This staining technique could, however, not be used in the procaine experiments because the procaine-exposed oocytes did not form normal pronuclei but instead produced condensed DNA fragments (Fig. 3).

Conclusion

For more than 25 years, co-incubation of mature oocytes with capacitated sperm has been the standard method for producing *in vitro* embryos in several species including

man, cattle, pigs and many laboratory animals. In the horse, however, conventional IVF still does not work. Inadequate capacitation of stallion spermatozoa under *in vitro* conditions is probably the major obstacle. It appears that several factors required to induce capacitation in other mammalian species are not involved in equine sperm capacitation. However, biological and chemical capacitation triggers have been identified to facilitate tail-associated protein tyrosine phosphorylation and hyperactivated motility in stallion sperm. However, since the acrosome reaction cannot be induced reliably using these factors, further research should focus on capacitation-related changes that prepare the sperm plasma and outer acrosomal membranes for fusion. It will be necessary to verify that 'improved' capacitation conditions support (1) cholesterol depletion from the sperm plasma membrane, (2) lateral redistribution of lipid rafts in the apical region and ultimately (3) the zona pellucida-induced acrosome reaction. It is likely that the induction of appropriate sperm plasma membrane changes will support the acrosome reaction. Only fully capacitated stallion spermatozoa will be able to penetrate the cumulus and zona pellucida barriers to accomplish fertilization.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was supported by the Research Foundation – Flanders (FWO-Flanders; grant number 12I0517N) and EU COST Action 16119 CellFit.

References

- Aitken RJ 2011 The capacitation-apoptosis highway: oxysterols and mammalian sperm function. *Biology of Reproduction* **85** 9–12. (<https://doi.org/10.1095/biolreprod.111.092528>)
- Aitken RJ, Wang YF, Liu J, Best F & Richardson DW 1983 The influence of medium composition, osmolarity and albumin content on the acrosome reaction and fertilizing capacity of human spermatozoa: development of an improved zona-free hamster egg penetration test. *International Journal of Andrology* **6** 180–193. (<https://doi.org/10.1111/j.1365-2605.1983.tb00337.x>)
- Aitken RJ, Ryan AL, Baker MA & McLaughlin EA 2004 Redox activity associated with the maturation and capacitation of mammalian spermatozoa. *Free Radical Biology and Medicine* **36** 994–1010. (<https://doi.org/10.1016/j.freeradbiomed.2004.01.017>)
- Alm H, Torner H, Blottner S, Nurnberg G & Kanitz W 2001 Effect of sperm cryopreservation and treatment with calcium ionophore or heparin on *in vitro* fertilization of horse oocytes. *Theriogenology* **56** 817–829. ([https://doi.org/10.1016/S0093-691X\(01\)00610-0](https://doi.org/10.1016/S0093-691X(01)00610-0))
- Ambruosi B, Accogli G, Douet C, Canepa S, Pascal G, Monget P, Moros Nicolas C, Holmskov U, Mollenhauer J, Robbe-Masselot C *et al.* 2013 Deleted in malignant brain tumor 1 is secreted in the oviduct and involved in the mechanism of fertilization in equine and porcine species. *Reproduction* **146** 119–133. (<https://doi.org/10.1530/REP-13-0007>)

- Arnoult C, Cardullo RA, Lemos JR & Florman HM 1996 Activation of mouse sperm T-type Ca²⁺ channels by adhesion to the egg zona pellucida. *PNAS* **93** 13004–13009. (<https://doi.org/10.1073/pnas.93.23.13004>)
- Arnoult C, Kazam IG, Visconti PE, Kopf GS, Villaz M & Florman HM 1999 Control of the low voltage-activated calcium channel of mouse sperm by egg ZP3 and by membrane hyperpolarization during capacitation. *PNAS* **96** 6757–6762. (<https://doi.org/10.1073/pnas.96.12.6757>)
- Austin CR 1952 The capacitation of the mammalian sperm. *Nature* **170** 326. (<https://doi.org/10.1038/170326a0>)
- Bailey JL, Bilodeau JF & Cormier N 2000 Semen cryopreservation in domestic animals: a damaging and capacitating phenomenon. *Journal of Andrology* **21** 1–7. (<https://doi.org/10.1002/j.1939-4640.2000.tb03268.x>)
- Balao da Silva CM, Spinaci M, Bucci D, Giarretta E, Pena FJ, Mari G & Galeati G 2013 Effect of sex sorting on stallion spermatozoa: heterologous oocyte binding, tyrosine phosphorylation and acrosome reaction assay. *Animal Reproduction Science* **141** 68–74. (<https://doi.org/10.1016/j.anireprosci.2013.07.008>)
- Ball BA, Gravance CG, Medina V, Baumber J & Liu IK 2000 Catalase activity in equine semen. *American Journal of Veterinary Research* **61** 1026–1030. (<https://doi.org/10.2460/ajvr.2000.61.1026>)
- Baumber J, Sabeur K, Vo A & Ball BA 2003 Reactive oxygen species promote tyrosine phosphorylation and capacitation in equine spermatozoa. *Theriogenology* **60** 1239–1247. ([https://doi.org/10.1016/S0093-691X\(03\)00144-4](https://doi.org/10.1016/S0093-691X(03)00144-4))
- Beek J, Nauwynck H, Maes D & Van Soom A 2012 Inhibitors of zinc-dependent metalloproteases hinder sperm passage through the cumulus oophorus during porcine fertilization *in vitro*. *Reproduction* **144** 687–697. (<https://doi.org/10.1530/REP-12-0311>)
- Benoff S 1998 Voltage dependent calcium channels in mammalian spermatozoa. *Frontiers in Bioscience* **3** D1220–D1240. (<https://doi.org/10.2741/A358>)
- Betteridge KJ 2006 Farm animal embryo technologies: achievements and perspectives. *Theriogenology* **65** 905–913. (<https://doi.org/10.1016/j.theriogenology.2005.09.005>)
- Betz H & Becker CM 1988 The mammalian glycine receptor: biology and structure of a neuronal chloride channel protein. *Neurochemistry International* **13** 137–146. ([https://doi.org/10.1016/0197-0186\(88\)90048-4](https://doi.org/10.1016/0197-0186(88)90048-4))
- Bézar J, Magistrini M, Battut I, Duchamp G & Palmer E 1992 *In vitro* fertilization in the mare. *Recueil De Medecine Veterinaire* **168** 993–1003.
- Bielfeld P, Anderson RA, Mack SR, De Jonge CJ & Zaneveld LJ 1994 Are capacitation or calcium ion influx required for the human sperm acrosome reaction? *Fertility and Sterility* **62** 1255–1261. ([https://doi.org/10.1016/S0015-0282\(16\)57195-7](https://doi.org/10.1016/S0015-0282(16)57195-7))
- Birck A, Labouriau R & Christensen P 2009 Dynamics of the induced acrosome reaction in boar sperm evaluated by flow cytometry. *Animal Reproduction Science* **115** 124–136. (<https://doi.org/10.1016/j.anireprosci.2008.10.017>)
- Bleil JD & Wassarman PM 1990 Identification of a ZP3-binding protein on acrosome-intact mouse sperm by photoaffinity crosslinking. *PNAS* **87** 5563–5567. (<https://doi.org/10.1073/pnas.87.14.5563>)
- Boehm S, Harvey RJ, von Holst A, Rohrer H & Betz H 1997 Glycine receptors in cultured chick sympathetic neurons are excitatory and trigger neurotransmitter release. *Journal of Physiology* **504** 683–694. (<https://doi.org/10.1111/j.1469-7793.1997.683bd.x>)
- Boerke A, Tsai PS, Garcia-Gil N, Brewis IA & Gadella BM 2008 Capacitation-dependent reorganization of microdomains in the apical sperm head plasma membrane: functional relationship with zona binding and the zona-induced acrosome reaction. *Theriogenology* **70** 1188–1196. (<https://doi.org/10.1016/j.theriogenology.2008.06.021>)
- Boerke A, Brouwers JF, Olkkonen VM, van de Lest CH, Sostaric E, Schoevers EJ, Helms JB & Gadella BM 2013 Involvement of bicarbonate-induced radical signaling in oxysterol formation and sterol depletion of capacitating mammalian sperm during *in vitro* fertilization. *Biology of Reproduction* **88** 21. (<https://doi.org/10.1095/biolreprod.112.101253>)
- Breining E, Cetica PD & Beconi MT 2010 Capacitation inducers act through diverse intracellular mechanisms in cryopreserved bovine sperm. *Theriogenology* **74** 1036–1049. (<https://doi.org/10.1016/j.theriogenology.2010.04.034>)
- Breitbart H & Naor Z 1999 Protein kinases in mammalian sperm capacitation and the acrosome reaction. *Reviews of Reproduction* **4** 151–159. (<https://doi.org/10.1530/ror.0.0040151>)
- Brinsko SP, Ball BA, Ignatz GG, Thomas PG, Currie WB & Ellington JE 1995 Initiation of transcription and nucleogenesis in equine embryos. *Molecular Reproduction and Development* **42** 298–302. (<https://doi.org/10.1002/mrd.1080420306>)
- Bromfield EG, Aitken RJ, Gibb Z, Lambourne SR & Nixon B 2014 Capacitation in the presence of methyl-beta-cyclodextrin results in enhanced zona pellucida-binding ability of stallion spermatozoa. *Reproduction* **147** 153–166. (<https://doi.org/10.1530/REP-13-0393>)
- Bromfield EG, McLaughlin EA, Aitken RJ & Nixon B 2016 Heat shock protein member A2 forms a stable complex with angiotensin converting enzyme and protein disulfide isomerase A6 in human spermatozoa. *Molecular Human Reproduction* **22** 93–109. (<https://doi.org/10.1093/molehr/gav073>)
- Brouwers JF, Boerke A, Silva PF, Garcia-Gil N, van Gestel RA, Helms JB, van de Lest CH & Gadella BM 2011 Mass spectrometric detection of cholesterol oxidation in bovine sperm. *Biology of Reproduction* **85** 128–136. (<https://doi.org/10.1095/biolreprod.111.091207>)
- Burnaugh L, Ball BA, Sabeur K, Thomas AD & Meyers SA 2010 Osmotic stress stimulates generation of superoxide anion by spermatozoa in horses. *Animal Reproduction Science* **117** 249–260. (<https://doi.org/10.1016/j.anireprosci.2009.05.014>)
- Buschiazio J, Ialy-Radio C, Auer J, Wolf JP, Serres C, Lefevre B & Ziyat A 2013 Cholesterol depletion disorganizes oocyte membrane rafts altering mouse fertilization. *PLoS One* **8** e62919. (<https://doi.org/10.1371/journal.pone.0062919>)
- Byrd W 1981 *In vitro* capacitation and the chemically induced acrosome reaction in bovine spermatozoa. *Journal of Experimental Zoology* **215** 35–46. (<https://doi.org/10.1002/jez.1402150105>)
- Carlson AE, Westenbroek RE, Quill T, Ren D, Clapham DE, Hille B, Garbers DL & Babcock DF 2003 CatSper1 required for evoked Ca²⁺ entry and control of flagellar function in sperm. *PNAS* **100** 14864–14868. (<https://doi.org/10.1073/pnas.2536658100>)
- Carlson AE, Quill TA, Westenbroek RE, Schuh SM, Hille B & Babcock DF 2005 Identical phenotypes of CatSper1 and CatSper2 null sperm. *Journal of Biological Chemistry* **280** 32238–32244. (<https://doi.org/10.1074/jbc.M501430200>)
- Carrera A, Gerton GL & Moss SB 1994 The major fibrous sheath polypeptide of mouse sperm: structural and functional similarities to the A-kinase anchoring proteins. *Developmental Biology* **165** 272–284. (<https://doi.org/10.1006/dbio.1994.1252>)
- Carrera A, Moos J, Ning XP, Gerton GL, Tesarik J, Kopf GS & Moss SB 1996 Regulation of protein tyrosine phosphorylation in human sperm by a calcium/calmodulin-dependent mechanism: identification of A kinase anchor proteins as major substrates for tyrosine phosphorylation. *Developmental Biology* **180** 284–296. (<https://doi.org/10.1006/dbio.1996.0301>)
- Castellano LE, Trevino CL, Rodriguez D, Serrano CJ, Pacheco J, Tsutsumi V, Felix R & Darszon A 2003 Transient receptor potential (TRPC) channels in human sperm: expression, cellular localization and involvement in the regulation of flagellar motility. *FEBS Letters* **541** 69–74. ([https://doi.org/10.1016/S0014-5793\(03\)00305-3](https://doi.org/10.1016/S0014-5793(03)00305-3))
- Chang MC 1951 Fertilizing capacity of spermatozoa deposited into the fallopian tubes. *Nature* **168** 697–698. (<https://doi.org/10.1038/168697b0>)
- Cheng FP, Fazeli A, Voorhout WF, Marks A, Bevers MM & Colenbrander B 1996 Use of peanut agglutinin to assess the acrosomal status and the zona pellucida-induced acrosome reaction in stallion spermatozoa. *Journal of Andrology* **17** 674–682.
- Cheng FP, Fazeli AR, Voorhout WF, Tremoleda JL, Bevers MM & Colenbrander B 1998a Progesterone in mare follicular fluid induces the acrosome reaction in stallion spermatozoa and enhances *in vitro* binding to the zona pellucida. *International Journal of Andrology* **21** 57–66. (<https://doi.org/10.1046/j.1365-2605.1998.00096.x>)
- Cheng FP, Gadella BM, Voorhout WF, Fazeli A, Bevers MM & Colenbrander B 1998b Progesterone-induced acrosome reaction in stallion spermatozoa is mediated by a plasma membrane progesterone receptor. *Biology of Reproduction* **59** 733–742. (<https://doi.org/10.1095/biolreprod59.4.733>)
- Choi YH, Okada Y, Hochi S, Braun J, Sato K & Oguiri N 1994 *In vitro* fertilization rate of horse oocytes with partially removed zonae. *Theriogenology* **42** 795–802. ([https://doi.org/10.1016/0093-691X\(94\)90448-R](https://doi.org/10.1016/0093-691X(94)90448-R))
- Christensen P, Whitfield CH & Parkinson TJ 1996 *In vitro* induction of acrosome reactions in stallion spermatozoa by heparin and A23187.

- Theriogenology* **45** 1201–1210. ([https://doi.org/10.1016/0093-691X\(96\)00075-1](https://doi.org/10.1016/0093-691X(96)00075-1))
- Contreras L, Drago I, Zampese E & Pozzan T 2010 Mitochondria: the calcium connection. *Biochimica et Biophysica Acta* **1797** 607–618. (<https://doi.org/10.1016/j.bbabi.2010.05.005>)
- Costello S, Michelangeli F, Nash K, Lefievre L, Morris J, Machado-Oliveira G, Barratt C, Kirkman-Brown J & Publicover S 2009 Ca²⁺-stores in sperm: their identities and functions. *Reproduction* **138** 425–437. (<https://doi.org/10.1530/REP-09-0134>)
- Darszon A, Lopez-Martinez P, Acevedo JJ, Hernandez-Cruz A & Trevino CL 2006 T-type Ca²⁺ channels in sperm function. *Cell Calcium* **40** 241–252. (<https://doi.org/10.1016/j.ceca.2006.04.028>)
- Davila MP, Muñoz PM, Bolaños JM, Stout TA, Gadella BM, Tapia JA, da Silva CB, Ferrusola CO & Peña FJ 2016 Mitochondrial ATP is required for the maintenance of membrane integrity in stallion spermatozoa, whereas motility requires both glycolysis and oxidative phosphorylation. *Reproduction* **152** 683–694. (<https://doi.org/10.1530/REP-16-0409>)
- de Lamirande E & Gagnon C 1993 A positive role for the superoxide anion in triggering hyperactivation and capacitation of human spermatozoa. *International Journal of Andrology* **16** 21–25. (<https://doi.org/10.1111/j.1365-2605.1993.tb01148.x>)
- Dell'Aquila ME, Fusco S, Lacalandra GM & Maritato F 1996 *In vitro* maturation and fertilization of equine oocytes recovered during the breeding season. *Theriogenology* **45** 547–560. ([https://doi.org/10.1016/0093-691X\(95\)00402-T](https://doi.org/10.1016/0093-691X(95)00402-T))
- Dell'Aquila ME, Cho YS, Minoia P, Traina V, Fusco S, Lacalandra GM & Maritato F 1997a Intracytoplasmic sperm injection (ICSI) versus conventional IVF on abattoir-derived and *in vitro*-matured equine oocytes. *Theriogenology* **47** 1139–1156. ([https://doi.org/10.1016/S0093-691X\(97\)00095-2](https://doi.org/10.1016/S0093-691X(97)00095-2))
- Dell'Aquila ME, Cho YS, Minoia P, Traina V, Lacalandra GM & Maritato F 1997b Effects of follicular fluid supplementation of *in vitro* maturation medium on the fertilization and development of equine oocytes after *in vitro* fertilization or intracytoplasmic sperm injection. *Human Reproduction* **12** 2766–2772. (<https://doi.org/10.1093/humrep/12.12.2766>)
- du Plessis SS, Agarwal A, Mohanty G & van der Linde M 2015 Oxidative phosphorylation versus glycolysis: what fuel do spermatozoa use? *Asian Journal of Andrology* **17** 230–235. (<https://doi.org/10.4103/1008-682X.135123>)
- Ellington JE, Ball BA & Yang X 1993 Binding of stallion spermatozoa to the equine zona pellucida after coculture with oviductal epithelial cells. *Journal of Reproduction and Fertility* **98** 203–208. (<https://doi.org/10.1530/jrf.0.0980203>)
- Ficarro S, Chertihin O, Westbrook VA, White F, Jayes F, Kalab P, Marto JA, Shabanowitz J, Herr JC, Hunt DF *et al.* 2003 Phosphoproteome analysis of capacitated human sperm. Evidence of tyrosine phosphorylation of a kinase-anchoring protein 3 and valosin-containing protein/p97 during capacitation. *Journal of Biological Chemistry* **278** 11579–11589. (<https://doi.org/10.1074/jbc.M202325200>)
- Flesch FM & Gadella BM 2000 Dynamics of the mammalian sperm plasma membrane in the process of fertilization. *Biochimica et Biophysica Acta* **1469** 197–235. ([https://doi.org/10.1016/S0304-4157\(00\)00018-6](https://doi.org/10.1016/S0304-4157(00)00018-6))
- Flesch FM, Colenbrander B, van Golde LM & Gadella BM 1999 Capacitation induces tyrosine phosphorylation of proteins in the boar sperm plasma membrane. *Biochemical and Biophysical Research Communications* **262** 787–792. (<https://doi.org/10.1006/bbrc.1999.1300>)
- Flesch FM, Brouwers JF, Nievelstein PF, Verkleij AJ, van Golde LM, Colenbrander B & Gadella BM 2001 Bicarbonate stimulated phospholipid scrambling induces cholesterol redistribution and enables cholesterol depletion in the sperm plasma membrane. *Journal of Cell Science* **114** 3543–3555.
- Florman HM, Arnoult C, Kazam IG, Li C & O'Toole CM 1998 A perspective on the control of mammalian fertilization by egg-activated ion channels in sperm: a tale of two channels. *Biology of Reproduction* **59** 12–16. (<https://doi.org/10.1095/biolreprod59.1.12>)
- Fraser LR, Abeydeera LR & Niwa K 1995 Ca(2+)-regulating mechanisms that modulate bull sperm capacitation and acrosomal exocytosis as determined by chlortetracycline analysis. *Molecular Reproduction and Development* **40** 233–241. (<https://doi.org/10.1002/mrd.1080400213>)
- Funahashi H & Nagai T 2001 Regulation of *in vitro* penetration of frozen-thawed boar spermatozoa by caffeine and adenosine. *Molecular Reproduction and Development* **58** 424–431. ([https://doi.org/10.1002/1098-2795\(20010401\)58:4<424::AID-MRD10>3.0.CO;2-1](https://doi.org/10.1002/1098-2795(20010401)58:4<424::AID-MRD10>3.0.CO;2-1))
- Gadella BM & Harrison RA 2000 The capacitating agent bicarbonate induces protein kinase A-dependent changes in phospholipid transbilayer behavior in the sperm plasma membrane. *Development* **127** 2407–2420.
- Gadella BM & Harrison RA 2002 Capacitation induces cyclic adenosine 3',5'-monophosphate-dependent, but apoptosis-unrelated, exposure of aminophospholipids at the apical head plasma membrane of boar sperm cells. *Biology of Reproduction* **67** 340–350. (<https://doi.org/10.1095/biolreprod67.1.340>)
- Gadella BM, Gadella TW Jr, Colenbrander B, van Golde LM & Lopes-Cardozo M 1994 Visualization and quantification of glycolipid polarity dynamics in the plasma membrane of the mammalian spermatozoon. *Journal of Cell Science* **107** (Pt 8) 2151–2163.
- Gadella BM, Lopes-Cardozo M, van Golde LM, Colenbrander B & Gadella TW, Jr 1995 Glycolipid migration from the apical to the equatorial subdomains of the sperm head plasma membrane precedes the acrosome reaction. Evidence for a primary capacitation event in boar spermatozoa. *Journal of Cell Science* **108** (Pt 3) 935–946.
- Gadella BM, Tsai PS, Boerke A & Brevis IA 2008 Sperm head membrane reorganisation during capacitation. *International Journal of Developmental Biology* **52** 473–480. (<https://doi.org/10.1387/ijdb.082583bg>)
- Galli C, Duchi R, Crotti G, Turini P, Ponderato N, Colleoni S, Lagutina I & Lazzari G 2003 Bovine embryo technologies. *Theriogenology* **59** 599–616. ([https://doi.org/10.1016/S0093-691X\(02\)01243-8](https://doi.org/10.1016/S0093-691X(02)01243-8))
- Gibb Z, Lambourne SR & Aitken RJ 2014 The paradoxical relationship between stallion fertility and oxidative stress. *Biology of Reproduction* **91** 77. (<https://doi.org/10.1095/biolreprod.114.118539>)
- Gibb Z, Lambourne SR, Quadrelli J, Smith ND & Aitken RJ 2015 L-carnitine and pyruvate are prosurvival factors during the storage of stallion spermatozoa at room temperature. *Biology of Reproduction* **93** 104. (<https://doi.org/10.1095/biolreprod.115.131326>)
- Gibb Z, Lambourne SR, Curry BJ, Hall SE & Aitken RJ 2016 Aldehyde dehydrogenase plays a pivotal role in the maintenance of stallion sperm motility. *Biology of Reproduction* **94** 133. (<https://doi.org/10.1095/biolreprod.116.140509>)
- Giroux-Widemann V, Jouannet P, Pignot-Paintrand I & Feneux D 1991 Effects of pH on the reactivation of human spermatozoa demembrated with Triton X-100. *Molecular Reproduction and Development* **29** 157–162. (<https://doi.org/10.1002/mrd.1080290211>)
- Gonzalez-Fernandez L, Macias-Garcia B, Velez IC, Varner DD & Hinrichs K 2012 Calcium-calmodulin and pH regulate protein tyrosine phosphorylation in stallion sperm. *Reproduction* **144** 411–422. (<https://doi.org/10.1530/REP-12-0067>)
- Gonzalez-Fernandez L, Macias-Garcia B, Loux SC, Varner DD & Hinrichs K 2013 Focal adhesion kinases and calcium/calmodulin-dependent protein kinases regulate protein tyrosine phosphorylation in stallion sperm. *Biology of Reproduction* **88** 138. (<https://doi.org/10.1095/biolreprod.112.107078>)
- Graham JK 1996 Methods for induction of capacitation and the acrosome reaction of stallion spermatozoa. *Veterinary Clinics of North America. Equine Practice* **12** 111–117. ([https://doi.org/10.1016/S0749-0739\(17\)30298-5](https://doi.org/10.1016/S0749-0739(17)30298-5))
- Gualtieri R, Boni R, Tosti E, Zagami M & Talevi R 2005 Intracellular calcium and protein tyrosine phosphorylation during the release of bovine sperm adhering to the fallopian tube epithelium *in vitro*. *Reproduction* **129** 51–60. (<https://doi.org/10.1530/rep.1.00374>)
- Harayama H, Noda T, Ishikawa S & Shidara O 2012 Relationship between cyclic AMP-dependent protein tyrosine phosphorylation and extracellular calcium during hyperactivation of boar spermatozoa. *Molecular Reproduction and Development* **79** 727–739. (<https://doi.org/10.1002/mrd.22106>)
- Hemachand T & Shaha C 2003 Functional role of sperm surface glutathione S-transferases and extracellular glutathione in the haploid spermatozoa under oxidative stress. *FEBS Letters* **538** 14–18. ([https://doi.org/10.1016/S0014-5793\(03\)00103-0](https://doi.org/10.1016/S0014-5793(03)00103-0))
- Heras S, Forier K, Rombouts K, Braeckmans K & Van Soom A 2014 DNA counterstaining for methylation and hydroxymethylation immunostaining in bovine zygotes. *Analytical Biochemistry* **454** 14–16. (<https://doi.org/10.1016/j.ab.2014.03.002>)

- Heras S, Smits K, Leemans B & Van Soom A 2015 Asymmetric histone 3 methylation pattern between paternal and maternal pronuclei in equine zygotes. *Analytical Biochemistry* **471** 67–69. (<https://doi.org/10.1016/j.ab.2014.11.005>)
- Hernandez-Gonzalez EO, Trevino CL, Castellano LE, de la Vega-Beltran JL, Ocampo AY, Wertheimer E, Visconti PE & Darszon A 2007 Involvement of cystic fibrosis transmembrane conductance regulator in mouse sperm capacitation. *Journal of Biological Chemistry* **282** 24397–24406. (<https://doi.org/10.1074/jbc.M701603200>)
- Hess KC, Jones BH, Marquez B, Chen Y, Ord TS, Kamenetsky M, Miyamoto C, Zippin JH, Kopf GS, Suarez SS *et al.* 2005 The “soluble” adenylyl cyclase in sperm mediates multiple signaling events required for fertilization. *Developmental Cell* **9** 249–259. (<https://doi.org/10.1016/j.devcel.2005.06.007>)
- Hinrichs K, Love CC, Brinsko SP, Choi YH & Varner DD 2002 *In vitro* fertilization of *in vitro*-matured equine oocytes: effect of maturation medium, duration of maturation, and sperm calcium ionophore treatment, and comparison with rates of fertilization *in vivo* after oviductal transfer. *Biology of Reproduction* **67** 256–262. (<https://doi.org/10.1095/biolreprod67.1.256>)
- Ho HC & Suarez SS 2001a Hyperactivation of mammalian spermatozoa: function and regulation. *Reproduction* **122** 519–526. (<https://doi.org/10.1530/rep.0.1220519>)
- Ho HC & Suarez SS 2001b An inositol 1,4,5-trisphosphate receptor-gated intracellular Ca(2+) store is involved in regulating sperm hyperactivated motility. *Biology of Reproduction* **65** 1606–1615. (<https://doi.org/10.1095/biolreprod65.5.1606>)
- Ho HC & Suarez SS 2003 Characterization of the intracellular calcium store at the base of the sperm flagellum that regulates hyperactivated motility. *Biology of Reproduction* **68** 1590–1596. (<https://doi.org/10.1095/biolreprod.102.011320>)
- Ho HC, Granish KA & Suarez SS 2002 Hyperactivated motility of bull sperm is triggered at the axoneme by Ca2+ and not cAMP. *Developmental Biology* **250** 208–217. (<https://doi.org/10.1006/dbio.2002.0797>)
- Humes HD & Weinberg JM 1980 Ionophore A23187 induced reductions in toad urinary bladder epithelial cell oxidative phosphorylation and viability: implications for A23187 related declines in epithelial active transport. *Pflügers Archiv* **388** 217–230. (<https://doi.org/10.1007/BF00658484>)
- Ignatz GG & Suarez SS 2005 Calcium/calmodulin and calmodulin kinase II stimulate hyperactivation in demembranated bovine sperm. *Biology of Reproduction* **73** 519–526. (<https://doi.org/10.1095/biolreprod.105.040733>)
- Ijiri TW, Mahub Hasan AK & Sato K 2012 Protein-tyrosine kinase signaling in the biological functions associated with sperm. *Journal of Signal Transduction* **2012** 181560. (<https://doi.org/10.1155/2012/181560>)
- Inoue N, Satouh Y, Ikawa M, Okabe M & Yanagimachi R 2011 Acrosome-reacted mouse spermatozoa recovered from the perivitelline space can fertilize other eggs. *PNAS* **108** 20008–20011. (<https://doi.org/10.1073/pnas.1116965108>)
- Ishijima S & Witman GB 1987 Flagellar movement of intact and demembranated, reactivated ram spermatozoa. *Cell Motility and the Cytoskeleton* **8** 375–391. (<https://doi.org/10.1002/cm.970080410>)
- Ishijima S, Mohri H, Overstreet JW & Yudin AI 2006 Hyperactivation of monkey spermatozoa is triggered by Ca2+ and completed by cAMP. *Molecular Reproduction and Development* **73** 1129–1139. (<https://doi.org/10.1002/mrd.20420>)
- Jin M, Fujiwara E, Kakiuchi Y, Okabe M, Satouh Y, Baba SA, Chiba K & Hirohashi N 2011 Most fertilizing mouse spermatozoa begin their acrosome reaction before contact with the zona pellucida during *in vitro* fertilization. *PNAS* **108** 4892–4896. (<https://doi.org/10.1073/pnas.1018202108>)
- Jones DP 2006 Redefining oxidative stress. *Antioxidants and Redox Signaling* **8** 1865–1879. (<https://doi.org/10.1089/ars.2006.8.1865>)
- Kim KS & Gerton GL 2003 Differential release of soluble and matrix components: evidence for intermediate states of secretion during spontaneous acrosomal exocytosis in mouse sperm. *Developmental Biology* **264** 141–152. (<https://doi.org/10.1016/j.ydbio.2003.08.006>)
- Krumschnabel G, Schwarzbach PJ & Wieser W 1999 Energetics of trout hepatocytes during A23187-induced disruption of Ca2+ homeostasis. *Comparative Biochemistry and Physiology. Part C, Pharmacology, Toxicology and Endocrinology* **124** 187–195. ([https://doi.org/10.1016/S0742-8413\(99\)00058-4](https://doi.org/10.1016/S0742-8413(99)00058-4))
- Lange-Consiglio A & Cremonesi F 2012 163 Hyperactivation of stallion sperm in follicular fluid for *in vitro* fertilization of equine oocytes. *Reproduction, Fertility and Development* **24** 193–194. (<https://doi.org/10.1071/RDv24n1Ab163>)
- Leclerc P, de Lamirande E & Gagnon C 1996 Cyclic adenosine 3',5'-monophosphate-dependent regulation of protein tyrosine phosphorylation in relation to human sperm capacitation and motility. *Biology of Reproduction* **55** 684–692. (<https://doi.org/10.1095/biolreprod55.3.684>)
- Leemans B, Gadella BM, Sostaric E, Nelis H, Stout TA, Hoogewijs M & Van Soom A 2014 Oviduct binding and elevated environmental pH induce protein tyrosine phosphorylation in stallion spermatozoa. *Biology of Reproduction* **91** 13. (<https://doi.org/10.1095/biolreprod.113.116418>)
- Leemans B, Gadella BM, Stout TA, Heras S, Smits K, Ferrer-Buitrago M, Claes E, Heindryckx B, De Vos WH, Nelis H *et al.* 2015a Procaine induces cytokinesis in horse oocytes via a pH dependent mechanism. *Biology of Reproduction* **93** 23. (<https://doi.org/10.1095/biolreprod.114.127423>)
- Leemans B, Gadella BM, Stout TA, Nelis H, Hoogewijs M & Van Soom A 2015b An alkaline follicular fluid fraction induces capacitation and limited release of oviduct epithelium-bound stallion sperm. *Reproduction* **150** 193–208. (<https://doi.org/10.1530/REP-15-0178>)
- Leemans B, Gadella BM, Stout TA, De Schauwer C, Nelis H, Hoogewijs M & Van Soom A 2016a Why doesn't conventional IVF work in the horse? The equine oviduct as a microenvironment for capacitation/fertilization. *Reproduction* **152** R233–R245. (<https://doi.org/10.1530/REP-16-0420>)
- Leemans B, Gadella BM, Stout TA, Sostaric E, Schauwer CD, Nelis H, Hoogewijs M & Van Soom A 2016b Combined albumin and bicarbonate induces head-to-head sperm agglutination which physically prevents equine sperm-oviduct binding. *Reproduction* **151** 313–330. (<https://doi.org/10.1530/REP-15-0471>)
- Li LY, Meintjes M, Graff KJ, Paul JB, Denniston RS & Godke RA 1995 *In vitro* fertilization and development of *in vitro*-matured oocytes aspirated from pregnant mares. *Biology of Reproduction* **1** 309–317. (https://doi.org/10.1093/biolreprod/52.monograph_series1.309)
- Liguori L, Rambotti MG, Bellezza I & Minelli A 2004 Electron microscopic cytochemistry of adenylyl cyclase activity in mouse spermatozoa. *Journal of Histochemistry and Cytochemistry* **52** 833–836. (<https://doi.org/10.1369/jhc.3B6141.2004>)
- Lindemann CB & Lesich KA 2010 Flagellar and ciliary beating: the proven and the possible. *Journal of Cell Science* **123** 519–528. (<https://doi.org/10.1242/jcs.051326>)
- Lishko PV, Botchkina IL, Fedorenko A & Kirichok Y 2010 Acid extrusion from human spermatozoa is mediated by flagellar voltage-gated proton channel. *Cell* **140** 327–337. (<https://doi.org/10.1016/j.cell.2009.12.053>)
- Lishko PV, Botchkina IL & Kirichok Y 2011 Progesterone activates the principal Ca2+ channel of human sperm. *Nature* **471** 387–391. (<https://doi.org/10.1038/nature09767>)
- Litvin TN, Kamenetsky M, Zarifyan A, Buck J & Levin LR 2003 Kinetic properties of “soluble” adenylyl cyclase. Synergism between calcium and bicarbonate. *Journal of Biological Chemistry* **278** 15922–15926. (<https://doi.org/10.1074/jbc.M212475200>)
- Liu B, Wang P, Wang Z & Zhang W 2011 The use of anti-VDAC2 antibody for the combined assessment of human sperm acrosome integrity and ionophore A23187-induced acrosome reaction. *PLoS One* **6** e16985. (<https://doi.org/10.1371/journal.pone.0016985>)
- Loux SC, Crawford KR, Ing NH, Gonzalez-Fernandez L, Macias-Garcia B, Love CC, Varner DD, Velez IC, Choi YH & Hinrichs K 2013 CatSper and the relationship of hyperactivated motility to intracellular calcium and pH kinetics in equine sperm. *Biology of Reproduction* **89** 123. (<https://doi.org/10.1095/biolreprod.113.111708>)
- Loux SC, Macias-Garcia B, Gonzalez-Fernandez L, Canesin HD, Varner DD & Hinrichs K 2014 Regulation of axonemal motility in demembranated equine sperm. *Biology of Reproduction* **91** 152. (<https://doi.org/10.1095/biolreprod.114.122804>)
- Maas DH, Storey BT & Mastroianni L Jr 1977 Hydrogen ion and carbon dioxide content of the oviductal fluid of the rhesus monkey (*Macaca mulatta*). *Fertility and Sterility* **28** 981–985. ([https://doi.org/10.1016/S0015-0282\(16\)42801-3](https://doi.org/10.1016/S0015-0282(16)42801-3))
- Macias Garcia B, Gonzalez Fernandez L, Loux SC, Rocha AM, Guimaraes T, Pena FJ, Varner DD & Hinrichs K 2015 Effect of calcium, bicarbonate and albumin on capacitation-related events in equine sperm. *Reproduction* **149** 87–99. (<https://doi.org/10.1530/REP-14-0457>)

- Mahony MC & Gwathmey T 1999 Protein tyrosine phosphorylation during hyperactivated motility of cynomolgus monkey (*Macaca fascicularis*) spermatozoa. *Biology of Reproduction* **60** 1239–1243. (<https://doi.org/10.1095/biolreprod60.5.1239>)
- Mandal A, Naaby-Hansen S, Wolkowicz MJ, Klotz K, Shetty J, Retief JD, Coonrod SA, Kinter M, Sherman N, Cesar F *et al.* 1999 FSP95, a testis-specific 95-kilodalton fibrous sheath antigen that undergoes tyrosine phosphorylation in capacitated human spermatozoa. *Biology of Reproduction* **61** 1184–1197. (<https://doi.org/10.1095/biolreprod61.5.1184>)
- Mann T 1975 Biochemistry of stallion semen. *Journal of Reproduction and Fertility Supplement* 47–52.
- Marin-Briggiler CI, Jha KN, Chertihin O, Buffone MG, Herr JC, Vazquez-Levin MH & Visconti PE 2005 Evidence of the presence of calcium/calmodulin-dependent protein kinase IV in human sperm and its involvement in motility regulation. *Journal of Cell Science* **118** 2013–2022. (<https://doi.org/10.1242/jcs.02326>)
- Marquez B & Suarez SS 2004 Different signaling pathways in bovine sperm regulate capacitation and hyperactivation. *Biology of Reproduction* **70** 1626–1633. (<https://doi.org/10.1095/biolreprod.103.026476>)
- Marquez B & Suarez SS 2007 Bovine sperm hyperactivation is promoted by alkaline-stimulated Ca²⁺ influx. *Biology of Reproduction* **76** 660–665. (<https://doi.org/10.1095/biolreprod.106.055038>)
- McPartlin LA, Littell J, Mark E, Nelson JL, Travis AJ & Bedford-Guaus SJ 2008 A defined medium supports changes consistent with capacitation in stallion sperm, as evidenced by increases in protein tyrosine phosphorylation and high rates of acrosomal exocytosis. *Theriogenology* **69** 639–650. (<https://doi.org/10.1016/j.theriogenology.2007.11.016>)
- McPartlin LA, Suarez SS, Czaya CA, Hinrichs K & Bedford-Guaus SJ 2009 Hyperactivation of stallion sperm is required for successful *in vitro* fertilization of equine oocytes. *Biology of Reproduction* **81** 199–206. (<https://doi.org/10.1095/biolreprod.108.074880>)
- McPartlin LA, Visconti PE & Bedford-Guaus SJ 2011 Guanine-nucleotide exchange factors (RAPGEF3/RAPGEF4) induce sperm membrane depolarization and acrosomal exocytosis in capacitated stallion sperm. *Biology of Reproduction* **85** 179–188. (<https://doi.org/10.1095/biolreprod.110.085555>)
- Meisel S 2004 The sperm, a neuron with a tail: 'neuronal' receptors in mammalian sperm. *Biological Reviews of the Cambridge Philosophical Society* **79** 713–732. (<https://doi.org/10.1017/S1464793103006407>)
- Melendrez CS & Meisel S 1995 Studies of porcine and human sperm suggesting a role for a sperm glycine receptor/Cl⁻ channel in the zona pellucida-initiated acrosome reaction. *Biology of Reproduction* **53** 676–683. (<https://doi.org/10.1095/biolreprod53.3.676>)
- Meyers SA, Liu IK, Overstreet JW, Vadas S & Drobnis EZ 1996 Zona pellucida binding and zona-induced acrosome reactions in horse spermatozoa: comparisons between fertile and subfertile stallions. *Theriogenology* **46** 1277–1288. ([https://doi.org/10.1016/S0093-691X\(96\)00299-3](https://doi.org/10.1016/S0093-691X(96)00299-3))
- Mugnier S, Dell'Aquila ME, Pelaez J, Douet C, Ambruosi B, De Santis T, Lacalandra GM, Lebos C, Sizaret PY, Delaleu B *et al.* 2009a New insights into the mechanisms of fertilization: comparison of the fertilization steps, composition, and structure of the zona pellucida between horses and pigs. *Biology of Reproduction* **81** 856–870. (<https://doi.org/10.1095/biolreprod.109.077651>)
- Mugnier S, Kervella M, Douet C, Canepa S, Pascal G, Deleuze S, Duchamp G, Monget P & Goudet G 2009b The secretions of oviduct epithelial cells increase the equine *in vitro* fertilization rate: are osteopontin, atrial natriuretic peptide A and oviductin involved? *Reproductive Biology and Endocrinology* **7** 129. (<https://doi.org/10.1186/1477-7827-7-129>)
- Mujica A, Neri-Bazan L, Tash JS & Uribe S 1994 Mechanism for procaine-mediated hyperactivated motility in guinea pig spermatozoa. *Molecular Reproduction and Development* **38** 285–292. (<https://doi.org/10.1002/mrd.1080380309>)
- Mukai C & Okuno M 2004 Glycolysis plays a major role for adenosine triphosphate supplementation in mouse sperm flagellar movement. *Biology of Reproduction* **71** 540–547. (<https://doi.org/10.1095/biolreprod.103.026054>)
- Naaby-Hansen S, Wolkowicz MJ, Klotz K, Bush LA, Westbrook VA, Shibahara H, Shetty J, Coonrod SA, Reddi PP, Shannon J *et al.* 2001 Colocalization of the inositol 1,4,5-trisphosphate receptor and calreticulin in the equatorial segment and in membrane bounded vesicles in the cytoplasmic droplet of human spermatozoa. *Molecular Human Reproduction* **7** 923–933. (<https://doi.org/10.1093/molehr/7.10.923>)
- Navarro B, Kirichok Y & Clapham DE 2007 K_{Sper}, a pH-sensitive K⁺ current that controls sperm membrane potential. *PNAS* **104** 7688–7692. (<https://doi.org/10.1073/pnas.0702018104>)
- Navarro B, Kirichok Y, Chung JJ & Clapham DE 2008 Ion channels that control fertility in mammalian spermatozoa. *International Journal of Developmental Biology* **52** 607–613. (<https://doi.org/10.1387/ijdb.072554bn>)
- Olds-Clarke P 1989 Sperm from tw32/+ mice: capacitation is normal, but hyperactivation is premature and nonhyperactivated sperm are slow. *Developmental Biology* **131** 475–482. ([https://doi.org/10.1016/S0012-1606\(89\)80018-1](https://doi.org/10.1016/S0012-1606(89)80018-1))
- Olson SD, Suarez SS & Fauci LJ 2010 A model of CatSper channel mediated calcium dynamics in mammalian spermatozoa. *Bulletin of Mathematical Biology* **72** 1925–1946. (<https://doi.org/10.1007/s11538-010-9516-5>)
- O'Rand MG & Fisher SJ 1987 Localization of zona pellucida binding sites on rabbit spermatozoa and induction of the acrosome reaction by solubilized zonae. *Developmental Biology* **119** 551–559. ([https://doi.org/10.1016/0012-1606\(87\)90058-3](https://doi.org/10.1016/0012-1606(87)90058-3))
- Ortega-Ferrusola C, Anel-Lopez L, Martin-Munoz P, Ortiz-Rodriguez JM, Gil MC, Alvarez M, de Paz P, Ezquerro LJ, Masot AJ, Redondo E *et al.* 2017 Computational flow cytometry reveals that cryopreservation induces spermatosis but subpopulations of spermatozoa may experience capacitation-like changes. *Reproduction* **153** 293–304. (<https://doi.org/10.1530/REP-16-0539>)
- Ortega Ferrusola C, Gonzalez Fernandez L, Morrell JM, Salazar Sandoval C, Macias Garcia B, Rodriguez-Martinez H, Tapia JA & Pena FJ 2009 Lipid peroxidation, assessed with BODIPY-C11, increases after cryopreservation of stallion spermatozoa, is stallion-dependent and is related to apoptotic-like changes. *Reproduction* **138** 55–63. (<https://doi.org/10.1530/REP-08-0484>)
- Osheroff JE, Visconti PE, Valenzuela JP, Travis AJ, Alvarez J & Kopf GS 1999 Regulation of human sperm capacitation by a cholesterol efflux-stimulated signal transduction pathway leading to protein kinase A-mediated up-regulation of protein tyrosine phosphorylation. *Molecular Human Reproduction* **5** 1017–1026. (<https://doi.org/10.1093/molehr/5.11.1017>)
- Palmer E, Bezard J, Magistrini M & Duchamp G 1991 *In vitro* fertilization in the horse. A retrospective study. *Journal of Reproduction and Fertility Supplement* **44** 375–384.
- Parrish JJ, Susko-Parrish J, Winer MA & First NL 1988 Capacitation of bovine sperm by heparin. *Biology of Reproduction* **38** 1171–1180. (<https://doi.org/10.1095/biolreprod38.5.1171>)
- Perry G 2013 2012 statistics of embryo collection and transfer in domestic farm animals. *Embryo Transfer Newsletter* **31** 1–67.
- Plaza Davila M, Martin Munoz P, Tapia JA, Ortega Ferrusola C, Balao da Silva CC & Pena FJ 2015 Inhibition of mitochondrial Complex I leads to decreased motility and membrane integrity related to increased hydrogen peroxide and reduced ATP production, while the inhibition of glycolysis has less impact on sperm motility. *PLoS One* **10** e0138777. (<https://doi.org/10.1371/journal.pone.0138777>)
- Pommer AC, Rutllant J & Meyers SA 2003 Phosphorylation of protein tyrosine residues in fresh and cryopreserved stallion spermatozoa under capacitating conditions. *Biology of Reproduction* **68** 1208–1214. (<https://doi.org/10.1095/biolreprod.102.011106>)
- Qi H, Moran MM, Navarro B, Chong JA, Krapivinsky G, Krapivinsky L, Kirichok Y, Ramsey IS, Quill TA & Clapham DE 2007 All four CatSper ion channel proteins are required for male fertility and sperm cell hyperactivated motility. *PNAS* **104** 1219–1223. (<https://doi.org/10.1073/pnas.0610286104>)
- Quill TA, Sugden SA, Rossi KL, Doolittle LK, Hammer RE & Garbers DL 2003 Hyperactivated sperm motility driven by CatSper2 is required for fertilization. *PNAS* **100** 14869–14874. (<https://doi.org/10.1073/pnas.2136654100>)
- Rathi R, Colenbrander B, Bevers MM & Gadella BM 2001 Evaluation of *in vitro* capacitation of stallion spermatozoa. *Biology of Reproduction* **65** 462–470. (<https://doi.org/10.1095/biolreprod65.2.462>)
- Rathi R, Colenbrander B, Stout TA, Bevers MM & Gadella BM 2003 Progesterone induces acrosome reaction in stallion spermatozoa via a protein tyrosine kinase dependent pathway. *Molecular Reproduction and Development* **64** 120–128. (<https://doi.org/10.1002/mrd.10216>)

- Ren D, Navarro B, Perez G, Jackson AC, Hsu S, Shi Q, Tilly JL & Clapham DE 2001 A sperm ion channel required for sperm motility and male fertility. *Nature* **413** 603–609. (<https://doi.org/10.1038/35098027>)
- Roasa LM, Choi YH, Love CC, Romo S, Varner DD & Hinrichs K 2007 Ejaculate and type of freezing extender affect rates of fertilization of horse oocytes *in vitro*. *Theriogenology* **68** 560–566. (<https://doi.org/10.1016/j.theriogenology.2007.04.057>)
- Roggero CM, De Blas GA, Dai H, Tomes CN, Rizo J & Mayorga LS 2007 Complexin/synaptotagmin interplay controls acrosomal exocytosis. *Journal of Biological Chemistry* **282** 26335–26343. (<https://doi.org/10.1074/jbc.M700854200>)
- Saaranen MJ, Calvo L, Dennison L, Banks S, Bustillo M, Dorfmann AD, Goldstein M, Thorsell L, Schulman JD & Sherins RJ 1993 Acrosome reaction inducing activity in follicular fluid correlates with progesterone concentration but not with oocyte maturity or fertilizability. *Human Reproduction* **8** 1448–1454. (<https://doi.org/10.1093/oxfordjournals.humrep.a138277>)
- Schlingmann K, Michaut MA, McElwee JL, Wolff CA, Travis AJ & Turner RM 2007 Calmodulin and CaMKII in the sperm principal piece: evidence for a motility-related calcium/calmodulin pathway. *Journal of Andrology* **28** 706–716. (<https://doi.org/10.2164/jandrol.106.001669>)
- Shingyoji C, Murakami A & Takahashi K 1977 Local reactivation of Triton-extracted flagella by ionophoretic application of ATP. *Nature* **265** 269–270. (<https://doi.org/10.1038/265269a0>)
- Si Y 1999 Hyperactivation of hamster sperm motility by temperature-dependent tyrosine phosphorylation of an 80-kDa protein. *Biology of Reproduction* **61** 247–252. (<https://doi.org/10.1095/biolreprod61.1.247>)
- Si Y & Okuno M 1999 Role of tyrosine phosphorylation of flagellar proteins in hamster sperm hyperactivation. *Biology of Reproduction* **61** 240–246. (<https://doi.org/10.1095/biolreprod61.1.240>)
- Si Y & Olds-Clarke P 2000 Evidence for the involvement of calmodulin in mouse sperm capacitation. *Biology of Reproduction* **62** 1231–1239. (<https://doi.org/10.1095/biolreprod62.5.1231>)
- Signorelli J, Diaz ES & Morales P 2012 Kinases, phosphatases and proteases during sperm capacitation. *Cell and Tissue Research* **349** 765–782. (<https://doi.org/10.1007/s00441-012-1370-3>)
- Simons K & Toomre D 2000 Lipid rafts and signal transduction. *Nature Reviews. Molecular Cell Biology* **1** 31–39. (<https://doi.org/10.1038/35036052>)
- Smith GD, Wolf DP, Trautman KC, da Cruz e Silva EF, Greengard P & Vijayaraghavan S 1996 Primate sperm contain protein phosphatase 1, a biochemical mediator of motility. *Biology of Reproduction* **54** 719–727. (<https://doi.org/10.1095/biolreprod54.3.719>)
- Stauss CR, Votta TJ & Suarez SS 1995 Sperm motility hyperactivation facilitates penetration of the hamster zona pellucida. *Biology of Reproduction* **53** 1280–1285. (<https://doi.org/10.1095/biolreprod53.6.1280>)
- Stephoe PC & Edwards RG 1978 Birth after the reimplantation of a human embryo. *Lancet* **2** 366. ([https://doi.org/10.1016/S0140-6736\(78\)92957-4](https://doi.org/10.1016/S0140-6736(78)92957-4))
- Storey BT 2008 Mammalian sperm metabolism: oxygen and sugar, friend and foe. *International Journal of Developmental Biology* **52** 427–437. (<https://doi.org/10.1387/ijdb.072522bs>)
- Suarez SS 2002 Formation of a reservoir of sperm in the oviduct. *Reproduction in Domestic Animals* **37** 140–143. (<https://doi.org/10.1046/j.1439-0531.2002.00346.x>)
- Suarez SS 2008 Control of hyperactivation in sperm. *Human Reproduction Update* **14** 647–657. (<https://doi.org/10.1093/humupd/dmn029>)
- Suarez SS & Dai X 1992 Hyperactivation enhances mouse sperm capacity for penetrating viscoelastic media. *Biology of Reproduction* **46** 686–691. (<https://doi.org/10.1095/biolreprod46.4.686>)
- Suarez SS & Pacey AA 2006 Sperm transport in the female reproductive tract. *Human Reproduction Update* **12** 23–37. (<https://doi.org/10.1093/humupd/dmi047>)
- Suarez SS, Vincenti L & Ceglia MW 1987 Hyperactivated motility induced in mouse sperm by calcium ionophore A23187 is reversible. *Journal of Experimental Zoology* **244** 331–336. (<https://doi.org/10.1002/jez.1402440218>)
- Suarez SS, Varosi SM & Dai X 1993 Intracellular calcium increases with hyperactivation in intact, moving hamster sperm and oscillates with the flagellar beat cycle. *PNAS* **90** 4660–4664. (<https://doi.org/10.1073/pnas.90.10.4660>)
- Takakura K, Beckman JS, MacMillan-Crow LA & Crow JP 1999 Rapid and irreversible inactivation of protein tyrosine phosphatases PTP1B, CD45, and LAR by peroxynitrite. *Archives of Biochemistry and Biophysics* **369** 197–207. (<https://doi.org/10.1006/abbi.1999.1374>)
- Tapia JC, Espinoza F & Aguayo LG 1997 Differential intracellular regulation of cortical GABA(A) and spinal glycine receptors in cultured neurons. *Brain Research* **769** 203–210. ([https://doi.org/10.1016/S0006-8993\(97\)00672-0](https://doi.org/10.1016/S0006-8993(97)00672-0))
- Tash JS 1989 Protein phosphorylation: the second messenger signal transducer of flagellar motility. *Cell Motility and the Cytoskeleton* **14** 332–339. (<https://doi.org/10.1002/cm.970140303>)
- Tash JS & Means AR 1983 Cyclic adenosine 3',5' monophosphate, calcium and protein phosphorylation in flagellar motility. *Biology of Reproduction* **28** 75–104. (<https://doi.org/10.1095/biolreprod28.1.75>)
- Tash JS & Bracho GE 1994 Regulation of sperm motility: emerging evidence for a major role for protein phosphatases. *Journal of Andrology* **15** 505–509.
- Tash JS, Krinks M, Patel J, Means RL, Klee CB & Means AR 1988 Identification, characterization, and functional correlation of calmodulin-dependent protein phosphatase in sperm. *Journal of Cell Biology* **106** 1625–1633. (<https://doi.org/10.1083/jcb.106.5.1625>)
- Tateno H, Kräpf D, Hino T, Sanchez-Cardenas C, Darszon A, Yanagimachi R & Visconti PE 2013 Ca²⁺ ionophore A23187 can make mouse spermatozoa capable of fertilizing *in vitro* without activation of cAMP-dependent phosphorylation pathways. *PNAS* **110** 18543–18548. (<https://doi.org/10.1073/pnas.1317113110>)
- Tremoleda JL, Gadella BM, Stout T, Colenbrander B & Bevers MM 2003 *Evaluation of Sperm-Oocyte Interaction During in Vitro Fertilization in the Horse*. Utrecht University, pp. 67–91.
- Trevino CL, Santi CM, Beltran C, Hernandez-Cruz A, Darszon A & Lomeli H 1998 Localisation of inositol trisphosphate and ryanodine receptors during mouse spermatogenesis: possible functional implications. *Zygote* **6** 159–172. (<https://doi.org/10.1017/S0967199498000094>)
- Tsai PS, Garcia-Gil N, van Haeften T & Gadella BM 2010 How pig sperm prepares to fertilize: stable acrosome docking to the plasma membrane. *PLoS One* **5** e11204. (<https://doi.org/10.1371/journal.pone.0011204>)
- Tsai PS, Brewis IA, van Maaren J & Gadella BM 2012 Involvement of complexin 2 in docking, locking and unlocking of different SNARE complexes during sperm capacitation and induced acrosomal exocytosis. *PLoS One* **7** e32603. (<https://doi.org/10.1371/journal.pone.0032603>)
- Turner RM 2006 Moving to the beat: a review of mammalian sperm motility regulation. *Reproduction, Fertility, and Development* **18** 25–38. (<https://doi.org/10.1071/RD05120>)
- van Gestel RA, Brewis IA, Ashton PR, Helms JB, Brouwers JF & Gadella BM 2005a Capacitation-dependent concentration of lipid rafts in the apical ridge head area of porcine sperm cells. *Molecular Human Reproduction* **11** 583–590. (<https://doi.org/10.1093/molehr/gah200>)
- van Gestel RA, Helms JB, Brouwers JF & Gadella BM 2005b Effects of methyl-beta-cyclodextrin-mediated cholesterol depletion in porcine sperm compared to somatic cells. *Molecular Reproduction and Development* **72** 386–395. (<https://doi.org/10.1002/mrd.20351>)
- van Gestel RA, Brewis IA, Ashton PR, Brouwers JF & Gadella BM 2007 Multiple proteins present in purified porcine sperm apical plasma membranes interact with the zona pellucida of the oocyte. *Molecular Human Reproduction* **13** 445–454. (<https://doi.org/10.1093/molehr/gam030>)
- Vigil P 1989 Gamete membrane fusion in hamster spermatozoa with reacted equatorial segment. *Gamete Research* **23** 203–213. (<https://doi.org/10.1002/mrd.1120230207>)
- Vijayaraghavan S, Stephens DT, Trautman K, Smith GD, Khatra B, da Cruz e Silva EF & Greengard P 1996 Sperm motility development in the epididymis is associated with decreased glycogen synthase kinase-3 and protein phosphatase 1 activity. *Biology of Reproduction* **54** 709–718. (<https://doi.org/10.1095/biolreprod54.3.709>)
- Visconti PE, Bailey JL, Moore GD, Pan D, Olds-Clarke P & Kopf GS 1995a Capacitation of mouse spermatozoa. I. Correlation between the capacitation state and protein tyrosine phosphorylation. *Development* **121** 1129–1137.
- Visconti PE, Moore GD, Bailey JL, Leclerc P, Connors SA, Pan D, Olds-Clarke P & Kopf GS 1995b Capacitation of mouse spermatozoa. II.

- Protein tyrosine phosphorylation and capacitation are regulated by a cAMP-dependent pathway. *Development* **121** 1139–1150.
- Visconti PE, Johnson LR, Oyaski M, Fornes M, Moss SB, Gerton GL & Kopf GS** 1997 Regulation, localization, and anchoring of protein kinase A subunits during mouse sperm capacitation. *Developmental Biology* **192** 351–363. (<https://doi.org/10.1006/dbio.1997.8768>)
- Visconti PE, Galantino-Homer H, Ning X, Moore GD, Valenzuela JP, Jorgez CJ, Alvarez JG & Kopf GS** 1999a Cholesterol efflux-mediated signal transduction in mammalian sperm. *Journal of Biological Chemistry* **274** 3235–3242. (<https://doi.org/10.1074/jbc.274.5.3235>)
- Visconti PE, Stewart-Savage J, Blasco A, Battaglia L, Miranda P, Kopf GS & Tezon JG** 1999b Roles of bicarbonate, cAMP, and protein tyrosine phosphorylation on capacitation and the spontaneous acrosome reaction of hamster sperm. *Biology of Reproduction* **61** 76–84. (<https://doi.org/10.1095/biolreprod61.1.76>)
- Vjugina U & Evans JP** 2008 New insights into the molecular basis of mammalian sperm-egg membrane interactions. *Frontiers in Bioscience* **13** 462–476. (<https://doi.org/10.2741/2693>)
- Wargo MJ & Smith EF** 2003 Asymmetry of the central apparatus defines the location of active microtubule sliding in *Chlamydomonas* flagella. *PNAS* **100** 137–142. (<https://doi.org/10.1073/pnas.0135800100>)
- Watson PF** 2000 The causes of reduced fertility with cryopreserved semen. *Animal Reproduction Science* **60–61** 481–492. ([https://doi.org/10.1016/S0378-4320\(00\)00099-3](https://doi.org/10.1016/S0378-4320(00)00099-3))
- Weaver CD, Partridge JG, Yao TL, Moates JM, Magnuson MA & Verdoorn TA** 1998 Activation of glycine and glutamate receptors increases intracellular calcium in cells derived from the endocrine pancreas. *Molecular Pharmacology* **54** 639–646.
- White DR & Aitken RJ** 1989 Relationship between calcium, cyclic AMP, ATP, and intracellular pH and the capacity of hamster spermatozoa to express hyperactivated motility. *Gamete Research* **22** 163–177. (<https://doi.org/10.1002/mrd.1120220205>)
- Wiesner B, Weiner J, Middendorff R, Hagen V, Kaupp UB & Weyand I** 1998 Cyclic nucleotide-gated channels on the flagellum control Ca²⁺ entry into sperm. *Journal of Cell Biology* **142** 473–484. (<https://doi.org/10.1083/jcb.142.2.473>)
- Yanagimachi R** 1994 Mammalian fertilization. In *The Physiology of Reproduction*, pp. 189–317. Eds JD Neill & E Knobil. New York: Raven Press.
- Zeng Y, Oberdorf JA & Florman HM** 1996 pH regulation in mouse sperm: identification of Na(+)-, Cl(-), and HCO₃(-)-dependent and arylaminobenzoate-dependent regulatory mechanisms and characterization of their roles in sperm capacitation. *Developmental Biology* **173** 510–520. (<https://doi.org/10.1006/dbio.1996.0044>)
- Zhang JJ, Muzs LZ & Boyle MS** 1990 *In vitro* fertilization of horse follicular oocytes matured *in vitro*. *Molecular Reproduction and Development* **26** 361–365. (<https://doi.org/10.1002/mrd.1080260411>)

Received 15 October 2018

First decision 10 December 2018

Revised manuscript received 25 December 2018

Accepted 4 February 2019