ORALLY ACTIVE OESTROGENS AND PROGESTINS IN PREVENTION OF PREGNANCY IN RATS

U. K. BANIK, C. REVESZ AND F. HERR

Ayerst Laboratories, Montreal, Canada

(Received 20th May 1968, revised 17th September 1968)

Summary. Orally active synthetic oestrogens and progestins were studied in tests to prevent pregnancy in rats. By daily administration the contraceptive effect of ethynyloestradiol and a new oestrogen, 17 α -[3-furyl]-estra-1,3,5(10),7-tetraene-3,17-diol 3-acetate (AY-11483), was potentiated by three progestins, namely, chlormadinone acetate, 6-chloro-3 β ,17 α -dihydroxypregn-4,6-dien-20-one diacetate (AY-11440) and medrogestone. AY-11483 given alone, or together with AY-11440, was more potent in preventing ovulation and implantation than ethynyl-oestradiol alone or in combination with other compounds. Chlormadinone acetate or AY-11440 alone were completely inactive in all the assays. The mode of action of oestrogen or oestrogen and progestin in the prevention of ovulation and implantation is discussed. In the rat, implantation seems more vulnerable to oestrogens than does ovulation.

INTRODUCTION

Recently, Diczfalusy (1968) reviewed the use of various oestrogens and progestins and their possible mode of action in blocking pregnancy both in humans and laboratory animals. The present paper reports experiments on the prevention of pregnancy in rats in which a series of tests was used to study a new oestrogen, AY-11483, and to compare its effects with other known oestrogens and progestins. The chemical and biological properties of AY-11483 were reported earlier (Lefebvre, Marshall, Revesz, Banik & Deghenghi, 1967).

MATERIALS AND METHODS

All experiments were performed on cyclic adult (180 to 250 g) Sprague-Dawley rats. Standard (Purina) diet and water were given unrationed and animals were kept in uniform conditions with exposure to 12 hr light and 12 hr dark. All compounds were dissolved or suspended in sesame oil and administered orally by stomach tube. Doses are expressed as the weight of compounds administered to each rat. Unless otherwise stated, treatments were started on the day of pro-oestrus of the normal cycle. Day 1 of pregnancy was designated as the day when either a vaginal plug or spermatozoa were found after placing overnight with males known to be fertile. Statistical analysis of the data from

rats with ova or implantations was made using the four-fold contingency test tables of Mainland & Murray (1952); the average number of ova or implantation sites in each rat in the experimental series was compared with those of the controls by using Student's *t* test. The structural formulae of different oestrogens and progestins used in these studies are presented in Text-fig. 1.

TEXT-FIG. 1. Structural formulae of oestrogens and progestins used in prevention of pregnancy in rats.

Daily administration of oestrogens and progestins

Groups of ten female rats were given the test compound every day for 35 days during cohabitation with males. Two males were put in each cage containing five females which were checked every morning for evidence of mating. Mated animals were killed on Day 9 or 10 of pregnancy and the number of implantations counted. All other animals were killed 10 days after the cohabitation period of 35 days and uteri examined for implantation sites.

Anti-ovulatory effect of oestrogens and progestins given alone or in different combinations

The test compound was administered to 4-day cyclic rats, either 1 day before pro-oestrus or on each day starting on the morning of oestrus until the next

pro-oestrus. The vaginal cytology was studied every morning between 09.00 and 10.00 hours. Treated and control animals were killed on the day of post-oestrous after treatment. The oviduct and uterus were examined for the presence of ova.

Anti-implantation effect of oestrogens and progestins given alone or in different combinations

In these experiments, all compounds were tested, in rats after mating, by the method of Banik & Pincus (1962). As before, different oestrogens were given alone or in combination with a progestin on Day 1, or on Days 1 to 5 of pregnancy. All animals were killed on Day 9 or 10 of pregnancy and the number of implantations was counted.

RESULTS

Daily administration of oestrogens and progestins

Results shown in Table 1 reveal that none of the oestrogens alone (10 µg/rat, Groups 2, 3 and 4) and none of the progestins alone (100 µg/rat, Groups 5, 6 and 7) were completely effective in preventing pregnancy when given over a period of 35 days. A marginally significant reduction of implantations was

TABLE 1
DAILY ORAL ADMINISTRATION OF OESTROGENS AND PROGESTINS IN PREVENTION OF PREGNANCY*

Group	Oestrogen (10 µg/rat)	Progestin (100 µg/rat)	No. of rats pregnant	Implantations (mean ± S.E.)
1	Oil control	Oil control	9/10	10.7 ± 0.55
2	Ethynyloestradiol	—	7/10	7.4 ± 1.65
3	AY-11483	—	6/10	6.5 ± 0.99†
4	Mestranol	—	8/10	9.5 ± 0.60
5	—	Chlormadinone acetate	9/10	12.3 ± 0.73
6	—	AY-11440	10/10	10.1 ± 1.26
7	—	Medrogestone	8/10	9.6 ± 0.24
8	Ethynyloestradiol	Chlormadinone acetate	0/10†	0†
9	Mestranol	Chlormadinone acetate	8/10	9.7 ± 0.86
10	AY-11483	AY-11440	0/10†	0†
11	AY-11483	Medrogestone	0/10†	0†
12	AY-11483	Chlormadinone acetate	0/12†	0†

* Treatment was continued daily for 35 days.

† Significantly different from control, $P < 0.01$.

found only in the group treated with AY-11483. However, combinations of ethynyloestradiol and chlormadinone acetate, AY-11483 and AY-11440, AY-11483 and medrogestone, and AY-11483 and chlormadinone acetate were all significantly effective in preventing pregnancy (Groups 8, 10, 11 and 12). However, neither the oestrous cycle nor the mating behaviour of these animals was disturbed. The mestranol-chlormadinone acetate combination was inactive in this test (Group 9).

Anti-ovulatory effect of oestrogens and progestins given alone or in different combinations

All ova were collected from the oviduct and none was found in the uterus. The data in Table 2 show that neither ethynyloestradiol nor AY-11483 (10 or

100 µg/rat, Groups 2 to 5) given 1 day before pro-oestrus, was effective in preventing ovulation. These oestrogens were only effective if given at the 100 µg dose level over the entire cycle (Table 3); at the 10 µg dose level AY-11483 was significantly more potent than ethynyloestradiol. Neither chlormadinone acetate nor AY-11440 given either 1 day before pro-oestrus (Table 2) or during

TABLE 2
ANTI-OVULATORY EFFECT OF OESTROGENS AND PROGESTINS GIVEN ALONE OR IN DIFFERENT COMBINATIONS*

Group	Oestrogen (µg/rat)	Progestin (100 µg/rat)	No. of rats that ovulated	No. of ova (mean ± S.E.)
1	Oil control	Oil control	13/13	11.7 ± 0.60
2	Ethynyloestradiol (10)	—	6/6	10.5 ± 1.21
3	Ethynyloestradiol (100)	—	10/12	11.9 ± 1.4
4	AY-11483 (10)	—	5/6	10.7 ± 0.94
5	AY-11483 (100)	—	11/11	11.7 ± 0.34
6	—	Chlormadinone acetate	6/6	11.0 ± 0.48
7	—	AY-11440	6/6	10.5 ± 0.89
8	Ethynyloestradiol (10)	Chlormadinone acetate	5/5	8.6 ± 1.44†
9	AY-11483 (10)	Chlormadinone acetate	6/6	9.1 ± 1.11
10	Ethynyloestradiol (10)	AY-11440	5/6	10.2 ± 1.50
11	AY-11483 (10)	AY-11440	5/6	10.2 ± 0.50

* Treatment given 1 day before pro-oestrus.

† Significantly different from control, $P < 0.05$.

TABLE 3
ANTI-OVULATORY EFFECT OF OESTROGENS AND PROGESTINS GIVEN ALONE OR IN DIFFERENT COMBINATIONS*

Group	Oestrogen (µg/rat)	Progestin (100 µg/rat)	No. of rats that ovulated	No. of ova (mean ± S.E.)
1	Oil control	Oil control	13/13	11.7 ± 0.60
2	Ethynyloestradiol (10)	—	6/8	10.1 ± 0.40
3	Ethynyloestradiol (100)	—	2/7†	11.0 ± 0.0
4	AY-11483 (10)	—	2/8†	13.5 ± 0.50
5	AY-11483 (100)	—	0/7‡	0.0‡
6	—	Chlormadinone acetate	9/10	10.7 ± 0.83
7	—	AY-11440	10/10	10.1 ± 0.95
8	Ethynyloestradiol (10)	Chlormadinone acetate	8/14	6.5 ± 0.68‡
9	AY-11483 (10)	Chlormadinone acetate	10/20	10.0 ± 1.04
10	Ethynyloestradiol (10)	AY-11440	9/13	10.4 ± 0.91
11	AY-11483 (10)	AY-11440	5/20‡	9.2 ± 1.11‡

* Treatment given once daily from one post-oestrus to the next pro-oestrus; not more than 4 days.

† Significantly different from control, $P < 0.05$.

‡ Significantly different from control, $P < 0.01$.

the whole of the cycle (Table 3) prevented ovulation. Furthermore, 10 µg of either of these oestrogens plus 100 µg of either of these progestins, when given in a single dose 1 day before pro-oestrus, was found inactive (Table 2). Partial inhibition of ovulation was achieved with the oestrogen-progestin combinations

given over the entire length of the cycle (Table 3). Of these treatments, it seems that 10 μg of AY-11483 alone or in combination with 100 μg of AY-11440 (Table 3) was the most potent in this test.

Anti-implantation effect of different oestrogens given alone or in combination with a progestin

The results presented in Table 4 show that 10 μg of either ethynyloestradiol or AY-11483 is partially effective in preventing the implantation of ova in mated rats when administered over a period of 5 days (Groups 2 and 4). These two oestrogens were more effective if given in doses of 100 μg on Day 1

TABLE 4
ANTI-IMPLANTATION EFFECT OF DIFFERENT OESTROGENS GIVEN ALONE OR IN COMBINATION WITH A PROGESTIN*

Group	Oestrogen ($\mu\text{g}/\text{rat}$)	Progestin (mg/rat)	Days of treatment after mating	No. of rats with implantations	Implantations (mean $\pm$ S.E.)
1	Oil control	Oil control	1 to 5	11/12	12.2 $\pm$ 0.65
2	Ethynyloestradiol (10)	—	1 to 5	7/9	7.4 $\pm$ 1.64†
3	Ethynyloestradiol (100)	—	1	2/10‡	11.0 $\pm$ 0
4	AY-11483 (10)	—	1 to 5	4/9†	7.0 $\pm$ 1.22‡
5	AY-11483 (100)	—	1	0/6‡	0.0 $\pm$ 0.0‡
6	Mestranol (100)	—	1	7/11	9.4 $\pm$ 1.41
7	—	Chlormadinone acetate (0.1)	1 to 5	8/9	9.7 $\pm$ 0.58†
8	—	Chlormadinone acetate (2.5)	1	5/6	12.3 $\pm$ 0.62
9	—	AY-11440 (0.1)	1 to 5	6/9	13.1 $\pm$ 0.87
10	—	AY-11440 (2.5)	1	5/5	13.2 $\pm$ 0.48
11	Ethynyloestradiol (10)	Chlormadinone acetate (0.1)	1 to 5	5/5	11.0 $\pm$ 0.31
12	AY-11483 (10)	Chlormadinone acetate (0.1)	1 to 5	3/9†	3.6 $\pm$ 1.20‡
13	Ethynyloestradiol (10)	AY-11440 (0.1)	1 to 5	4/8	9.2 $\pm$ 0.75†
14	AY-11483 (10)	AY-11440 (0.1)	1 to 5	0/6‡	0‡

* Treatment given orally in mated rats during the pre-implantation period.

† Significantly different from control, $P < 0.05$.

‡ Significantly different from control, $P < 0.01$.

of pregnancy (Groups 3 and 5). Mestranol (100 μg) given on Day 1 of pregnancy (Group 6) was not significantly effective in the inhibition of implantation. AY-11483 was more potent than ethynyloestradiol or mestranol in this test. When ethynyloestradiol (10 μg) was given in combination with chlormadinone acetate (100 μg) for 5 days, implantation was not inhibited (Group 11); however, corresponding doses of AY-11483 and chlormadinone acetate gave a significant inhibition of implantation (Group 12). When ethynyloestradiol (10 μg) was given with AY-11440 (100 μg), only a partial inhibition of implantation was achieved (Group 13), but treatment with the same quantity of AY-11483 together with AY-11440 resulted in the complete inhibition of implantation (Group 14). In a subsequent study, we failed to collect ova on Day 5 of pregnancy from the reproductive tract of rats treated daily with AY-11483 (10 μg) and AY-11440 (100 μg) during the preceding 4 days.

Chlormadinone acetate or AY-11440 given in doses up to 2.5 mg was inactive in preventing implantation (Groups 7 to 10).

DISCUSSION

The results presented in this paper demonstrate the anti-fertility effect of a new orally active oestrogen AY-11483 given alone in moderately high doses or in combination with a progestin in lower doses. The new oestrogen was more effective than various other oestrogens given in similar doses (Tables 3 and 4). It is pertinent to point out here that we have found AY-11483 to be more active than mestranol or ethynyloestradiol as percentage of the median cornifying dose in rats (Revesz, Banik & Herr, unpublished observations). The effect, in preventing pregnancy, of continuous treatment over 35 days with AY-11483 or ethynyloestradiol (Table 1) was potentiated by AY-11440 and chlormadinone acetate, perhaps by interfering with both ovulation and implantation. It was also noted that the anti-implantation effect of AY-11483 was potentiated more by AY-11440 than by chlormadinone acetate (Table 4), although the progestational activity of these two progestins in rabbits is similar (Revesz, Banik & Herr, 1967). Synergism in anti-ovulatory tests between ethynyloestradiol and chlormadinone acetate in rabbits (Chambon & Veve, 1966) and between ethynyloestradiol and norgestrel in rats (Edgren, Jones & Peterson, 1967) has been reported before and was confirmed in these experiments.

Data obtained from the anti-ovulatory assay (Tables 2 and 3) revealed that a prolonged treatment with oestrogen was more effective in preventing ovulation than the single dose. Oestrogens could conceivably inhibit the release of both FSH and LH (Dorfman & Dorfman, 1963; McCann & Talecnik, 1961) and thus prevent ovulation. The other possibility is that oestrogens maintain the life of the corpus luteum. Thus, the progesterone produced by the corpora lutea may act independently, or together with the oestrogen to inhibit the release of LH and block ovulation (Watnick, Gibson, Vinegra & Tolksdorf, 1964). However, in a subsequent experiment (Tables 2 and 3), a 'break-through' in ovulation was observed in some rats when 10 µg of ethynyloestradiol or AY-11483 was given daily during the pre-ovulatory period in combination with 100 µg of chlormadinone acetate or AY-11440. Similar effects with oral contraceptives were suggested in human trials (Rudel, Martinez-Manautou & Maquico-Topete, 1965). Since a single administration of 100 µg of either ethynyloestradiol or AY-11483 did not prevent ovulation (Table 2), but precluded implantation (Table 4), it is believed that the process of implantation in rats is more vulnerable to these oestrogens than is ovulation.

The primary contraceptive effect of excessive oestrogen or oestrogen plus progestin in mated rats before nidation was perhaps due to the expulsion of fertilized eggs (Banik & Pincus, 1964; Pincus, Banik & Jacques, 1964; Bennett, Kendle, Vallance & Vickery, 1966). Although a potent anti-oestrogenic progestin, on the other hand, delays the implantation of eggs (Nutting & Sollman, 1967), data presented in this paper (Table 4) are in agreement with our previous findings (Revesz *et al.*, 1967) that both chlormadinone acetate and

AY-11440 alone do not inhibit implantation or cause delayed implantation in rats. It may also be pertinent that oestrogen or oestrogen plus progestin may create an unfavourable uterine environment and preclude implantation. Although the perfect functional equilibrium between oestrogen and progesterone is required for ovum implantation (Psychoyos, 1966; Mayer, 1966), the expulsion of eggs by AY-11483 alone or with AY-11440 (Table 4) is thought to be the primary mode of action.

The present study indicates that AY-11483 is orally active in preventing pregnancy in rats by interfering with both ovulation and implantation.

ACKNOWLEDGMENTS

We are grateful to Dr R. I. Dorfman of Syntex Research Laboratories for a gift of chlormadinone acetate. Our sincere thanks are due to Mrs Irene Haltrecht and to Mrs Monique Politick for their efficient technical assistance.

REFERENCES

- BANIK, U. K. & PINCUS, G. (1962) The effect of steroidal anti-progestins on the implantation of fertilized eggs of rats and mice. *Proc. Soc. exp. Biol. Med.* **111**, 595.
- BANIK, U. K. & PINCUS, G. (1964) Estrogen and transport of ova in the rat. *Proc. Soc. exp. Biol. Med.* **116**, 1032.
- BENNETT, J. P., KENDLE, K. E., VALLANCE, D. K. & VICKERY, B. H. (1966) A comparison of the anti-fertility and hormonal activities of some new synthetic oestrogens. *Acta endocr., Copenh.* **53**, 443.
- CHAMBON, Y. & VEVE, Y. (1966) Effets anti-ovulatoires de l'éthynyl oestradiol et du mestranol, isolement ou en association avec le chlormadinone, chez le lapin et chez le rat. *C. r. Séanc. Soc. Biol.* **160**, 2411.
- DICZFALUSY, E. (1968) Mode of action of contraceptive drugs. *Am. J. Obstet. Gynec.* **100**, 136.
- DORFMAN, R. I. & DORFMAN, A. S. (1963) Pituitary gonadotropic inhibitory action of estrone in para-biotic rats. *Endocrinology*, **72**, 115.
- EDGREN, R., JONES, R. C. & PETERSON, D. L. (1967) A biological classification of progestational agents. *Fert. Steril.* **18**, 238.
- LEFEBVRE, Y., MARSHALL, D. J., REVESZ, C., BANIK, U. K. & DEGHENGI, R. (1967) Preparation and biological properties of 17-substituted oestradiol derivatives in the 8-normal and 8-iso series. *Am. Chem. Soc. Meet. Abstract No. 154*, p. 10.
- MAINLAND, D. & MURRAY, I. M. (1952) Tables for use in four-fold contingency tests. *Science, N.Y.* **116**, 591.
- MAYER, G. (1966) Techniques de nidation retardée chez la ratte et possibilités d'utilisation en tératogénèse expérimentale. *Bull. schweiz. Akad. med. Wiss.* **22**, 123.
- MCCANN, S. M. & TALEISNIK, S. (1961) The effect of estrogen on plasma luteinizing hormone (LH) activity in the rats. *Endocrinology*, **69**, 909.
- NUTTING, E. T. & SOLLMAN, P. B. (1967) Delay of implantation in intact rats treated with progestins. *Acta endocr., Copenh.* **54**, 8.
- PINCUS, G., BANIK, U. K. & JACQUES, J. (1964) Further studies on implantation inhibitors. *Steroids*, **4**, 657.
- PSYCHOYOS, A. (1966) Recent researches on egg implantation. *Ciba Fdn Study Grps*, **23**, 4.
- REVESZ, C., BANIK, U. K. & HERR, F. (1967) Hormonal and biological activities of 3 β ,17 α -diacetoxy-6-chloropregna-4,6-diene-20-one (AY-11,440). *Steroids*, **10**, 291.
- RUDEL, H. W., MARTINEZ-MANAUTOU, J. & MAQUIGO-TOPEPE, M. (1965) The role of progestogens in the hormonal control of fertility. *Fert. Steril.* **16**, 158.
- WATNICK, A. S., GIBSON, J., VINEGRA, M. & TOLKSDORF, S. (1964) Ethynyl estradiol: A potent orally active contraceptive in rats. *Proc. Soc. exp. Biol. Med.* **116**, 343.