Oocyte recovery, maturation and fertilization *in vitro* in the puma (*Felis concolor*)*

A. M. Miller†, M. E. Roelke‡, K. L. Goodrowe§, J. G. Howard† and D. E. Wildt†

†National Zoological Park, Smithsonian Institution, Washington DC 20008, USA; ‡Florida Game and Fresh Water Fish Commission, Gainesville, FL 32601, USA; and §Metro Toronto Zoo, Box 280, West Hill, Ontario M1E 4R5, Canada

Summary. Eight female pumas were treated i.m. with 1000 (N = 5) or 2000 (N = 3) i.u. PMSG followed 84 h later by 800 i.u. hCG. Eggs were recovered 24–26 h after hCG from ovarian follicles by using laparoscopy and transabdominal aspiration. Mature eggs were inseminated *in vitro* 4–6 h later whereas immature eggs were cultured for 24 h and then inseminated. Electroejaculates from 3 pumas were diluted with mKRB before insemination to evaluate the influence of sperm concentration on fertilization. Seven of 8 pumas responded with follicle development, and 140 eggs were recovered from 145 follicles (96.6%; 77 mature, 43 immature, 20 degenerate eggs; mean $\pm$ s.e.m., 20.0 ± 5.9 eggs/female). Overall fertilization rate was 43.5% (total eggs fertilized = 40) despite using inseminates containing 82–99% pleiomorphic spermatozoa. Of the 36 immature oocytes matured *in vitro* and inseminated, 12 were fertilized even though 50% of the inseminating spermatozoa contained an acrosomal defect. Fertilization rate of mature oocytes collected from follicles appeared unrelated ($P > 0.05$) to PMSG dose or number of spermatozoa/inseminate. This study demonstrates that a high proportion of follicular eggs can be recovered laparoscopically from adult pumas treated with PMSG and hCG. These gametes are capable of being fertilized *in vitro* (immediately or after maturation *in vitro*) even with low quality semen with a high incidence of sperm pleiomorphisms.

Keywords: *in vitro* fertilization; puma; oocyte maturation; teratospermia

Introduction

The puma (*Felis concolor*) has been taxonomically subdivided into 30 subspecies and is geographically distributed in the western hemisphere from British Columbia, Alberta and Manitoba in the north to Chile and Argentina in the south (Young & Goldman, 1946). Although existing in considerable numbers in the western United States (Guggisberg, 1975; Anderson, 1983), there are few free-living pumas east of the Mississippi River. One extremely rare subspecies (*F. c. coryi*) exists precariously in the Big Cypress Swamp and Everglades regions of southern Florida (Belden, 1986). Fewer than 50 of these pumas, called Florida panthers, free-range in what is one of North America's most rapidly growing human population areas. Florida panther electroejaculates contain a high proportion of pleiomorphic spermatozoa (overall mean, $\sim 94\%$; 8 ejaculates from 5 males; J. G. Howard, M. E. Roelke & D. E. Wildt, unpublished data), a general finding made previously for the cheetah (*Acinonyx jubatus*: Wildt *et al.*, 1983, 1987b), clouded leopard (*Neofelis nebulosa*: Wildt *et al.*, 1986a), 2 subspecies of leopard (*Panthera pardus*: Wildt *et al.*, 1988; Brown

*Reprint requests to: Dr D. E. Wildt.

et al., 1989), certain geographically-isolated lions (*Panthera leo*: Wildt et al., 1987a) and pumas from the western regions of North America (*F. concolor*: Wildt et al., 1988). However, Florida panthers appear unique in the unusually high number of spermatozoa afflicted with deranged acrosomes (mean, ~50%), a much greater rate than observed in western pumas (mean, 1.4%, Wildt et al., 1988).

The rarity of the Florida panther restricts conventional, 'controlled' reproductive biology studies. Current efforts focus on opportunistic access to animals and parallel studies of the western puma as a model subspecies. Some baseline information is available on the reproductive physiology of the western puma in captivity. Analysis of circulating oestradiol and progesterone patterns suggests that the oestrous cycle is 17–25 days in length and that ovulation does not occur spontaneously (Bonney et al., 1981). Western pumas respond to exogenous follicle-stimulating hormone (FSH-P: Phillips et al., 1982) or pregnant mares' serum gonadotrophin (PMSG: Bonney et al., 1981; Moore et al., 1981) by developing ovarian follicles. The first successful artificial insemination of a non-domestic felid was achieved in the puma (Moore et al., 1981). Three adult females were treated with 1250 i.u. PMSG (i.m.) followed 72 h later by 1000 i.u. hCG (i.m.). Laparoscopy revealed that 2–6 follicles ruptured/female at 30–40 h after hCG. Each puma was inseminated *in utero* at laparotomy using $20\text{--}40 \times 10^6$ motile, electroejaculated spermatozoa/ml; one pregnancy and the birth of a cub resulted.

The present study examined the feasibility of in-vitro fertilization in a mixed population of western pumas and Florida panthers. This approach may have artificial breeding potential for the Florida panther because domestic cat eggs, recovered by laparoscopic aspiration, are capable of being fertilized *in vitro* and result in live-born offspring after embryo transfer (Goodrowe et al., 1988b). Because pumas exhibit an extreme case of teratospermia, this study also permitted evaluation of the impact of structurally abnormal spermatozoa on sperm/egg interaction. Spermatozoa from teratospermic domestic cats appear to be compromised in ability to penetrate homologous oocytes *in vitro* (Howard et al., 1988, 1989). Our specific objectives were to (1) determine the effects of PMSG and hCG on ovarian follicle development and oocyte maturation in pumas maintained chronically in captivity or recently captured from the wild; and (2) evaluate the ability of laparoscopically-recovered follicular oocytes to become fertilized *in vitro* with electroejaculates containing high proportions of pleiomorphic spermatozoa.

Materials and Methods

Animals. The study was conducted in Gainesville, Florida (USA) in June 1988 using 8 adult female pumas, 6 of the Texas subspecies (*F. c. stanleyana*), one Florida panther (*F. c. coryi*) and one hybrid female of unknown origin (Table 1). Three of the western females (Nos 3, 5 and 7) were wild caught in southwest Texas 3–6 weeks before study onset; therefore, reproductive histories were unknown. The 3 remaining western pumas (Nos 1, 4, 6) and one hybrid female (No. 2) had been maintained in captivity in northern Florida for more than 2 years or since birth, respectively. All were presumed nulliparous except Puma 6 which was a known proven breeder. The Florida panther (No. 8) had been in captivity since April 1987 and had not reproduced or been observed in behavioural oestrus. Three adult males served as sperm donors: (1) a wild-caught puma (No. 1) from Texas with an unknown reproductive history and captured 4 weeks before study onset; (2) a captive born hybrid male and proven breeder (No. 2); and (3) a Florida panther (No. 3) maintained in captivity since 1984 with no history of reproducing. On 3 occasions from 1985 to 1987, Male 3 was electroejaculated under anaesthesia and produced the following average semen characteristics: semen volume, 1.3 ml; sperm concentration, 13×10^6 /ml ejaculate; sperm motility rating, 85%; proportion of pleiomorphic spermatozoa, 96.3%; proportion of spermatozoa with abnormal acrosomes, 48.3%.

Captive pumas in Florida were maintained individually or in pairs (females) in outdoor enclosures and provided with continuous access to water and either Nebraska Brand Feline, Canine Racing Diet (Lincoln, NE, USA) or Western Plateau Feline Diet (Amarillo, TX, USA) (1.4–2.4 kg/animal/day). Before transport from Texas to Florida, the newly captured pumas were maintained in individual outdoor enclosures and fed red meat (2.3–3.6 kg/day). These animals were placed in crates 12 h before scheduled gonadotrophin treatment and shipped by air transport (over a 6-h interval) to Gainesville.

Induction of ovarian activity, laparoscopy and oocyte recovery. Female pumas received i.m. injections of 1000 i.u. (N = 5) or 2000 i.u. (N = 3) PMSG (Equitech Inc., Atlanta, GA, USA). To ensure complete hormone delivery,

females were sedated with 40 mg diazepam i.m. (Valium®: Elkins-Sinn Inc., Cherryhill, NJ, USA) and 200–400 mg ketamine hydrochloride (HCl) i.m. (Vetalar®: Parke-Davis, Morris Plains, NJ, USA) before the PMSG dose. Precisely 84 h after PMSG, each puma was injected with 800 i.u. hCG i.m. (Sigma Chemical Co., St Louis, MO, USA) but without pre-treatment sedation. For the western pumas being transported from Texas to Florida, PMSG was administered immediately after translocation. The 84-h period between PMSG and hCG was based on findings suggesting that this interval maximized intrafollicular egg maturation in the domestic cat (Miller *et al.*, 1988).

All oocyte recovery attempts were made 24–26 h after hCG. A surgical plane of anaesthesia was induced with ketamine HCl i.m. (10 mg/kg body weight), and each female was intubated and maintained on inhalation anaesthesia of halothane and oxygen. Standard laparoscopy procedures were used (Wildt *et al.*, 1981; Phillips *et al.*, 1982). A pneumoperitoneum was established using room air, and the ovaries were observed using a 10-mm diameter, 180° laparoscope assembly placed mid-ventrally through a 20-mm skin incision. A Verres needle probe (2 mm in diameter) was used to manipulate the reproductive organs into view and aid counting ovarian follicles or corpora haemorrhagica or estimating follicle size. Laparoscopic oocyte aspirations were performed using the procedures described by Goodrowe *et al.* (1988b) for the domestic cat. For aspirations, a 4-cm long, 22-gauge needle attached to size 100 polyethylene tubing (i.d. 0.86 mm; Clay Adams, Parsippany, NJ, USA) was rinsed with 2–3 ml of a modified Krebs–Ringer bicarbonate solution (mKRB; Toyoda & Chang, 1974; Niwa *et al.*, 1985; NIH Media Unit, Bethesda, MD, USA) containing 4 mg bovine serum albumin/ml (Fraction V, Miles Laboratories Inc., Elkhart, IN, USA) and 40 units heparin/ml medium. A siliconized 7 ml collection tube (Terumo Medical Corp., Elkton, MD, USA) was attached to the free end of the polyethylene tubing, and the aspiration system was driven by a vacuum pump (Gast Manufacturing Corp., Benton Harbor, MI, USA). The Verres probe was used to stabilize the reproductive tract, and the aspiration needle was inserted through the abdominal wall ventro-medial to each ovary. Distinct follicles ≥ 2 mm in diameter were perforated with the needle while applying negative pressure (100 mmHg) with the vacuum pump. After aspirating follicles from one ovary, the collection tube and aspiration needle were replaced and the procedure repeated for the contralateral ovary. Collection tubes from each animal were emptied into separate plastic culture dishes which were examined by stereomicroscopy. All eggs were transferred immediately into fresh mKRB medium (without heparin) and placed in a 5% CO₂ in air, humidified environment at 38°C. Each cumulus cell–oocyte complex was evaluated for maturational status and classified as: (1) mature, corona radiata and cumulus oophorus cells loosened and expanded; (2) immature, tightly compacted corona radiata; or (3) degenerate, egg appeared abnormal, pale and/or lacked a corona radiata. Eggs were washed 3 times in mKRB under lightweight paraffin oil (Fisher Scientific Co., Fair Lawn, NJ, USA), placed in fresh medium and returned to the incubator.

Semen collection and processing. Spermatozoa were obtained by electroejaculation using a previously described, standardized procedure (Wildt *et al.*, 1983, 1987b, 1988; Howard *et al.*, 1986). In brief, this involved inducing a surgical plane of anaesthesia with ketamine HCl (10 mg/kg body weight; i.m.) or Telazol® (8 mg/kg, A. H. Robbins, Richmond, VA, USA). Each animal was given 80 electrical stimuli of similar voltage (2–8 V) and milliamperage (20–175 mA) administered over ~26 min. An AC, 60-Hz sine-wave electroejaculator (P-T Electronics, Boring, OR, USA) and rectal probe were used to deliver stimuli in 3 series of 30, 30 and 20 stimuli, respectively, with a 3–5 min rest period permitted between series.

Electroejaculate volume was recorded, and a 10 μ l sample was used to calculate sperm concentration using procedures described previously (Wildt *et al.*, 1983, 1988; Howard *et al.*, 1986). An undiluted 5- μ l sample of ejaculate was examined ($\times 250$) immediately after collection by 2 technicians to provide subjective estimates of sperm percentage motility and progressive status (0 to 5 scale; 0 = no forward progression or movement to 5 = rapid, linear forward progression; Wildt *et al.*, 1983, 1988; Howard *et al.*, 1986). An average motility rating was calculated for each male. Gross morphological assessments were made by fixing a 25 μ l sample of the semen in 1% glutaraldehyde and later evaluating 200 spermatozoa/ejaculate by phase-contrast microscopy ($\times 1000$).

In earlier studies of domestic cat fertilization *in vitro* (Goodrowe *et al.*, 1988b), electroejaculates were subjected to 'swim-up' processing (low-speed centrifugation and supernatant removal followed by sperm migration from the pellet into fresh culture medium). This permitted recovery of mostly motile, structurally normal spermatozoa. Preliminary studies indicated that swim-up processing was ineffective in the puma, probably because of fewer spermatozoa/ml of ejaculate, poorer sperm motility ratings and more abnormal sperm forms than observed in the domestic cat. Because puma electroejaculates were low in volume and overall quality, sufficient sperm numbers were available to permit only one semen processing treatment/male: (1) diluted in mKRB, insemination concentration was 2.0×10^6 motile spermatozoa/ml (Male 1); (2) semen centrifugation (300 g, 8 min), supernatant discarded and pellet diluted in mKRB to 1.0×10^6 motile spermatozoa/ml (Male 2); and (3) semen centrifugation (300 g, 8 min), supernatant discarded and pellet diluted in mKRB to 0.2×10^6 motile spermatozoa/ml (Male 3). During the processing to pre-insemination interval, spermatozoa were maintained at room temperature in mKRB, and eggs from at least 3 donors were inseminated with spermatozoa from each male.

Insemination and egg/sperm co-culture. Mature eggs were inseminated with a 100 μ l sample of the diluted sperm suspension from Males 1 and 2. Therefore, these eggs were placed with 0.2×10^6 and 0.1×10^6 total spermatozoa, respectively, under oil in a 35 $\times$ 10 mm Petri dish. Eggs from individual females were maintained in separate sperm cell drops (≤ 10 oocytes/drop) and the fertilization dishes were placed into a humidified, 5% CO₂ in air incubator at 38°C. The same incubator system was used to culture immature eggs in mKRB for 24 h to advance maturation. Eggs with an expanded corona radiata and cumulus cell complex were considered to have matured *in vitro*, were inseminated with 100 μ l of the sperm suspension from Male 3 (0.02×10^6 spermatozoa) and cultured under the same conditions as described for mature oocytes. Degenerate eggs were excluded from culture. To determine the rate of

parthenogenesis, control eggs (representative oocytes from each PMSG treatment and maturational status were used; total $n = 12$) were cultured under standard conditions in medium containing no spermatozoa. At 18–20 h after insemination, eggs were removed from the incubator and washed 3 times in a 0.2% hyaluronidase solution (Type 1-S, from bovine testes; Sigma Chemical Co.) for 3 min to remove residual cumulus cells and loosely attached spermatozoa. Eggs were returned to the incubator in 100 μ l drops of fresh, equilibrated mKRB under oil and examined 24 h after insemination.

Assessment of fertilization. Fertilization criteria were those described previously (Goodrowe *et al.*, 1988b) and were dependent on the presence of 2 polar bodies, 2 pronuclei or cleavage to at least the 2-cell stage. Eggs with more than 2 nuclear structures within the cytoplasm were considered polyspermic. Puma eggs, like oocytes from other felid species (Goodrowe *et al.*, 1988b, 1989), were extremely dark and opaque making it impossible to identify accurately the germinal vesicle or pronuclei using conventional light microscopy. Therefore, uncleaved eggs were treated with a DNA-specific fluorescent stain, Hoechst 33342 (H342; bisbenzamine; Sigma Chemical Co.), as described for a variety of mammals (Pursel *et al.*, 1985) including the domestic cat (Goodrowe *et al.*, 1988b). Eggs were counterstained with 0.1% Trypan blue (Sigma Chemical Co.) for 1–2 min, incubated in H342 (0.09 mg/ml, 15 min, 23°C) and examined using differential interference contrast (DIC) and fluorescence optics ($\times 250$ and $\times 400$) for germinal vesicles, pronuclei, polar bodies or polyspermic fertilization.

Statistics. Average values are reported as means $\pm$ the standard error of the mean (s.e.m.). A Student's *t* test was used to evaluate differences in the mean numbers of mature or immature eggs collected between PMSG dosages. The proportion of mature, immature, degenerate or fertilized eggs was compared between PMSG treatments by χ^2 analysis. Potential differences among males in the numbers of morphologically abnormal spermatozoa produced, fertilization rate *in vitro* and incidence of polyspermia also were evaluated by χ^2 statistics.

Results

The combined use of PMSG and hCG was effective in stimulating ovarian activity in 7 of 8 pumas (Table 1). Only Female 8 (the Florida panther) had no follicular or luteal activity at the laparoscopy after hCG; this animal died abruptly 2 months later from kidney disease and a hyperthyroidism condition. In the pumas responding to gonadotrophin stimulation, follicle (≥ 2 mm in diameter) numbers ranged from 8 to 52/female (Table 1). One animal (No. 1) apparently began ovulating before laparoscopic aspiration, as fresh corpora haemorrhagica were observed on both ovaries. The 3 wild-caught pumas tended to produce more follicles (30.6 ± 10.7) than the females chronically maintained in captivity (13.3 ± 2.6) but the difference was not significant ($P = 0.12$). A total of 140 eggs from 145 follicles (96.6% recovery) was collected from the 7 pumas (mean, 20.0 ± 5.9 eggs/female). Of these, 77 were classified as mature (55.0%), 43 as immature (30.7%) and 20 (14.3%) were characterized as degenerate. The immature eggs were collected from 4 pumas and 93% of these were recovered from the recently captured animals.

A total of 108 eggs (71 matured *in vivo*; 37 matured *in vitro*) was inseminated, and 16 of these were damaged, lost or could not be precisely classified after staining. Overall fertilization rate for all oocytes inseminated was 43.5% (total eggs fertilized = 40), averaging 50.0% for those matured *in vivo* and 33.3% for those matured *in vitro* (Table 1). A total of 10 eggs had cleaved to the 2-cell stage of development within 24 h of insemination. Parthenogenetic cleavage was not observed in any control egg.

Although limited numbers of animals were tested, the data were examined on the basis of PMSG dose. Pumas treated with 1000 i.u. PMSG and demonstrating ovarian activity ($N = 4$) tended to produce more ovarian follicles (mean, 24.8 ± 9.5) than those given 2000 i.u. (mean, 15.3 ± 2.9) but the difference was not significant ($P = 0.14$). Neither mean number of immature eggs recovered (1000 i.u., 6.5 ± 3.3 versus 2000 i.u., 5.7 ± 5.7), mean number of mature eggs recovered/female (1000 i.u., 14.3 ± 4.1 versus 2000 i.u., 6.7 ± 2.1) or overall fertilization rate *in vitro* was influenced by PMSG treatment ($P > 0.05$). Of the 39 mature eggs inseminated in the 1000 i.u. group, 6 cleaved and an additional 13 fertilized as determined by Hoechst staining (overall rate, 48.7%). Of the 17 mature eggs recovered from the 2000 i.u. PMSG group and co-cultured with spermatozoa, 4 cleaved and 5 exhibited intracellular evidence of fertilization (overall rate, 52.9%).

Semen characteristics for the 3 sperm donors are depicted in Table 2. Overall, 50% or fewer of all spermatozoa demonstrated forward progressive motility, and 82% or more of all cells exhibited

Table 1. Ovarian activity, egg recovery and fertilization for individual female pumas after PMSG/hCG treatment

Females	Animal status	Ovarian activity on day of aspiration*	Eggs				No. of fertilized eggs†	
			No. collected	No. mature	No. immature	No. degenerate†	Matured <i>in vivo</i>	Matured <i>in vitro</i>
1§	Captive, 2 years	10 follicles 3 corpora haemorrhagica	11	10	0	1	7/10 (70.0%)	—
2§	Captive born	16 follicles	7	7	0	0	1/5 (20.0%)	—
3§	Wild-caught, 4 weeks	20 follicles	23	3	17	3	1/2 (50.0%)	6/13 (46.2%)
4¶	Captive, 2 years	8 follicles 1 corpus albicans	6	5	0	1	3/5 (60.0%)	—
5¶	Wild-caught, 3 weeks	52 follicles	52	24	15	13	9/21 (42.8%)	1/12 (8.3%)
6¶	Captive, 2 years	19 follicles 5 corpora albicantia	22	17	3	2	5/7 (71.4%)	0/3 (0%)
7¶	Wild-caught, 6 weeks	20 follicles	19	11	8	0	2/6 (33.3%)	5/8 (62.5%)
8¶	Captive, 2 years	no activity	0	—	—	—	—	—

*Follicles ≥ 2 mm in diameter.

†Degenerate eggs were not inseminated.

‡Control eggs ($n = 12$) and damaged or lost eggs ($n = 16$) are excluded.

§Received 2000 i.u. PMSG.

¶Received 1000 i.u. PMSG.

Table 2. Ejaculate characteristics and in-vitro fertilization results for individual male pumas

	Male 1 (2.0×10^6)*	Male 2 (1.0×10^6)*	Male 3† (0.2×10^6)*
Ejaculate volume (ml)	0.37	1.52	1.42
Spermatozoa/ml ejaculate ($\times 10^{-6}$)	27.0	1.3	4.0
Sperm motility (%)	40	50	40
Sperm progressive status	2.5	2.5	3.0
Morphologically abnormal spermatozoa (%)	98	82	99
Abnormal head shape	0	0	12
Abnormal acrosome	5	6	50
Abnormal/missing midpiece	1	6	3
Bent midpiece with droplet	16	6	1
Bent midpiece without droplet	5	14	1
Proximal/distal droplet	4	20	14
Tightly coiled flagellum	66	20	18
Bent flagellum	1	10	0
No. of fertilized eggs/ total eggs (%)	11/21 (52.4)	16/33 (48.5)	13/38‡ (34.2)
No. of polyspermic eggs/ total eggs (%)	2/21 (9.5)	3/33 (9.1)	3/38 (7.9)

*Motile spermatozoa/ml; insemination concentration/ml.

†*Felis concolor coryi* (Florida panther) male.

‡Eggs were matured *in vitro* 24 h before insemination.

structural deformities. In Males 1 and 2, the majority of morphological defects consisted of mid-piece and flagellar abnormalities (93 and 76%, respectively) whereas 62% of the spermatozoa from Male 3 (the Florida panther) had head deformities, including 50% with deranged acrosomes. Although processing procedures and total number of spermatozoa inseminated varied among males, the overall fertilization rate of oocytes did not differ, being 52.4% for Male 1, 48.5% for Male 2 and 34.2% for Male 3 ($P > 0.05$). However, spermatozoa from Male 3 were used to inseminate in-vitro-matured eggs. Even under these conditions, more than one-third of the eggs exposed to the low quality spermatozoa of Male 3 demonstrated evidence of fertilization. The overall incidence of polyspermy was 8.8% and, among males, appeared unrelated to individual or numbers of spermatozoa inseminated (Table 2).

Discussion

As observed in the domestic cat (Goodrowe *et al.*, 1988b), a single injection of PMSG followed by hCG stimulated sufficient ovarian activity in the puma to permit collection of high quality, intra-follicular oocytes capable of being fertilized *in vitro*. The PMSG to hCG interval, laparoscopic aspiration technique and egg/sperm co-culture system developed for the domestic cat were effective for the puma, confirming previous assertions that the cat is a suitable model for developing embryo strategies for non-domestic felid species (Wildt *et al.*, 1986b). The data, however, re-emphasized the variability to be expected in ovarian response and egg/embryo quality following hormonal treatment, both among and within species. Striking differences in intrafollicular egg integrity and in-vitro fertilizability exist between the domestic cat and leopard cat (*F. bengalensis*), a small, non-domestic felid indigenous to southeast Asia (Goodrowe *et al.*, 1989). Although capable of interbreeding with domestic cats and producing hybrid offspring, leopard cats given the same PMSG/hCG regimen and subjected to the same laparoscopic recovery and culture procedures as

the domestic cat produce inferior quality eggs and overall lower fertilization rates *in vitro*. Furthermore, a higher proportion of leopard cat eggs were classified as degenerate (~45%) compared to domestic cat eggs (~14%; Goodrowe *et al.*, 1988b) and the puma eggs (14.3%) studied here. Although the basic mechanisms associated with follicular development, egg maturation and fertilizability probably are similar among felid species, subtle and perhaps critical species-specific differences may exist. This variation illustrates the evolutionary diversity of the felids and indicates that there are physiological differences among species which may present formidable challenges to the practical application of artificial breeding strategies.

The variation in PMSG-induced ovarian responses among domestic cats has been attributed, in part, to individual differences in animal sensitivity (Wildt *et al.*, 1978; Cline *et al.*, 1980). The greatest variability among pumas was in the numbers of ovarian follicles induced to develop and eggs recovered/female. Some of this variation may have been due to differences in reproductive status at the start of treatment. Pumas cycle throughout the year in North America (Anderson, 1983), and it is possible that some females were in a follicular phase which was enhanced further by the PMSG-hCG treatment. A single injection of 1000 i.u. PMSG resulted in a range of 8–52 follicles which was greater than the 2–6 CL observed by Bonney *et al.* (1981) after treating pumas with 1250 i.u. PMSG. Although this difference may be attributable to variations in PMSG source and potency, Bonney *et al.* (1981) administered hCG 72 h after PMSG compared to 84 h in our study. In domestic cats, extending the PMSG to hCG interval tends to increase the number of ovarian follicles produced and the total number of eggs recovered (Miller *et al.*, 1988).

The finding that one puma was ovulating 26 h after hCG suggests that any further delay in follicle aspiration after hCG would result in ovulation and loss of eggs before laparoscopic recovery. Even so, ~30% of all eggs were immature at aspiration, indicating that the asynchronous follicles were being recruited which resulted in eggs of varied maturational status. The rate of human intrafollicular egg maturation *in vivo* is directly related to the interval of follicle exposure to hCG (Botero-Ruiz *et al.*, 1981). Perhaps the more immature puma follicles were slow to respond to PMSG, experienced an inadequate exposure interval to hCG and, thus, produced immature eggs. The kinetics of in-vivo or in-vitro egg maturation have not been studied extensively in carnivores. However, we recently developed an in-vitro maturation system for follicular oocytes recovered from ovariectomized domestic cats (Johnston *et al.*, 1989). Of the immature antral oocytes cultured with gonadotrophins, 54% achieved metaphase II by 48 h. In the present study, based on subjective evaluation of cumulus complexes and corona radiata expansion, 100% of the oocytes appeared to have matured *in vitro* after 24 h in culture. Treating the females with gonadotrophins before collection and a high selectivity of oocytes before culture probably enabled all puma eggs to mature visibly *in vitro*. The in-vitro fertilization rate for these eggs (~33%) was very similar to that observed for domestic cat eggs matured *in vitro* (36%).

Wild-caught pumas responded well to the gonadotrophin treatment despite recent capture and long-distance transport. Even though receiving the PMSG treatment immediately after transit, these females produced high numbers of quality eggs which fertilized at a rate no different from that of pumas maintained in captivity for more than 2 years. Adrenal activity in captive pumas is very sensitive to an acute stress (Wildt *et al.*, 1988): males are capable of secreting massive concentrations of glucocorticoids as demonstrated by a 6-fold increase in circulating cortisol values within 1.5 h of combined anaesthesia and electroejaculation (Wildt *et al.*, 1988). Unlike in certain felid species (Brown *et al.*, 1988, 1989; Wildt *et al.*, 1988), elevated cortisol has no effect on testosterone production in pumas (Wildt *et al.*, 1988). A wild species like the puma is exposed continuously to stressors, and survival is highly dependent on adaptive abilities to minimize any adverse effects of hyperadrenal activity on reproduction. No doubt, the transported pumas in our study were experiencing stress, yet the females readily responded to exogenous gonadotrophins. Our collective observations of the puma suggest that gonadal functions in both the male (testosterone production) and female (PMSG-induced follicle development) appear relatively resilient to factors which potentially elicit stress.

Sperm capacitation is a prerequisite to fertilization, and capacitating spermatozoa of many species requires extensive processing *in vitro* (Brackett, 1981; Yanagimachi, 1981). We have demonstrated that domestic cat spermatozoa readily capacitate after low-speed centrifugation and 1-h 'swim-up' processing (Goodrowe *et al.*, 1988a). Puma spermatozoa also were easily capacitated, and eggs were fertilized by ejaculates which either were centrifuged only and the seminal plasma removed or simply used undiluted and unprocessed. This was fortunate because preliminary studies indicated that the overall poor ejaculate quality of pumas precluded using swim-up processing for recovering spermatozoa. Howard *et al.* (1989) reported that removing the seminal plasma from domestic cat ejaculates is not necessary for cat spermatozoa to bind and penetrate homologous eggs. Apparently the decapacitation factor which plays such an important role in controlling fertilization in some species (Fraser, 1984; Thomas *et al.*, 1986; Shivaji & Bhagava, 1987) either does not exist or has little consequence in either the domestic cat or puma.

All male pumas produced high numbers of pleiomorphic spermatozoa/ejaculate, a finding which was not surprising in view of similar observations in a variety of other felids of large size (Wildt *et al.*, 1983, 1987a, 1988). The aetiology of this characteristic is unknown, although several studies have related the number of abnormal spermatozoa in felid ejaculates to a lack of genetic variability (O'Brien *et al.*, 1986; Wildt *et al.*, 1983, 1987a, b). Florida panthers produce fewer polymorphic loci than do any other puma subspecies studied to-date (Roelke, 1988; M. E. Roelke & S. J. O'Brien, unpublished data). Additionally, 8 of 17 (47%) free-ranging males examined during field capture episodes are unilaterally cryptorchid (M. E. Roelke, unpublished observations), a condition which is highly heritable in other species (Farrer *et al.*, 1985; Knap, 1986; MacKellar, 1987). These findings suggest that the Florida panther has a narrow genetic base and that inbreeding may be contributing to certain reproductive characteristics including the high incidence of abnormal acrosomes.

It is important to define more clearly the impact of teratospermia on the ability of a spermatozoon to participate in fertilization and normal embryo development. A high incidence of teratospermia reduces fertility in non-felid species (Hulet *et al.*, 1965; Singleton & Shelby, 1972; Munro, 1981; Saacke *et al.*, 1988; Bader *et al.*, 1988) and decreases in-vitro fertilization rates in humans (Bostofte *et al.*, 1982; Jeulin *et al.*, 1986). Spermatozoa from teratospermic, domestic cat ejaculates are less capable of binding and penetrating zona pellucida-free hamster eggs and zona-intact domestic cat eggs *in vitro* than are those from normospermic ejaculates (Howard *et al.*, 1988, 1989). The in-vitro fertilization rate of puma eggs using western puma (50%) and Florida panther (34%) spermatozoa was considerably less than the ~75% rate for domestic cat eggs using normospermic ejaculates and the same culture system. None the less, it was rather remarkable that 1 in 3 in-vitro matured puma eggs co-cultured with Florida panther spermatozoa demonstrated evidence of fertilization, especially considering that 50% of all spermatozoa contained a severely deformed acrosome. Although these ejaculates contain extreme numbers of structurally abnormal spermatozoa, it is apparent that fertility is not completely compromised. This would explain why certain males have sired litters in the wild, even though opportunistic semen evaluations have indicated a comparable level of teratospermia (M. E. Roelke, J. G. Howard & D. E. Wildt, unpublished observations). The actual reproductive performance of the Florida panther and the level at which teratospermia begins to contribute to suboptimal fertility remain to be determined. Field observations indicate that not all adult Florida panthers reproduce (M. E. Roelke, unpublished observations). Male 3, in captivity for more than 4 years, has failed to breed naturally even though exposed to several oestrous females. It will be difficult to determine the impact of teratospermia on fertility of the current Florida panther population. However, it is encouraging that ejaculates which would be considered extremely poor quality by conventional assessment standards could be used to produce embryos. Although the biological competence of these embryos *in vivo* remains to be determined, the present results suggest that in-vitro fertilization may be a viable alternative for ensuring that such individuals contribute to a subspecies recovery plan.

We thank Dr Donald Forrester, Dr George Kollias, Dr Elliot Jacobson, Dr Collin Burrows, Dr Martin Drost and Dr Victor Shille and the University of Florida, College of Veterinary Medicine, Gainesville, for providing generous access to equipment, laboratories and surgical facilities; Tom Logan, Bureau Chief of Research, Steve Parker and Robert Belden of the Florida Game and Fresh Water Fish Commission; and John Lucas and the staff at White Oak Plantation and the Tallahassee Museum for co-operation and assistance. Portions of this project were supported by grants from the National Institutes of Health (HD 23853) and Friends of the National Zoo (to D.E.W.), the U.S. Fish and Wildlife Service through Endangered Species Grant funding (to M.E.R.) and the Blackstock Foundation (to K.L.G.). A.M.M. is supported, in part, by the Women's Committee of the Smithsonian Associates.

References

- Anderson, A.E. (1983) A critical review of literature on the puma (*Felis concolor*). *Spec. Rep. No. 54, Colo. Div. Wdlf.*
- Bader, H., Hoppen, H.O., Wockener, A. & Merkt, H. (1988) Case studies in stallions with fertility problems: endocrine and spermatological aspects. *Proc. 11th Int. Cong. Anim. Reprod. A.I., Dublin 3*, 367, abstr.
- Belden, R.C. (1986) Florida panther recovery plan implementation: a 1983 progress report. In *Cats of the World: Biology, Conservation and Management Proceedings of the 2nd International Cat Symposium*, pp. 159–172. Eds S. D. Miller & D. D. Everitt. Kingsville.
- Bonney, R.C., Moore, H.D.M. & Jones, D.M. (1981) Plasma concentrations of oestradiol-17 β and progesterone, and laparoscopic observations of the ovary in the puma (*Felis concolor*) during oestrus, pseudopregnancy and pregnancy. *J. Reprod. Fert.* **63**, 523–531.
- Bostofte, E., Serup, J. & Rebbe, H. (1982) Relation between morphologically abnormal spermatozoa and pregnancies obtained during a twenty-year follow-up period. *Int. J. Androl.* **5**, 379–386.
- Botero-Ruiz, W., Laufer, N., Decherney, A.H., Polan, M.L., Haseltine, F.P. & Behrman, H.R. (1981) The relationship between follicular fluid steroid concentration and successful fertilization of human oocytes *in vitro*. *Fert. Steril.* **41**, 820–826.
- Brackett, B.G. (1981) Applications of *in vitro* fertilization. In *New Technologies in Animal Breeding*, pp. 141–161. Eds G. E. Seidel, Jr, S. Seidel & B. G. Brackett. Academic Press, New York.
- Brown, J.L., Goodrowe, K.L., Simmons, L.G., Armstrong, D.L. & Wildt, D.E. (1988) Evaluation of the pituitary-gonadal response to GnRH, and adrenal status, in the leopard (*Panthera pardus japonensis*) and tiger (*Panthera tigris*). *J. Reprod. Fert.* **82**, 227–236.
- Brown, J.L., Wildt, D.E., Phillips, L.G., Seidensticker, J., Fernando, S.B.U., Miththapala, S. & Goodrowe, K.L. (1989) Ejaculate characteristics, and adrenal pituitary-gonadal interrelationships in captive leopards (*Panthera pardus kotiya*) isolated on the island of Sri Lanka. *J. Reprod. Fert.* **85**, 605–614.
- Cline, E.M., Jennings, L.L. & Sojka, N.J. (1980) Breeding laboratory cats during artificially induced estrus. *Lab. Anim. Sci.* **30**, 1003–1005.
- Farrer, J.H., Sikka, S.C., Xie, H.W., Constantinide, D. & Rajfer, J. (1985) Impaired testosterone biosynthesis in cryptorchidism. *Fert. Steril.* **44**, 125–132.
- Fraser, L.R. (1984) Mouse sperm capacitation *in vitro* involves loss of a surface-associated inhibitory component. *J. Reprod. Fert.* **72**, 373–384.
- Goodrowe, K.L., Miller, A.M. & Wildt, D.E. (1988a) Capacitation of domestic cat spermatozoa as determined by homologous zona pellucida penetration. *Proc. 11th Int. Cong. Anim. Reprod. A.I., Dublin 3*, 245, abstr.
- Goodrowe, K.L., Wall, R.J., O'Brien, S.J., Schmidt, P.M. & Wildt, D.E. (1988b) Developmental competence of domestic cat follicular oocytes after fertilization *in vitro*. *Biol. Reprod.* **39**, 355–372.
- Goodrowe, K.L., Miller, A.M. & Wildt, D.E. (1989) *In vitro* fertilization of gonadotropin-stimulated leopard cat (*Felis bengalensis*) follicular oocytes. *J. exp. Zool.* (in press).
- Guggisberg, C.A.W. (1975) *Wild Cats of the World*, pp. 328. Taplinger Publishing Co., New York.
- Howard, J.G., Bush, M. & Wildt, D.E. (1986) Semen collection, analysis and cryopreservation in non-domestic mammals. In *Current Therapy in Theriogenology*, pp. 1047–1053. Ed. D. A. Morrow. W. B. Saunders Co., Philadelphia.
- Howard, J.G., Barone, M.A., Clingerman, K.J., Bush, M. & Wildt, D.E. (1988) Influence of teratospermia, culture media and a sperm swim-up technique on penetration of zona-free hamster ova by domestic cat spermatozoa. *Proc. 11th Int. Cong. Anim. Reprod. A.I., Dublin 3*, 332, abstr.
- Howard, J.G., Hurlbut, S.L. & Bush, M. (1989) Compromised zona binding and penetration of homologous oocytes by spermatozoa from teratospermic domestic cats. *Biol. Reprod.* **40** (Suppl. 1), 161, abstr.
- Hulet, C.V., Foote, W.C. & Blackwell, R.L. (1965) Relationship of semen quality and fertility in the ram to fecundity in the ewe. *J. Reprod. Fert.* **9**, 311–341.
- Jeulin, C., Feneux, D., Serres, C., Jouannet, P., Guillet-Rosso, R., Belaisch-Allart, J., Fryman, R. & Testart, J. (1986) Sperm factors related to failure in human *in-vitro* fertilization. *J. Reprod. Fert.* **76**, 735–744.
- Johnston, L.A., O'Brien, S.J. & Wildt, D.E. (1989) *In vitro* maturation and fertilization of domestic cat follicular oocytes. *Gamete Res.* **24**, 343–356.
- Knap, P.W. (1986) Congenital defects inheritance of AI boars: genetic parameters and breeding value estimation procedures. *Livestock Prod. Sci.* **15**, 337–352.

- MacKellar, A.** (1987) Cryptorchidism. In *Andrology*, pp. 53–68. Eds J. P. Pryor & L. I. Lipshultz. Butterworths & Co., London.
- Miller, A.M., Goodrowe, K.L. & Wildt, D.E.** (1988) Influence of gonadotropin source and treatment interval on *in vitro* fertilization of domestic cat oocytes. *Biol. Reprod.* **38** (Suppl. 1), 73, abstr.
- Moore, H.D.M., Bonney, R.C. & Jones, D.M.** (1981) Induction of oestrus and successful artificial insemination in the cougar, *Felis concolor*. *Vet. Rec.* **108**, 282–283.
- Munro, I.B.** (1981) Bovine semen characteristics and fertility. *J. Reprod. Fert.* **2**, 513–571.
- Niwa, K., Ohara, K., Hosoi, Y. & Iritani, A.** (1985) Early events of *in-vitro* fertilization of cat eggs by epididymal spermatozoa. *J. Reprod. Fert.* **74**, 657–660.
- O'Brien, S.J., Wildt, D.E. & Bush, M.** (1986) The cheetah in genetic peril. *Sci. Amer.* **5**, 84–92.
- Phillips, L.G., Simmons, L.G., Bush, M., Howard, J.G. & Wildt, D.E.** (1982) Gonadotropin regimen for inducing ovarian activity in captive wild felids. *J. Am. vet. med. Assoc.* **181**, 1246–1250.
- Pursel, V.G., Wall, R.J., Rexroad, C.E., Jr, Hammer, R.E. & Brinster, R.L.** (1985) A rapid whole-mount staining procedure for nuclei of mammalian embryos. *Theriogenology* **24**, 687–691.
- Roelke, M.E.** (1988) Florida Panther Recovery Florida Panther Biomedical Investigation, E-1-12, pp. 31–33. Fl. Game Fresh Wat. Fish Comm.
- Saacke, R.G., Bame, J.H., Karabinus, D.S., Mullins, K.J. & Whitman, S.** (1988) Transport of abnormal spermatozoa in the artificially inseminated cow based upon accessory spermatozoa in the zona pellucida. *Proc. 11th Int. Cong. Reprod. A.I., Dublin* **3**, 292, abstr.
- Shivaji, S. & Bhagava, P.M.** (1987) Antifertility factors of mammalian seminal fluid. *Bioessays* **7**, 13–17.
- Singleton, W.L. & Shelby, D.R.** (1972) Variation among boars in semen characteristics and fertility. *J. Anim. Sci.* **34**, 762–766.
- Thomas, T.S., Wilson, W.L. & Oliphant, G.** (1986) Chemical and physical characterization of rabbit sperm acrosome stabilizing factor. *Biol. Reprod.* **35**, 691–703.
- Toyoda, Y. & Chang, M.C.** (1974) Fertilization of rat eggs *in vitro* by epididymal spermatozoa and the development of eggs following transfer. *J. Reprod. Fert.* **36**, 9–22.
- Wildt, D.E., Kinney, G.M. & Seager, S.W.J.** (1978) Gonadotropin-induced reproductive cyclicity in the domestic cat. *Lab. Anim.* **38**, 301–307.
- Wildt, D.E., Platz, C.C., Seager, S.W.J. & Bush, M.** (1981) Induction of ovarian activity in the cheetah (*Acinonyx jubatus*). *Biol. Reprod.* **24**, 217–222.
- Wildt, D.E., Bush, M., Howard, J.G., O'Brien, S.J., Meltzer, D., van Dyk, A., Ebedes, H. & Brand, D.J.** (1983) Unique seminal quality in the south African cheetah and a comparative evaluation in the domestic cat. *Biol. Reprod.* **29**, 949–947.
- Wildt, D.E., Howard, J.G., Hall, L.L. & Bush, M.** (1986a) The reproductive physiology of the clouded leopard: I. Electroejaculates contain high proportions of pleiomorphic spermatozoa throughout the year. *Biol. Reprod.* **34**, 937–947.
- Wildt, D.E., Schiewe, M.C., Schmidt, P.M., Goodrowe, K.L., Howard, J.G., Phillips, L.G., O'Brien, S.J. & Bush, M.** (1986b) Developing animal model systems for embryo technologies in rare and endangered wildlife. *Theriogenology* **25**, 33–51.
- Wildt, D.E., Bush, M., Goodrowe, K.L., Packer, C., Pusey, A.E., Brown, J.L., Joslin, P. & O'Brien, S.J.** (1987a) Reproductive and genetic consequences of founding isolated lion populations. *Nature, Lond.* **329**, 328–331.
- Wildt, D.E., O'Brien, S.J., Howard, J.G., Caro, T.M., Roelke, M.E., Brown J.L. & Bush, M.** (1987b) Similarity in ejaculate-endocrine characteristics in captive versus free-ranging cheetahs of two subspecies. *Biol. Reprod.* **36**, 351–360.
- Wildt, D.E., Phillips, L.G., Simmons, L.G., Chakraborty, P.K., Brown, J.L., Howard, J.G., Teare, A. & Bush, M.** (1988) A comparative analysis of ejaculate and hormonal characteristics of the captive male cheetah, tiger, leopard and puma. *Biol. Reprod.* **38**, 245–255.
- Yanagimachi, R.** (1981) Mechanisms of fertilization in mammals. In *Fertilization and Embryonic Development In Vitro*, pp. 181–183. Eds L. Mastroianni, Jr & J. D. Biggers. Plenum Press, New York.
- Young, S.P. & Goldman, E.A.** (1946) *The Puma: Mysterious American Cat*. The American Wildlife Institute, Washington.

Received 3 May 1989