Possible roles for prostaglandins E_2 and $F_{2\alpha}$ in seasonal changes in ovarian steroidogenesis in the frog (*Rana esculenta*)

A. Gobbetti and M. Zerani

Department of Molecular, Cellular, and Animal Biology, University of Camerino, via F. Camerini 2,
62032 Camerino MC, Italy

Concentrations of PGE_2 , $PGF_{2\alpha}$, androgens and oestradiol in plasma, and ovary weights were measured in the female frog, *Rana esculenta*, during the annual breeding cycle. Experiments were carried out *in vivo* to study the effects of PGE_2 and $PGF_{2\alpha}$ on plasma sex steroids during the following stages: pre-reproduction (April), reproduction (May), post-reproduction (June) and recovery (October). Experiments were performed *in vitro* during these stages to evaluate the effects of these two prostaglandins on the secretion of ovarian steroids. Concentrations of PGE_2 were low in plasma during winter hibernation, the reproduction and post-reproduction stages, whereas they were high during the pre-reproduction and recovery stages. PGE_2 treatment *in vivo* increased androgen secretion in April, whereas $PGF_{2\alpha}$ treatment increased oestradiol secretion in June and October. In experiments *in vitro*, PGE_2 increased androgen secretion and decreased oestradiol secretion from ovaries collected in April, whereas $PGF_{2\alpha}$ increased oestradiol secretion from ovaries collected in October. These results suggest that a seasonal increase in plasma PGE_2 may inhibit breeding activity, probably by stimulating ovarian androgen secretion, whereas, as previously reported, a seasonal increase in plasma $PGF_{2\alpha}$ may inhibit breeding, by stimulating ovarian oestradiol secretion.

Introduction

In mammals, ovarian steroidogenesis (Espey *et al.*, 1986; Sedrani and El-Banna, 1987; Dharmarajan *et al.*, 1989; Greenhalgh, 1990) is regulated by PGs of both F and E series. Prostaglandins are also implicated in the control of ovulation in the chicken (Shimada *et al.*, 1984) and of reproductive function in reptiles (Guillette *et al.*, 1984, 1990, 1991; Jones *et al.*, 1990).

Little is known, however, about the role of PGs in the reproductive processes of amphibians. In the anuran frog, *Rana esculenta* (Gobbetti *et al.*, 1990, 1991a; Gobbetti and Zerani, 1991), and in the urodele crested newt, *Triturus cristatus* (Gobbetti *et al.*, 1991b, c), seasonal changes in concentrations of $PGF_{2\alpha}$ in plasma have been observed to be related to the stages of the annual breeding cycle. The possibility that these seasonal changes in concentrations of the PGs may be related to reproductive function has not been investigated. A role for PGE_2 is suggested by a study *in vitro* on the male *Triturus cristatus* abdominal gland, where it was found to induce pheromonal activity (Gobbetti and Zerani, 1992a).

The objective of this study was to establish whether in the female frog *Rana esculenta* there are seasonal changes in the concentration of PGE_2 in plasma and how this relates to seasonal changes in $PGF_{2\alpha}$. A further objective was to determine *in vivo* and *in vitro* the effects of PGE_2 and $PGF_{2\alpha}$ on the secretion of ovarian steroids at different stages of the annual breeding cycle.

Materials and Methods

Animals

Adult female frogs were collected in Umbria, Italy, from a pond 870 m above sea level. The frogs breed in May (reproductive stage), when the temperature increases and enter a post-reproductive phase in the summer. Gonad recrudescence is initiated in mid-summer, and continues into the autumn (recovery stage). The animals hibernate during the cold months of winter to emerge when the temperature increases in the next spring. At the beginning of spring, the frogs return to the pond (pre-reproductive stage).

Observations on free-living and captive animals

Fifteen adult female frogs, *Rana esculenta*, were collected every month during 1991. Immediately after capture, they were anaesthetized with 3-aminobenzoic acid ethyl ester (Tricaine: Sigma Chemical Co., St Louis, MO), weighed, and a blood sample was taken through a heparinized microtube inserted into the heart. Individual blood samples (600 μ l) were collected into chilled tubes containing acetylsalicylic acid (Aspirin; Sigma) and EDTA (5 μ g and 7 μ g ml^{-1} of blood, respectively) (Gobbetti *et al.*, 1990). After centrifugation (1500 g for 10 min), plasma samples were stored at $-20^\circ C$. Ovaries were removed and weighed. In April, May, June and October, frogs were captured and blood samples were taken in the field (control 1), while a further 15 animals were transferred to laboratory aquaria, kept under natural lighting and temperatures and fed *ad libitum* to

study the effects of captivity on concentrations of hormones in plasma. A week later, blood samples were taken from the animals in captivity, and at the same time blood samples were taken from 15 animals in the field (control 2).

Experiments in vivo

In April, May, June and October, 84 female frogs were captured each month, transferred to laboratory aquaria and maintained under natural lighting and temperature. The frogs collected in May had not ovulated. One week later, the animals were assigned to four groups: (a) 21 frogs received a single s.c. injection of 20 ng PGE₂ (Sigma) g⁻¹ body weight dissolved in 100 µl amphibian saline (0.64% w/v NaCl solution); (b) 21 frogs received a single s.c. injection of 15 ng PGF_{2α} (Sigma) g⁻¹ body weight dissolved in 100 µl amphibian saline; (c) 21 frogs received a single s.c. injection of 100 µl amphibian saline only; (d) 21 untreated control animals. Blood samples were taken from seven of the 21 frogs in each group at 6, 18 and 36 h after treatment. After centrifugation (1500 g for 10 min), plasma samples were stored at -20°C. Blood samples were taken from a further batch of seven untreated animals at the beginning of the experiments (controls, time 0). The times of treatment and the minimum effective doses of PGE₂ and PGF_{2α} were chosen after preliminary experiments (data not shown).

Experiments in vitro

In April, May, June and October, seven female frogs were captured each month and transferred to laboratory aquaria. The frogs collected in May had not ovulated. One week later, the animals were decapitated, the ovaries rapidly removed and placed in cold Dulbecco's modified Eagle medium (DME; Sigma) containing 10 mmol Hepes l⁻¹, 1 mg Penicillin G ml⁻¹, and 2 mg streptomycin ml⁻¹. Each month, the ovary from one animal was divided into equal sized fragments, pooled and equally distributed over 12 incubation wells. Multiwell tissue culture plates (Becton Dickinson and Co., Lincoln Park, NJ) were used. Each set of wells was divided into three experimental groups, each consisting of four wells. The experimental groups were: (a) ovarian tissue incubated with DME alone; (b) ovarian tissue incubated with DME plus PGE₂ (75 ng); (c) ovarian tissue incubated with DME plus PGF_{2α} (75 ng). The final volume of each well was 1 ml. Culture plates were wrapped in aluminium foil and incubated in a shaking water bath (19°C), at 30 revolutions min⁻¹. Incubation medium was removed from separate wells after 25, 50, 100 and 200 min of incubation. The incubation medium samples were stored at -20°C until hormone assays were performed; ovarian tissues were homogenized in amphibian saline, and protein contents were determined using a commercial kit (Bio-Rad, Richmond, CA). The experiment was repeated with incubation media without ovarian tissue. The experiment was repeated seven times for each month. The incubation conditions and the minimum effective doses of PGE₂ and PGF_{2α} were determined in preliminary experiments (data not shown).

Determination of PGE₂, PGF_{2α}, androgens and oestradiol

Concentrations of PGE₂ and PGF_{2α} were measured by radioimmunoassay in plasma and incubation media as described

by Gobbetti *et al.* (1990) and Gobbetti and Zerani (1992a), respectively. Determinations were carried out on duplicate plasma samples (100 µl) and in incubation media (500 µl) extracted with 10 volumes of diethyl ether for 5 min. After centrifugation (1000 g for 5 min), organic fractions were transferred into glass tubes and evaporated to dryness under nitrogen. Extracts were resuspended in 100 µl assay buffer before assay. The recoveries of added labelled PGE₂ and PGF_{2α} were 86.9 ± 0.81% and 85.3 ± 0.71%, respectively (means of 10 determinations ± SD). Standard curves, for the two prostaglandins in buffer and extracted incubation medium were parallel.

Concentrations of androgens, oestradiol in plasma and incubation media were measured by radioimmunoassay as described by d'Istria *et al.* (1974).

Intra- and interassay coefficients of variation and minimum detectable doses were PGE₂, 6.5%, 12%, 18 pg; PGF_{2α}, 5%, 11.5%, 17 pg; androgens, 6%, 9%, 9.0 pg; oestradiol, 4%, 9%, 8.5 pg, respectively. The PGF_{2α} testosterone and oestradiol antisera were provided by G. F. Bolelli and F. Franceschetti (CNR-Physiopathology of Reproduction Service, University of Bologna) and the PGE₂ antiserum was purchased from Cayman Chemical (Ann Arbor, MI). Testosterone was not separated from 5α-dihydrotestosterone and, therefore, as the antiserum used is not specific, the data are expressed as total androgens. Tritiated PGE₂, PGF_{2α}, testosterone and oestradiol were purchased from Amersham International (Buckinghamshire), non-radioactive PGE₂, PGF_{2α}, testosterone and oestradiol from Sigma.

Statistical analysis

An analysis of variance (ANOVA) followed by Duncan's multiple range test (Duncan, 1955; Sokal and Rohlf, 1981) was used to analyse the data. Correlation coefficients were calculated as described by Scossiroli and Palenzona (1979). Data on ovary weights are shown as the gonadosomatic index (GSI: gonad weight:body weight).

Results

Seasonal changes in concentrations of plasma prostaglandins, gonadal steroids and GSI

The concentration of plasma PGE₂ was low in November, December and January, increased in February ($P < 0.01$) and remained high until April; it then fell in May ($P < 0.01$), increasing again in July until October ($P < 0.01$) (Fig. 1). The concentration of plasma PGF_{2α} was high in January and February ($P < 0.01$), decreased from March to May ($P < 0.01$), then increased from June to October ($P < 0.01$), and decreased in November and December ($P < 0.01$) (Fig. 1). Androgen concentrations were highest from January to April ($P < 0.01$), decreased in May ($P < 0.01$), remained low until August and increased from September to December (Fig. 1). Oestradiol concentrations in plasma showed a peak in June ($P < 0.01$) (Fig. 1). GSI decreased between May and June ($P < 0.01$), and then increased until October, and decreased between November and January. GSI began to increase again in February, reaching its

Fig. 1. Concentrations of plasma (a) PGE_2 , (b) $PGF_{2\alpha}$, (c) androgens and (d) oestradiol, and (e) gonadosomatic index (GSI: ovary weight:body weight) during the annual breeding cycle in female frog, *Rana esculenta*. Each mean refers to 15 determinations $\pm$ SD.

maximum value in spring ($P < 0.01$) (Fig. 1). Values of plasma PGE_2 were positively correlated with those of GSI ($r = 0.253$, $df = 178$, $P < 0.01$).

Effects of captivity on plasma prostaglandins and gonadal steroids

After one week in captivity, concentrations of plasma PGE_2 decreased in April and October ($P < 0.01$), concentrations of plasma $PGF_{2\alpha}$ decreased in June and October ($P < 0.01$), concentrations of plasma androgens decreased in April and October ($P < 0.01$), and concentrations of plasma oestradiol decreased in June ($P < 0.01$) (Fig. 2).

Experiments in vivo

Treatment with PGE_2 increased androgen concentrations in April, 6, 18 and 36 h after injection ($P < 0.01$) (Fig. 3). Treatment with $PGF_{2\alpha}$ increased oestradiol concentrations in June and October, 18 and 36 h after injection ($P < 0.01$) (Fig. 4).

Experiments in vitro

The basal release of PGE_2 from the ovary was higher in April and October ($P < 0.01$) than in May and June. Values in October were higher ($P < 0.05$) than those of April (Fig. 5). The basal release of $PGF_{2\alpha}$ in June was lower ($P < 0.01$) than in

Fig. 2. Effects of one week in captivity on the concentrations of plasma (a) PGE_2 , (b) $PGF_{2\alpha}$, (c) androgens and (d) oestradiol in female frog, *Rana esculenta*, during April (pre-reproduction), May (reproduction), June (post-reproduction), and October (recovery). Experimental groups: (▨) frogs captured in the field and blood samples taken at once (control 1); (■) blood samples taken from frogs after one week in captivity in laboratory; (□) frogs captured and blood samples taken at once in the field, one week after control 1 (control 2). Each mean refers to 15 determinations $\pm$ SD. * $P < 0.01$ versus control 1 and 2 (Duncan's multiple range test).

Fig. 3. Effects *in vivo* of 20 ng PGE_2 g^{-1} body weight on concentrations of plasma androgens in female frog, *Rana esculenta*, during April (pre-reproduction). Experimental groups: (□) untreated frogs; (▨) amphibian-saline-only injected frogs; (■) PGE_2 injected frogs. Each mean refers to seven determinations $\pm$ SD. * $P < 0.01$ versus untreated and amphibian-saline-only injected frogs (Duncan's multiple range test).

other months (Fig. 5). The basal release of androgens was higher in April ($P < 0.01$) than in other months (Fig. 6). Values in October were higher ($P < 0.01$) than in May or June (Fig. 6).

Fig. 4. Effects *in vivo* of 15 ng PGE₂ g⁻¹ body weight on concentrations of plasma oestradiol in female frog, *Rana esculenta*, during (a) June (post-reproduction) and (b) October (recovery). Experimental groups: (□) untreated frogs; (▨) amphibian-saline-only injected frogs; (■) PGE₂ injected frogs. Each mean refers to seven determinations ± SD. **P* < 0.01 versus untreated and amphibian-saline-only injected frogs (Duncan's multiple range test).

The basal concentration of oestradiol was lower in June (*P* < 0.01), than in other months (Fig. 7).

Treatment with PGE₂ increased the secretion of androgens in April after 50, 100 and 200 min (*P* < 0.01) (Fig. 6), and decreased oestradiol concentrations in April after 50, 100 and 200 min (*P* < 0.01) (Fig. 7). Treatment with PGE₂ increased oestradiol concentrations in October after 100 and 200 min (*P* < 0.01) (Fig. 7).

Hormones were not detected in media incubated without ovarian tissue (data not shown).

Discussion

This work reports, for the first time, the presence of plasma PGE₂ in the female frog, *Rana esculenta*, and showed that the concentrations of this prostaglandin are related to the annual reproductive cycle. Concentrations of PGE₂ were high in plasma during the pre-reproductive and recovery stages and low during the breeding season and winter hibernation. The seasonal changes in plasma PGE₂, androgens, oestradiol and ovarian weight confirm the findings of Gobetti *et al.* (1990) and Zerani *et al.* (1991). However, Gobetti *et al.* (1990) did not observe a peak in plasma oestradiol in June; this may be because this hormone peak is transient. Concentrations of plasma PGE₂ were high during the recovery stage of the breeding cycle; concentrations of androgens were high during the pre-reproduction stage, whereas concentration of oestradiol peaked during the post-reproductive stage.

The data *in vitro* on the basal release of PGE₂ and androgens from the ovary correlated with seasonal changes in plasma concentrations of these hormones. The highest amounts of PGE₂

Fig. 5. Basal release *in vitro* of PGE₂ (■) and PGF_{2α} (□) from ovary of female frog, *Rana esculenta*, incubated (a) in April (pre-reproduction), (b) May (reproduction), (c) June (post-reproduction) and (d) October (recovery). Each mean refers to seven determinations ± SD. a, *P* < 0.01 versus same time in May and June; b, *P* < 0.05 versus same time April; c, *P* < 0.01 versus same time April, May and June (Duncan's multiple range test).

and androgens released in ovarian incubations were observed in April and October. In contrast, the lowest release of PGE₂ and oestradiol from ovarian tissues was found in June, despite the fact that concentrations of plasma PGE₂ and oestradiol are high during this month. In previous studies it was suggested that the major source of these two hormones, during June, may be the interrenal gland (Gobetti and Zerani, 1991).

The high concentrations of plasma PGE₂ and androgens in frogs, during the pre-reproductive phase of the breeding cycle, suggest a causal relationship between these two hormones. The ability of PGE₂ to stimulate androgen synthesis *in vivo* and *in vitro* during the pre-reproductive phase of the breeding cycle supports this hypothesis. On the contrary, in May, this phenomenon was not observed in experiments *in vivo* and *in vitro* with frogs that had not ovulated. It is suggested that PGE₂, by stimulating androgen synthesis, inhibits breeding during the pre-reproductive phase, when the environmental conditions are unfavourable for the survival of young. Androgens could act at ovarian, pituitary or hypothalamic levels, as has been proposed for oestradiol (Pavgi and Licht, 1989; Fasano *et al.*, 1991). In this context, it should be noted that during the breeding and post-reproductive phases of the annual cycle, concentrations of plasma PGE₂ and androgens were low, in agreement with the low basal release of androgens from the ovaries *in vitro*, and with the observations that treatment with PGE₂ failed to

Fig. 6. Effects *in vitro* of 75 ng PGE₂ or 75 ng PGF_{2α} on androgen release from ovary of female frog, *Rana esculenta*, incubated in (a) April (pre-reproduction), (b) May (reproduction), (c) June (post-reproduction) and (d) October (recovery). Experimental groups: (■) ovary incubated with medium alone; (□) ovary incubated with PGE₂; (▨) ovary incubated with PGF_{2α}. Each mean refers to seven determinations $\pm$ SD. a, $P < 0.01$ versus same time in May, June and October; b, $P < 0.01$ versus same time in May and June; * $P < 0.01$ versus same time with medium-alone (Duncan's multiple range test).

stimulate the release of androgens. The decrease in oestradiol secretion observed in April *in vitro*, after treatment with PGE₂, could be due to the enhanced synthesis of androgens.

The positive correlation between the seasonal change in concentration of plasma PGE₂ and ovarian weights is at present still unclear. In this context, it should be noted that the maturation of the amphibian oocytes is modulated by various hormones that control the amounts of 3',5'-cyclic monophosphate (cAMP) in the follicular cells. A receptor for PGE is present on the follicular cell membrane and mediates an increase in cAMP concentrations (Mori *et al.*, 1989).

PGF_{2α} injection *in vivo* in June and October increased the release of oestradiol, although experiments *in vitro* suggested that PGF_{2α} stimulated oestradiol secretion only from ovaries collected in October. The reason for this discrepancy may be that the target tissue for PGF_{2α} is the interrenal in June (Gobbetti and Zerani, 1991) and the ovary in October. During the post-reproductive phase the release of oestradiol from the interrenal glands by PGF_{2α} may be responsible for inhibiting ovarian function by exerting a negative feedback action at the ovarian, pituitary or hypothalamic levels (Pavgi and Licht, 1989; Fasano *et al.*, 1991). This phenomenon is an adaptive

Fig. 7. Effects *in vitro* of 75 ng PGE₂ or 75 ng PGF_{2α} on oestradiol release from ovary of female frog, *Rana esculenta*, incubated in (a) April (pre-reproduction), (b) May (reproduction), (c) June (post-reproduction) and (d) October (recovery). Experimental groups: (■) ovary incubated with medium alone; (□) ovary incubated with PGE₂; (▨) ovary incubated with PGF_{2α}. Each mean refers to seven determinations $\pm$ SD. a, $P < 0.01$ versus same time in April, May and October; * $P < 0.01$ versus same time with medium-alone (Duncan's multiple range test).

mechanism to prevent the presence of tadpoles in autumn, when the environmental conditions are unfavourable for their survival (Salthe and Mecham, 1974). The ability of PGF_{2α} to stimulate ovarian oestradiol synthesis in October is not adequately supported by evidence of increased circulating oestradiol concentrations at this time of year and the meaning of this phenomenon is unclear.

This study indicates that in *Rana esculenta* one week in captivity modified PGs and sex steroids. However, stress-related decreases in the secretion of these hormones were also reported in *Rana esculenta* (Paolucci *et al.*, 1991; Zerani *et al.*, 1991), in *Rana catesbeiana* (Licht *et al.*, 1983), in rough skinned newt, *Taricha granulosa* (Moore & Zoeller, 1985), and in crested newt, *Triturus cristatus* (Gobbetti *et al.*, 1991b, c). It is unlikely that the effects of PGs on steroidogenesis were merely a consequence of captivity as effects of PGs on steroidogenesis were observed irrespective of whether captivity affected steroidogenesis, in this and another amphibian species. Thus PGF_{2α} induced an increase in plasma oestradiol in October, when captivity did not affect the secretion of this steroid, in agreement with findings in male *Rana esculenta* (Gobbetti *et al.*, 1992a). In addition, in *Triturus cristatus*, treatment with PGE₂ decreased androgen secretion, which was modified by captivity (Gobbetti and Zerani, 1992b), whereas a progesterone secretion which was not modified by

captivity was increased after treatment with PGE₂ (Gobbetti *et al.*, 1992b).

In summary, this work indicates that seasonal changes in the concentration of these two PGs in blood may regulate the timing of breeding in *Rana esculenta* ensuring young are produced in the spring; PGE₂ inhibits reproductive function in April by stimulating androgen secretion, and PGF_{2α} may have a similar function in October by stimulating oestradiol secretion.

This work was supported by a grant to A. Gobbetti from M.U.R.S.T.

References

- Dharmarajan AM, Sueoka K, Miyazaki S, Atlas SJ, Ghodgaonkar RB, Dubin NH, Zirkin BR and Wallach EE (1989) Prostaglandins and progesterone secretion in the *in vitro* perfused pseudopregnant rabbit ovary *Endocrinology* **124** 1198–1203
- d'Istria M, Delrio G, Botte V and Chieffi G (1974) Radioimmunoassay of testosterone, 17β-oestradiol and estrone in the male and female plasma of *Rana esculenta* during sexual cycle *Steroids and Lipids Research* **5** 42–48
- Duncan DB (1955) Multiple range and multiple F test *Biometrics* **11** 1–42
- Espey LL, Norris C and Saphire D (1986) Effect of time and dose of indomethacin on follicular prostaglandins and ovulation in the rabbit *Endocrinology* **119** 746–754
- Fasano S, D'Antonio M and Pierantoni R (1991) Sites of action of local oestradiol feedback mechanism in the frog (*Rana esculenta*) testis *General and Comparative Endocrinology* **81** 492–499
- Gobbetti A and Zerani M (1991) Gonadotropin-releasing hormone stimulates biosynthesis of prostaglandin F_{2α} by the interrenal gland of the water frog, *Rana esculenta*, *in vitro* *General and Comparative Endocrinology* **84** 434–439
- Gobbetti A and Zerani M (1992a) PGF_{2α}, PGE₂ and sex steroids from the abdominal gland of the male crested newt *Triturus cristatus* (Laur.) *Prostaglandins* **43** 101–109
- Gobbetti A and Zerani M (1992b) Prostaglandin E₂ and reproduction in the male crested newt, *Triturus cristatus* (Laur.) *Journal of Experimental Zoology* **263** 186–193
- Gobbetti A, Zerani M, Carnevali O and Botte V (1990) Prostaglandin F_{2α} in female water frog, *Rana esculenta*: plasma levels during the annual cycle and effects of exogenous PGF_{2α} on circulating sex hormones *General and Comparative Endocrinology* **80** 175–180
- Gobbetti A, Zerani M, Bolelli GF and Botte V (1991a) Seasonal changes in plasma prostaglandin F_{2α} and sex hormones in the male water frog, *Rana esculenta*. *General and Comparative Endocrinology* **82** 331–336
- Gobbetti Z, Zerani M, Bellini-Cardellini L and Botte V (1991b) Plasma prostaglandin F_{2α} and reproduction in the female newt *Triturus cristatus* (Laur.) *Prostaglandins* **42** 269–277
- Gobbetti A, Zerani M and Botte V (1991c) Plasma prostaglandin F_{2α} in the male *Triturus cristatus* (Laur.) during the reproductive annual cycle and effects of exogenous prostaglandin on sex hormones *Prostaglandins* **41** 67–74
- Gobbetti A, Zerani M and Bellini-Cardellini L (1992a) A possible role of prostaglandin E₂ in reproduction of the male water frog, *Rana esculenta*: *in vivo* and *in vitro* studies *Prostaglandins* **44** 277–289
- Gobbetti A, Zerani M and Botte V (1992b) A possible involvement of prostaglandin E₂ in the reproduction of female crested newt, *Triturus cristatus* *Zoological Science* **9** 639–648
- Greenhalgh EA (1990) Luteal steroidogenesis and regression in the rat: progesterone secretion and lipid peroxidation induced in luteal cells by human chorionic gonadotrophin, phospholipase A₂ and prostaglandin F_{2α} *Journal of Endocrinology* **125** 397–402
- Guillette LJ Jr, Lavia LA, Walker NJ and Roberts DK (1984) Luteolysis induced by prostaglandin F_{2α} in the lizard, *Anolis carolinensis* *General and Comparative Endocrinology* **56** 271–277
- Guillette LJ Jr, Gross TS, Matter JM and Palmer BD (1990) Arginine vasotocin-induced prostaglandin synthesis *in vitro* by the reproductive tract of the viviparous lizard *Sceloporus jarrovi* *Prostaglandins* **39** 39–51
- Guillette LJ Jr, Bjorndal KA, Bolten AB, Gross TS, Palmer BD, Witherington BE and Matter JM (1991) Plasma oestradiol, progesterone, prostaglandin F_{2α} and prostaglandin E₂ concentration during natural oviposition in the loggerhead turtle (*Caretta caretta*) *General and Comparative Endocrinology* **82** 121–130
- Jones RE, Orlicky DJ, Austin HB, Rand MS and Lopez KH (1990) Indomethacin inhibits ovarian PGE secretion and gonadotropin-induced ovulation in a reptile (*Anolis carolinensis*) *Journal of Experimental Zoology* **255** 57–62
- Licht P, McCreery BR, Barnes R and Pang R (1983) Seasonal and stress related changes in plasma gonadotropins, sex steroids, and corticosterone in the bullfrog, *Rana catesbeiana* *General and Comparative Endocrinology* **50** 124–145
- Moore FL and Zoeller RT (1985) Stress-induced inhibition of reproduction: evidence of suppressed secretion of LH-RH in an amphibian *General and Comparative Endocrinology* **60** 252–258
- Mori K, Oka S, Tani A, Ito S and Watanabe Y (1989) E-series prostaglandins activate cAMP-mediated potassium currents in follicle-enclosed *Xenopus* oocyte *Biochemical and Biophysical Research Communications* **162** 1535–1540
- Paolucci M, Esposito V, Di Fiore MM and Botte V (1991) Effects of short post-capture confinement on plasma reproductive hormone and corticosterone profiles in *Rana esculenta* during the sexual cycle *Bollettino di Zoologia* **57** 253–259
- Pavgi S and Licht P (1989) Effects of gonadectomy and steroids on pituitary gonadotropin secretion in a frog, *Rana pipiens* *Biology of Reproduction* **40** 40–48
- Salthe SN and Mecham JS (1974) Reproductive and courtship patterns. In *Physiology of the Amphibia* pp 309–521 Ed. B. Lofts. Academic Press, New York
- Scossiroli RE and Palenzona DL (1979) *Manuale di Biometria*. Zanichelli, Bologna
- Sedrani SH and El-Banna AA (1987) Effect of indomethacin on plasma level of vitamin D metabolites, oestradiol and progesterone in rabbit during early pregnancy *Comparative Biochemistry and Physiology* **87** 635–639
- Shimada K, Olson DM and Etches RJ (1984) Follicular and uterine prostaglandin levels in relation to uterine contraction and the first ovulation of a sequence in the hen *Biology of Reproduction* **31** 76–82
- Sokal RR and Rohlf FJ (1981) *Biometry* (2nd Edn). Freeman, San Francisco
- Zerani M, Amabili F, Mosconi G and Gobbetti A (1991) Effects of captivity stress on plasma steroid levels in the green frog, *Rana esculenta*, during the annual reproductive cycle *Comparative Biochemistry and Physiology* **98** 491–496