Correlation of growth hormone secretion during pregnancy with circulating prolactin in rats

G. A. Jahn¹, A. M. Rastrilla² and R. P. Deis^{1*}

¹Laboratorio de Reproduccion y Lactancia, CRICYT-CONICET, Casilla de Correos 855, 5500 Mendoza, Argentina; and ²Facultad de Bioquímica, Química y Farmacia, Universidad Nacional de San Luis, San Luis, Argentina

Growth hormone (GH) concentrations were measured throughout pregnancy in rats. The effects of surgical stress, ovariectomy, and treatments with the antiprogesterone mifepristone (RU 486) or the antioestrogen tamoxifen on serum GH, progesterone and prolactin were studied. GH concentrations were low during the first 18 days of pregnancy, except on the morning of day 5, and increased progressively from day 19 reaching peak values on the mornings of days 21 and 22. Thereafter GH concentrations decreased progressively, reaching very low values at 24.00 h on day 22, in parallel with a rise in serum prolactin concentrations. Surgical stress, performed at 12.00 h on day 20 of pregnancy, diminished serum GH concentrations 10 min later, but these returned to values similar to those of the non-operated rats 1–24 h later. Surgical stress did not modify serum prolactin concentrations at any time. Ovariectomy performed on the morning of day 19 produced the expected fall in serum progesterone and a rise in prolactin which lasted until the night of day 20. Serum GH concentrations were significantly diminished with respect to controls on day 20 and the morning of day 21 and then increased. Treatment with mifepristone on day 19 produced a simultaneous rise in serum prolactin and a fall in serum progesterone and GH by 08.00 h on day 21. Treatment with tamoxifen on days 3 and 4, or given daily from day 17 onwards did not modify prolactin concentrations but diminished serum GH concentrations at 08.00 on day 5 and on days 19–22, with the exception of a peak on day 22 (08.00 h). Tamoxifen also decreased serum progesterone concentrations. These results show that pregnant rats have a reduced capacity of response to stress in terms of changes in GH and prolactin secretion. There are high serum concentrations of GH at the end of pregnancy. The regulation of GH secretion at this time is different from that of prolactin and does not seem to depend on the fall in progesterone concentrations. However, serum GH concentrations seem to be inversely correlated with serum prolactin concentrations, as they tended to fall after increases in prolactin above basal concentrations. Oestrogen may also have a stimulatory role on GH since administration of an anti-oestrogen also resulted in a fall in GH concentrations in spite of reduced prolactin secretion.

Introduction

Regulation of growth hormone (GH) secretion shows some similarities to and some differences from prolactin secretion. GH is secreted in an ultradian fashion as several daily peaks, and there is a marked sexual dimorphism which is caused mainly by sex steroids (Tannenbaum and Martin, 1976; Eden, 1979; Jansson *et al.*, 1984, 1985). GH secretion is stimulated by GH-releasing hormone, inhibited by somatostatin, which also inhibits prolactin, and GH and prolactin are stimulated by oestrogens (Jansson *et al.*, 1985; Hall *et al.*, 1986). Glucocorticoids have dual actions on GH release and inhibit prolactin release (Leung *et al.*,

1980; Nakagawa *et al.*, 1987a, b). Both hormones are also modulated by neurotransmitters; serotonin is mainly stimulatory for both (Martin *et al.*, 1978; Willoughby *et al.*, 1987), whereas catecholamines have opposite effects on GH and prolactin secretion (Martin *et al.*, 1978; Eden *et al.*, 1979; Willoughby and Day, 1981; Crowley *et al.*, 1982). Stress also has opposite effects, stimulating prolactin release and inhibiting GH (Schalch and Reichlin, 1966; Takahashi *et al.*, 1971; Krulich *et al.*, 1974; Terry *et al.*, 1976; Eden, 1978). Finally, the suckling stimulus induces secretion of prolactin and GH, although the latter only in a transient manner (Sar and Meites, 1969; Saunders *et al.*, 1976; Nagy *et al.*, 1986).

In rats, the fall in circulating progesterone at the end of pregnancy is followed by an increase in serum prolactin (Vermouth and Deis, 1972, 1974; Bussmann and Deis, 1979) which is

*Correspondence.
Received 20 May 1992.

mediated by adrenergic but not serotonergic pathways (Jahn and Deis, 1987, 1988, 1991). In the present study, we investigated the circulating concentrations of GH throughout pregnancy in rats and the regulation of GH by stress and gonadal hormones. Previous studies performed on selected days of pregnancy (Saunders *et al.*, 1976; Klindt *et al.*, 1981; Carlsson *et al.*, 1990) showed that there is an increase in GH secretion in the last days of pregnancy. We, therefore, studied the profile of GH secretion at different times on days 4–6, 11–13 and 18–22 of pregnancy. We also investigated whether procedures known to advance prolactin secretion through a fall in or blockade of progesterone (such as ovariectomy or treatment with the anti-progesterone mifepristone) could modify GH secretion at the end of pregnancy. In addition, we studied the effect of tamoxifen administration, as oestrogens appear to stimulate GH secretion (Simard *et al.*, 1986; Ho *et al.*, 1987). Some of these results have been presented in abstract form (Jahn *et al.*, 1987).

Materials and Methods

Animals

Virgin female rats, three to four months old (200–220 g) bred in our laboratory and originally of the Wistar strain, were used. The rats were kept in a light (lights on 06.00–20.00 h)- and temperature (22–24°C)-controlled room; rat chow (Cargill, Buenos Aires and Nutric, Cordoba) and tap water were available *ad libitum*. Vaginal smears were taken daily. Rats were caged individually with fertile males on the night of pro-oestrus, and the presence of spermatozoa was checked in the vaginal smear the following morning. This day was designated day 1 of pregnancy. In our laboratory, rats usually give birth on day 23. All the rats were handled daily, to minimize the effect of handling stress. The experiments were performed in accordance with the NIH Guide for the Care and Use of Laboratory Animals (NIH publication No. 86-23, revised 1985).

Experimental procedures

To study GH concentrations throughout pregnancy, serum samples were obtained after decapitation or cardiac puncture of conscious animals at 08.00, 12.00, 16.00, 20.00 and 24.00 h on days 4, 5, 6, 11, 12, 13, 18, 19, 20, 21 and 22 of pregnancy. Rats were bled once by cardiac puncture on days 4–6 or 11–13 and the second time after decapitation on days 19 to 22. All the groups sampled on days 4–6 and 11–13 included samples from decapitated rats, whose hormone concentration did not differ from the samples obtained by cardiac puncture. These samples can be considered as from unstressed rats, since they were obtained within 15 s and serum prolactin values obtained in these rats were not different from those obtained in decapitated rats from the same experimental groups. Blood was allowed to clot at room temperature; the serum was separated and stored frozen at –30°C until assayed for GH and prolactin.

Surgical procedures

Ovariectomy was performed through bilateral incisions under ether anaesthesia between 08.00 and 09.00 h on day 19

of pregnancy. Groups of intact or sham-operated rats were used as controls, and since the hormone concentrations of both groups did not differ, they were considered as one group for statistical evaluation and expression of results.

To determine the effect of the surgical stress on serum GH and prolactin concentrations, a group of daily handled pregnant rats were sham operated (laparotomized) at 12.00 h on day 20 of pregnancy under ether anaesthesia. Groups of rats were decapitated 10 min, 1 h, 4 h and 24 h after surgery and blood was obtained for GH and prolactin determinations. The values obtained were compared with those of intact pregnant rats obtained at the same times of day.

Drug treatments

Mifepristone (RU 38486, donated by Roussel-Uclaf, Romainville, France) dissolved in sunflower seed oil (2 g l⁻¹) was injected s.c. at a dose of 2 mg kg⁻¹ at 08.00 h on day 19 of pregnancy.

Tamoxifen (ICI 46474, donated by Gador, Buenos Aires) was dissolved in 0.5% Tween 80 (50 mg l⁻¹) and given per os at a dose of 0.5 mg kg⁻¹ to a group of rats at 08.00 and 18.00 h on days 3 and 4 of gestation. The rats were decapitated at 08.00 h on day 5 and trunk blood collected for hormone determinations.

For the experiments at the end of pregnancy, tamoxifen was dissolved in sunflower seed oil (1 g l⁻¹) and injected daily at a dose of 200 µg per rat at 09.00 h from day 17 of pregnancy until the morning the rats were killed.

Blood samples were obtained from groups of 8–10 animals subjected to the different treatments at 12.00, 16.00 or 20.00 h on day 19 and 08.00, 12.00, 16.00 or 20.00 h on days 20, 21 or 22. The first sample was obtained by cardiac puncture on days 19 or 20 and the second after decapitation at least 36 h later. In the mifepristone-treated rats, the last sample was obtained at 08.00 h on day 22 as by this time most of these animals had already given birth.

Prolactin, progesterone and GH determinations

Prolactin and GH were measured by double antibody radioimmunoassays using materials generously provided by the NIADDK (S. Raiti, NIADDK Rat Pituitary Hormone Distribution Program). The hormones were radio-iodinated using the chloramine T method and purified by passage through Sephadex G75. The results were expressed in terms of the rat prolactin RP-3 and rat GH RP-2 standard preparations. Assay sensitivity for both hormones was 0.5 ng ml⁻¹ serum and the inter- and intra-assay coefficients of variation were less than 10%.

Serum progesterone was measured using a radioimmunoassay developed in our laboratory (Bussmann and Deis, 1979) with an antiserum raised in rabbits against progesterone-11-BSA conjugate. Assay sensitivity was less than 5 ng ml⁻¹ serum and the inter- and intra-assay coefficients of variation were less than 10%.

Statistical analysis

Statistical analysis was performed using one- or two-way analysis of variance followed by the least significant difference

Fig. 1. Concentrations of growth hormone in serum during pregnancy in rats. Results are the means $\pm$ SEM of six determinations for days 4–18 and of 10–15 on days 19–22. *Significantly different ($P < 0.05$) from the means of days 4–18 (excluding day 5, 08.00 h).

Fig. 2. Effect of sham surgery (laparotomy) under ether anaesthesia performed on rats at 12.00 h on day 20 of pregnancy on serum GH and prolactin (PRL) concentrations (■) at various times after surgery. Comparison with serum hormone concentrations in intact rats (□) sampled at the same times. Results are expressed as means $\pm$ SEM of groups of five to six rats per sampling time. *Significantly different ($P < 0.05$) from control results.

between means test when more than two experimental groups were compared or by Duncan's multiple range test for analysis of the significance between different groups during pregnancy or after surgery (Snedecor and Cochran, 1967).

Results

Effect of surgical stress on serum GH and prolactin concentrations in pregnant rats

To ascertain the effect of surgery alone on serum GH and prolactin, groups of 5–6 rats were bled by decapitation 10 min,

1 h, 4 h and 24 h after sham surgery (laparotomy) under ether anaesthesia performed at 12.00 h on day 20 of pregnancy, when serum GH concentrations are high (Fig. 1) and compared with values for non-operated rats decapitated at the same times. Ten minutes after surgery there was a significant decrease in serum GH concentrations, whereas 1–24 h after surgery, values were not different from the non-operated group (Fig. 2). There were no significant changes in serum prolactin concentrations at any of the times studied (Fig. 2). As blood samples in the ovariectomized rats were taken from 4 h after surgery, the observed changes in serum GH and prolactin values were due to the different treatments and not to surgical stress.

Table 1. Serum GH, prolactin and progesterone concentrations (ng ml⁻¹) in rats on days 19–22 of pregnancy. Effects of ovariectomy, treatment with mifepristone or tamoxifen

Day of gestation and time of day when samples were taken	Treatments					
	Controls			Ovariectomy		
	GH	Prolactin	Progesterone	GH	Prolactin	Progesterone
<i>Day 19</i>						
12.00 h	15.7 ± 1.6	8.0 ± 1.8	64.3 ± 3.1	10.4 ± 1.6	10.6 ± 2.2	17.7 ± 2.0*
16.00 h	10.8 ± 1.3	5.6 ± 1.0	55.5 ± 2.9	16.8 ± 1.6	7.0 ± 1.9	13.7 ± 3.4*
20.00 h	22.2 ± 3.1	3.3 ± 0.2	54.7 ± 4.9	11.1 ± 1.3*	13.4 ± 4.2*	7.6 ± 1.0*
<i>Day 20</i>						
8.00 h	28.0 ± 7.0	7.1 ± 1.2	62.1 ± 4.3	12.0 ± 1.1*	24.5 ± 2.5*	10.3 ± 1.0*
12.00 h	32.2 ± 6.0	6.4 ± 1.1	46.6 ± 3.8	13.6 ± 4.3*	28.2 ± 7.2*	4.0 ± 1.1*
16.00 h	27.8 ± 3.6	9.1 ± 0.8	70.3 ± 3.0	15.1 ± 3.7*	21.4 ± 6.2*	7.2 ± 0.9*
20.00 h	30.1 ± 4.8	5.8 ± 0.6	61.5 ± 5.6	11.6 ± 1.8*	12.5 ± 1.0*	4.9 ± 0.7*
<i>Day 21</i>						
8.00 h	53.0 ± 5.3	7.5 ± 0.5	58.3 ± 3.1	10.7 ± 1.1*	10.4 ± 2.0	7.6 ± 1.3*
12.00 h	36.6 ± 3.3	5.4 ± 0.6	42.0 ± 4.2	34.2 ± 6.7	7.0 ± 1.0	ND
16.00 h	42.1 ± 7.6	7.6 ± 0.8	50.0 ± 5.0	49.5 ± 13.8	8.5 ± 1.3	ND
20.00 h	28.7 ± 2.7	5.7 ± 0.5	48.5 ± 2.2	30.6 ± 9.3	10.2 ± 3.4	14.2 ± 0.5*
<i>Day 22</i>						
8.00 h	29.2 ± 2.1	9.3 ± 0.4	41.2 ± 7.6	25.1 ± 9.3	12.5 ± 4.2	10.2 ± 2.8*
12.00 h	54.0 ± 11.0	15.6 ± 6.0	23.3 ± 4.7**	24.4 ± 7.1	6.9 ± 0.8	ND
16.00 h	25.1 ± 3.8**	36.4 ± 4.5**	18.8 ± 2.8**	28.8 ± 7.9	8.1 ± 1.2	ND
20.00 h	12.7 ± 1.5**	22.4 ± 6.4**	12.8 ± 2.1**	21.2 ± 4.6	10.1 ± 1.7*	7.6 ± 1.8
Mifepristone			Tamoxifen			
	GH	Prolactin	Progesterone	GH	Prolactin	Progesterone
<i>Day 19</i>						
12.00 h	21.8 ± 1.9	6.6 ± 1.2	59.6 ± 4.3	15.1 ± 2.2	3.9 ± 0.6	43.0 ± 2.9*
16.00 h	24.0 ± 4.0*	3.2 ± 0.8	47.4 ± 2.7	11.7 ± 2.7	5.1 ± 0.5	50.8 ± 3.4
20.00 h	16.2 ± 5.5	6.1 ± 0.9	37.5 ± 4.6	10.8 ± 1.4*	3.9 ± 0.5	29.7 ± 1.7*
<i>Day 20</i>						
8.00 h	35.9 ± 6.7	6.9 ± 0.9	51.4 ± 3.6	4.9 ± 0.7*	4.7 ± 0.8	32.2 ± 2.0*
12.00 h	41.7 ± 14.5	7.4 ± 0.6	36.3 ± 4.6*	17.7 ± 3.0*	2.6 ± 0.5	29.2 ± 2.9*
16.00 h	21.8 ± 5.9	12.5 ± 3.6	36.1 ± 4.8*	12.6 ± 2.4*	2.1 ± 0.5*	53.1 ± 3.8*
20.00 h	24.7 ± 4.0	6.0 ± 1.0	40.6 ± 3.3	9.7 ± 1.6*	5.2 ± 1.7	30.1 ± 6.7*
<i>Day 21</i>						
8.00 h	12.8 ± 2.2*	53.1 ± 8.8*	25.3 ± 4.1*	12.0 ± 3.6*	4.4 ± 1.4	27.6 ± 4.1*
12.00 h	16.4 ± 4.4*	28.7 ± 7.2*	13.7 ± 3.1*	13.9 ± 3.7*	5.5 ± 1.2	25.2 ± 8.3*
16.00 h	8.3 ± 2.2*	25.3 ± 11.4*	8.8 ± 2.1*	15.9 ± 3.1*	6.5 ± 1.2	8.7 ± 1.6*
20.00 h	6.0 ± 1.6*	23.6 ± 5.2*	12.8 ± 1.8*	14.3 ± 4.0*	14.6 ± 8.5	7.8 ± 1.4*
<i>Day 22</i>						
8.00 h	11.6 ± 2.5*	34.4 ± 8.4*	8.6 ± 0.6*	21.1 ± 5.0	3.0 ± 1.7	15.9 ± 3.1*
12.00 h	Abortion	Abortion	Abortion	25.6 ± 7.5*	5.5 ± 0.5	10.9 ± 4.5*
16.00 h				25.8 ± 6.2	1.8 ± 0.4*	7.4 ± 1.3*
20.00 h				4.8 ± 0.9*	4.6 ± 1.2*	7.2 ± 0.6

Results are the means ± SEM of 8–15 determinations at each time point.

P* < 0.05 compared with controls; *P* < 0.05 compared with values on day 21.

ND: not determined.

Serum GH concentrations during pregnancy

In the present study we have not attempted to study the ultradian rhythm that has been repeatedly described for GH secretion (Saunders *et al.*, 1976; Klindt *et al.*, 1981). Average values obtained in different rats sampled at the same hour are given and represent mean secretion of GH. Serum GH concentrations remained below 20 ng ml⁻¹ during days 4–6 and 11–13 of pregnancy, except on

the morning of day 5 (08.00 h) when values increased significantly to 60 ng ml⁻¹ (Fig. 1). Values were low on day 18, but from day 19 onwards, serum GH concentrations increased progressively, reaching concentrations as high as 53 ng ml⁻¹ at 08.00 h on day 21 with a new peak at 12.00 h on day 22, followed by a marked decrease during the afternoon, to a minimum of 10 ng ml⁻¹ at midnight of day 22, which was coincident with an increase in serum prolactin concentrations and a fall in progesterone (see Table 1).

Effect of ovariectomy on serum concentrations of GH, prolactin and progesterone

Ovariectomy performed on day 19 changed the rhythm of serum GH secretion, decreasing the values with respect to controls at 20.00 h on day 19, throughout day 20 and up to 08.00 h on day 21 of pregnancy. There was an acute increase at 12.00 h on day 21 to values similar to those of control rats; thereafter values remained high and not significantly different from control values (Table 1).

Serum prolactin concentrations measured at different times on days 19–22 in intact pregnant rats remained below 10 ng ml^{-1} until the morning of day 22, and increased significantly thereafter. Serum progesterone concentrations remained between 40 and 70 ng ml^{-1} until the morning of day 22, and then declined progressively throughout the rest of day 22. Ovariectomy produced an acute fall in serum progesterone concentrations, followed as expected (Vermouth and Deis, 1974) by significant increases in serum prolactin concentrations 8 h after surgery; prolactin concentrations remained high until 20.00 h on day 20. Ovariectomy also blocked the increase in serum prolactin observed on the afternoon of day 22 in control rats.

The increases in prolactin secretion produced by the fall in serum progesterone concentrations were accompanied by significant decreases in serum GH concentrations with respect to intact pregnant rats and, conversely, serum GH concentrations increased after serum prolactin had returned to basal (less than 7.5 ng ml^{-1}) concentrations.

Effect of mifepristone administration

Mifepristone administration induced a slight increase in serum GH concentration at 16.00 h on day 19 but did not modify them with respect to control rats at the other times measured until 20.00 h on day 20; serum concentrations of GH were significantly lower on days 21 and 22. These rats gave birth by 08.00 h on day 22. Serum progesterone concentrations started to decline in the mifepristone-treated rats by 12.00 h on day 20, reaching basal values 24 h later, whereas prolactin concentrations were low on day 19, but increased by 08.00 h on day 21 and remained high until 20.00 h on day 21 (Table 1). In this group of rats, serum GH concentrations were again significantly reduced with respect to control rats coincident with the increase in serum prolactin.

Effect of tamoxifen

The group of rats that had been given tamoxifen on days 3 and 4 of gestation showed a mean serum GH concentration of $15 \pm 4 \text{ ng ml}^{-1}$ ($n = 9$) after decapitation at 08.00 h on day 5, which was significantly lower ($P < 0.001$, Student's *t* test) than the values observed at the same time in the intact pregnant rats (Fig. 1).

Daily administration of tamoxifen starting on day 17 significantly reduced serum GH concentration, except for peaks at 08.00 h and 16.00 h on day 22 (Table 1). Serum prolactin concentrations in tamoxifen-treated rats were low at all times

studied and the increase seen in controls on day 21 was also blocked (Table 1). Serum progesterone concentrations were lower than those of controls at most of the times sampled, and reached values similar to those of controls on the afternoon of day 22, and on the afternoon of day 21. None of the rats had given birth by 20.00 h on day 22, but examination of the uterine content after decapitation on day 22 showed only dead and partially digested fetuses.

Discussion

Any study of GH concentrations in the blood should involve stress-free methods of obtaining samples since handling rats may cause a fall in the release of GH (Takahashi *et al.*, 1971; Terry *et al.*, 1976; Armario *et al.*, 1984; Jurcovicova *et al.*, 1984) and concentrations are extremely sensitive to anaesthetic. Thus, a brief 2 min exposure to ether has been reported to cause a significant fall in GH concentration (Schalch and Reichlin, 1966; Krulich *et al.*, 1974; Briski *et al.*, 1984). It is also known that major surgical procedures cause a reduction in GH release for two days (Obonsawin *et al.*, 1985) and even 4 days (Eden, 1978) after the operation. However, most previous studies have been performed on male or nonpregnant female rats, and female rats seem to be less sensitive to stress in terms of GH release, particularly when they are pregnant (Schalch and Reichlin, 1966). An interesting finding of the present work is that pregnant female rats seem to have a shorter response to stress in terms of GH release than has been reported for male rats, since after ether-laparotomy, serum GH concentrations were significantly reduced for only 10 min, whereas in male rats the duration of GH depression is longer (Eden, 1978; Obonsawin *et al.*, 1985). This may reflect a general insensitivity of pregnant rats to stress, since prolactin secretion was unaffected by ether-laparotomy, a procedure that readily increases secretion of this hormone in virgin animals (Boehm *et al.*, 1982). These results exclude the possibility that the modifications in hormone concentrations observed in our study were due to stress effects resulting from the methodology used.

Our results showed that a marked increase in serum GH concentrations occurred during the last 4 days of pregnancy, confirming and extending previous studies which were performed by serial sampling of individual rats on selected days of pregnancy (Saunders *et al.*, 1976; Klindt *et al.*, 1981; Carlsson *et al.*, 1990). This increase may be important in the final differentiation of the mammary gland before the initiation of milk secretion and may be related to the marked increase observed in prolactin and GH receptors in the liver observed in the last days of pregnancy (Husman *et al.*, 1985; Jahn *et al.*, 1991).

Regulation of GH release seems to be different to that of prolactin at the end of gestation. The increase observed on the last days of pregnancy could be a consequence of increased oestrogen secretion seen in rats from day 19 (Shaikh, 1971), as oestrogens have been shown to stimulate GH production by the pituitary (Simard *et al.*, 1986; Armario *et al.*, 1984). The high serum GH concentrations found on the morning of day 5 of pregnancy may also be a product of stimulation by oestrogens secreted on this day (Shaikh, 1971), that are necessary for implantation, since the high GH concentrations were inhibited by prior administration of tamoxifen. Furthermore, tamoxifen

administration in the last days of pregnancy produced significant decreases in serum GH and prolactin concentrations.

The changes in prolactin secretion followed the previously described increases observed after the progesterone fall induced by ovariectomy (Vermouth and Deis, 1974). Mifepristone administration did not induce an immediate increase in prolactin secretion (Deis *et al.*, 1989), but prolactin concentrations rose after 48 h as a consequence of the delayed fall in serum progesterone concentrations. Our results also indicate a role for oestrogens in the increase in prolactin secretion on day 22 (when serum progesterone concentrations are low; Bussmann and Deis, 1979; Nicholas and Hartmann, 1981), as ovariectomy, as well as tamoxifen administration, completely abolished this rise.

A potential stimulatory action for oestrogens and progesterone on the maintenance of luteal progesterone production in late pregnancy is indicated since both anti-hormones advanced the fall in serum progesterone. A role for both steroid hormones in the maintenance of luteal function in rats has already been described in early and mid-pregnancy (Gibori and Keyes, 1978, 1980; Rothchild, 1980; Kawano *et al.*, 1988; Singh *et al.*, 1988), and our results extend this role to the end of pregnancy.

There appears to be an inverse relationship between prolactin and GH secretion on the last days of pregnancy. In intact, pregnant rats the increase in GH secretion occurred 3 days before the release of prolactin, and when prolactin secretion increased, serum GH concentrations had already fallen. The increases in prolactin secretion observed after ovariectomy or mifepristone treatment were accompanied by decreases in serum GH, which returned to control concentrations when prolactin concentrations diminished. The reciprocal relationship between prolactin and GH was observed independent of experimental procedure or progesterone concentrations, except for the tamoxifen-treated rats, in which both hormones were depressed, indicating a stimulatory action of oestrogens for both hormones. The stimulatory action of oestrogens on GH release may be evident only in the presence of low concentrations of prolactin. In contrast, any action of progesterone seems to be mediated through its regulation of prolactin secretion.

It is known that many factors regulate GH secretion, but according to our results progesterone does not directly regulate GH release and oestrogen may not be the main factor involved. The inverse relationship between GH and prolactin secretion means that both hormones cannot be released simultaneously. The inhibitory effect of hyperprolactinaemia on GH secretion in female (Esquifino *et al.*, 1987a) and male rats (Esquifino *et al.*, 1987b) and in women (Andersen and Tabor, 1982; Ho *et al.*, 1985) has been reported already.

This work was supported by a grant (PID 9036/01) from the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina. The authors are indebted to F. E. Guinazu de Di Nasso for her excellent technical assistance. We thank A. F. Parlow and the NIADDK for the rat GH and prolactin kits.

References

- Andersen AN and Tabor A (1982) Prl, TSH, GH and LH responses to metoclopramide and breast-feeding in normal and hyperprolactinaemic women *Acta Endocrinologica* **100** 177–183
- Armario A, Castellanos JM and Balash J (1984) Dissociation between corticosterone and growth hormone adaptation to chronic stress in the rat *Hormone and Metabolic Research* **16** 142–145
- Boehm N, Plas-Roser S, Lazarus C and Aron C (1982) Influence of blood removal under ether anaesthesia on the release of prolactin during the oestrous cycle of the rat. Involvement of the ovaries *Acta Endocrinologica* **99** 179–186
- Briski KP, Quigley K and Meites J (1984) Counteraction by morphine of stress-induced inhibition of growth hormone release in the rat *Proceedings of the Society for Experimental Biology and Medicine* **177** 137–142
- Bussmann LE and Deis RP (1979) Studies concerning the hormonal induction of lactogenesis by prostaglandin $F_{2\alpha}$ in pregnant rats *Journal of Steroid Biochemistry* **11** 1485–1489
- Carlsson L, Eden S and Jansson J-O (1990) The plasma pattern of growth hormone in conscious rats during late pregnancy *Journal of Endocrinology* **124** 191–198
- Crowley W, Terry LC and Johnson MD (1982) Evidence for the involvement of central epinephrine systems in the regulation of luteinizing hormone, prolactin, and growth hormone release in female rats *Endocrinology* **110** 1102–1107
- Deis RP, Carrizo DG and Jahn GA (1989) Suckling-induced prolactin release potentiates mifepristone-induced lactogenesis in pregnant rats *Journal of Reproduction and Fertility* **87** 147–153
- Eden S (1978) The secretory pattern of growth hormone. An experimental study in the rat *Acta Physiologica Scandinavica Supplement* **458** 1–54
- Eden S (1979) Age- and sex-related differences in episodic growth hormone secretion in the rat *Endocrinology* **105** 555–560
- Eden S, Bolle P and Modigh K (1979) Monoaminergic control of episodic growth hormone secretion in the rat: effects of reserpine, α -methyl- p -tyrosine, p -chlorophenylalanine, and haloperidol *Endocrinology* **105** 523–529
- Esquifino AI, Vilanova MA and Agrasal C (1987a) Possible role of prolactin in growth regulation *Revista Espanola de Fisiologia* **43** 455–462
- Esquifino AI, Fernandez-Ruiz JJ, Cebeira M, Agrasal C, Tresguerres JAF and Ramos JA (1987b) Effect of experimentally induced hyperprolactinemia on growth hormone secretion *Revista Espanola de Fisiologia* **43** 463–468
- Gibori G and Keyes L (1978) Role of intraluteal estrogen in the regulation of the rat corpus luteum during pregnancy *Endocrinology* **102** 1176–1182
- Gibori G and Keyes L (1980) Luteotrophic effect of estrogen in the early pregnancy in the rat *Endocrinology* **106** 1584–1588
- Hall TR, Harvey S and Scanes CG (1986) Control of growth hormone secretion in the vertebrates: a comparative survey *Comparative Biochemistry and Physiology* **84A** 231–253
- Ho KY, Smythe GA and Lazarus L (1985) Enhanced hypothalamic dopaminergic inhibition of LH, TSH and GH release in patients with pathological hyperprolactinaemia *Acta Endocrinologica* **108** 289–296
- Ho KY, Evans WS, Blizzard RM, Veldhuis JD, Merriam GR, Samojlik E, Furlanetto E, Ropol AD, Kaiser DL and Thorner MO (1987) Effects of sex and age on the 24 hour profile of growth hormone secretion in man: importance of endogenous estradiol concentration *Journal of Clinical Endocrinology and Metabolism* **64** 51–58
- Husman B, Andersson G, Norstedt G and Gustafsson J-A (1985) Characterization and subcellular distribution of the somatogenic receptor in rat liver *Endocrinology* **116** 2605–2611
- Jahn GA and Deis RP (1987) A possible dual regulation of prolactin release by the serotonergic system in pro-oestrous and late pregnant rats: role of the ovarian hormones *Journal of Endocrinology* **112** 367–374
- Jahn GA and Deis RP (1988) Effect of serotonin antagonists on prolactin and progesterone secretion. Evidence that the stimulatory and inhibitory actions of serotonin upon prolactin release may be mediated through different receptors *Journal of Endocrinology* **117** 415–422
- Jahn GA and Deis RP (1991) The involvement of the adrenergic system on the release of prolactin and lactogenesis at the end of pregnancy in the rat *Journal of Endocrinology* **129** 343–350
- Jahn GA, Rastrilla AM and Deis RP (1987) Modificaciones de los niveles sericos de GH durante la preñez en la rata *Medicina* **47** Abstract 365
- Jahn GA, Edery M, Belair L and Djiane J (1991) Prolactin gene expression in rat mammary gland and liver during pregnancy and lactation *Endocrinology* **128** 2976–2984
- Jansson JO, Ekberg S, Isaksson O and Eden S (1984) Influence of gonadal steroids on age- and sex-related secretory patterns of growth hormone in the rat *Endocrinology* **114** 1287–1294
- Jansson JO, Eden S and Isaksson O (1985) Sexual dimorphism in the control of growth hormone secretion *Endocrine Reviews* **6** 128–150

- Jurcovicova J, Vigas M, Klir P and Jezova D (1984) Response of prolactin, growth hormone and corticosterone secretion to morphine administration or stress exposure in Wistar-AVN and Long Evans rats *Endocrinologia Experimentalis* **18** 209–214
- Kawano T, Okamura H, Tajima C, Fukuma K and Hatabuchi H (1988) Effect of RU 486 on luteal function in the early pregnant rat *Journal of Reproduction and Fertility* **83** 279–285
- Klindt J, Robertson MC and Friesen HG (1981) Secretion of placental lactogen, growth hormone, and prolactin in late pregnant rats *Endocrinology* **109** 1492–1495
- Krulich L, Hefco E, Illner P and Read CB (1974) The effects of acute stress on the secretion of LH, FSH, prolactin and GH in the normal male rat, with comments on their statistical evaluation *Neuroendocrinology* **16** 293–311
- Leung FC, Chen HT, Verkaik SJ, Steger RW, Peluso JJ, Campbell GA and Meites J (1980) Mechanism(s) by which adrenalectomy and corticosterone influence prolactin release in the rat *Journal of Endocrinology* **87** 131–140
- Martin JB, Duran D, Gurd W, Faille G, Audet J and Brazeau P (1978) Neuropharmacological modulation of episodic growth hormone and prolactin secretion in the rat *Endocrinology* **102** 106–113
- Nagy G, Kacsóh B, Kanyicska B, Toth BE and Korausz E (1986) Separation and suckling-induced changes in serum growth hormone levels of lactating rats and their pups *Endocrinologia Experimentalis* **20** 217–222
- Nakagawa K, Akikawa K, Matsubara M, Kubo M, Ishizuka T and Obara T (1987a) Biphasic effects of dexamethasone on growth hormone release *in vitro* *Endocrinologia Japonica* **34** 947–953
- Nakagawa K, Ishizuka T, Obara T, Matsubara M and Akikawa K (1987b) Dichotomic action of glucocorticoids on growth hormone secretion *Acta Endocrinologica* **116** 165–171
- Nicholas PR and Hartmann PE (1981) Progesterone control of the initiation of lactose synthesis in the rat *Australian Journal of Biological Sciences* **34** 435–443
- Obonsawin MC, Shin SH and Arrowsmith J (1985) Surgery depresses pulsatile growth hormone release in rats for up to 2 days *Acta Endocrinologica* **110** 42–45
- Rothchild I (1980) The regulation of the mammalian corpus luteum *Recent Progress in Hormone Research* **37** 183–298
- Sar M and Meites J (1969) Effects of suckling on pituitary release of prolactin, GH and TSH in postpartum lactating rats *Neuroendocrinology* **4** 25–31
- Saunders A, Terry LC, Audet J, Brazeau P and Martin JB (1976) Dynamic studies of growth hormone and prolactin secretion in the female rat *Neuroendocrinology* **21** 193–203
- Schalch DS and Reichlin S (1966) Plasma growth hormone concentration in the rat determined by radio-immunoassay: influence of sex, pregnancy, lactation, anesthesia, hypophysectomy and extracellular pituitary transplants *Endocrinology* **79** 275–280
- Shaikh A (1971) Estrone and estradiol levels in ovarian venous blood from rats during the estrous cycle and pregnancy *Biology of Reproduction* **5** 297–307
- Simard J, Hubert JF, Hosseinzadeh T and Labrie F (1986) Stimulation of growth hormone release and synthesis by estrogens in rat anterior pituitary cells in culture *Endocrinology* **119** 2004–2011
- Singh G, Singh MM, Maitra SC, Elger W, Kalra V, Upadhyay SN, Chowdhury SR and Kamboj VP (1988) Luteolytic action of two antiprogesterone agents (RU 486 and ZK-93734) in the rat *Journal of Reproduction and Fertility* **83** 73–83
- Snedecor GW and Cochran WG (1967) *Statistical Methods*. Iowa State University Press
- Takahashi K, Daughaday WH and Kipnis DM (1971) Regulation of immunoreactive growth hormone secretion in male rats *Endocrinology* **88** 909–917
- Tannenbaum GS and Martin JB (1976) Evidence for an endogenous ultradian rhythm governing growth hormone secretion in the rat *Endocrinology* **98** 562–570
- Terry LC, Willoughby JO, Brazeau P, Martin JB and Patel Y (1976) Antiserum to somatostatin prevents stress-induced inhibition of growth hormone secretion in the rat *Science* **192** 565–567
- Vermouth NT and Deis RP (1972) Prolactin release induced by prostaglandin F_{2a} in pregnant rats *Nature* **238** 248–250
- Vermouth NT and Deis RP (1974) Prolactin release and lactogenesis after ovariectomy in pregnant rats: effect of ovarian hormones *Journal of Endocrinology* **63** 13–20
- Willoughby JO and Day TA (1981) Central catecholamine depletion: effects on physiological growth hormone and prolactin secretion *Neuroendocrinology* **32** 65–69
- Willoughby JO, Menadue MF and Liebelt H (1987) Activation of serotonin receptors in the medial basal hypothalamus stimulates growth hormone secretion in the unanesthetized rat *Brain Research* **404** 319–322