Activation of temperature-dependent flagellar movement of demembranated fowl spermatozoa: involvement of an endogenous serine protease

K. Ashizawa¹, S. Katayama¹, G. J. Wishart², A. Shiraishi¹ and Y. Tsuzuki¹

¹Laboratory of Animal Reproduction, Faculty of Agriculture, Miyazaki University, Miyazaki 889-21, Japan; and ²Department of Molecular and Life Sciences, Dundee Institute of Technology, Dundee DD1 1HG, UK

In the presence of ATP, the motility of demembranated fowl spermatozoa was vigorous at 30°C, but negligible at 40°C. Motility could be restored at 40°C by the addition of 10–100 ng trypsin ml⁻¹. Chymotrypsin also stimulated the motility, but neither papain nor carboxypeptidase B appreciably affected motility. Conversely, at 30°C sperm motility was inhibited by aprotinin or phenylmethylsulfonyl fluoride. These results suggest that endogenous protease, presumably serine protease, activity is instrumental in the regulation of fowl sperm motility. It seems likely that the site of action of this protease is axonemal, but a direct effect of added protease on dynein ATPase activity could not be demonstrated.

Introduction

Flagellar movement of spermatozoa is based on the active sliding of microtubules as a result of ATP hydrolysis by dynein ATPase (for review, see Tash and Means, 1983; Lindemann and Kanous, 1989). Although this basic axonemal mechanism is fairly well understood, the factors and mechanisms of its regulation have still to be clarified. There appear to be several regulatory mechanisms that individually or synergistically control sperm motility: Ca²⁺ and Ca²⁺-associated compounds, cAMP and intracellular pH (Majumder *et al.*, 1990). In addition, current evidence suggests that a protease activity with a Lys- and Arg-ester bond specificity is required for sperm motility. It is assumed that this regulatory system is probably located near the dynein arms, but does not directly involve the force-generating dynein ATPase (de Lamirande *et al.*, 1990).

Unlike mammalian spermatozoa, fowl spermatozoa show definitive temperature-dependent changes in their motility: in most synthetic diluents, they become immotile at the avian body temperature of 40–41°C, and motility is restored by decreasing the temperature (Munro, 1938; Nevo and Schindler, 1968; Ashizawa and Nishiyama, 1978; Takeda, 1982; Ashizawa and Okauchi, 1984; Ashizawa and Wishart, 1987; Wishart and Ashizawa, 1987; Ashizawa *et al.*, 1989a, b; Thomson and Wishart, 1989, 1991). The axoneme itself appears to be directly involved in this regulatory system, since the motility of demembranated spermatozoa is, as with intact spermatozoa, negligible at 40°C and restored at 30°C (Ashizawa *et al.*, 1989a, b). Furthermore, cAMP-independent phosphorylation of a 43 kDa axonemal protein is likely to be a regulatory step in the maintenance of fowl sperm motility (Ashizawa *et al.*, 1992).

If the activity of endogenous proteases is involved in the above temperature-dependent regulatory system, then fowl spermatozoa might be expected to be made immotile at 30°C by the addition of protease inhibitors and, conversely, motile at 40°C by the addition of appropriate exogenous proteases. In this study, the effects of some proteases and protease inhibitors on the motility of demembranated fowl spermatozoa were therefore examined at 30°C and 40°C, and information was obtained concerning the locus of action of such proteases.

Materials and Methods

Animals and preparation of spermatozoa

Commercial White Leghorn roosters (Babcock strain, Akagi Poultry Breeding Farm, Miyazaki) were used throughout the study. All birds were housed in individual cages and fed *ad libitum* on a commercial breeder diet. They were exposed to a 14 h light:10 h dark cycle.

Semen was collected by the method of Bogdonoff and Shaffner (1954). Samples of semen pooled from four to six males were diluted approximately tenfold in 150 mmol NaCl l⁻¹ with 20 mmol TES (N-Tris-[hydroxymethyl]-methyl-2-aminoethanesulfonic acid) l⁻¹ at pH 7.4 and centrifuged at 700 g for 13 min at room temperature (20–25°C). The washed spermatozoa were reconstituted in the same buffer to give a final concentration of approximately 1 × 10⁹ cells ml⁻¹. Samples of 3–4 ml were poured into 30 ml Erlenmeyer flasks with a screw cap.

Chemicals

Trypsin (approximately 10 000 BALL A₂₅₃ units mg⁻¹ protein) was obtained from Mochida Pharmaceutical Co.

(Tokyo). Chymotrypsin (40–60 BTEE $\mu\text{mol mg}^{-1}$ protein), papain (10–20 BAEE $\mu\text{mol mg}^{-1}$ protein), carboxypeptidase B (approximately 100 HA $\mu\text{mol mg}^{-1}$ protein), aprotinin, PMSF, adenosine 5'-triphosphate (ATP), dithiothreitol, potassium glutamate, TES, Triton X-100, phosphodiesterase and desiccated firefly tails were purchased from Sigma Chemical Co. (St Louis, MO). [γ - ^{32}P]ATP was purchased from Du Pont-New England Nuclear (Boston, MA). Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) molecular weight standards were obtained from Bio-Rad Laboratories (Richmond, CA). Other chemicals were of reagent grade from Nacalai Tesque, Inc. (Kyoto).

Measurement of motility of demembranated spermatozoa

Intact spermatozoa were preincubated at 30°C or 40°C for 10 min. After the preincubation, demembration and reactivation were performed according to the method described by Ashizawa *et al.* (1989b). The extraction medium used consisted of 0.1% Triton X-100, 0.2 mol sucrose l^{-1} , 25 mmol potassium glutamate l^{-1} , 1 mmol MgSO_4 l^{-1} , 1 mmol dithiothreitol l^{-1} and 20 mmol Tris-HCl buffer l^{-1} (pH 7.9). The reactivation medium consisted of 0.5 mmol ATP l^{-1} , 0.2 mol sucrose l^{-1} , 25 mmol potassium glutamate l^{-1} , 1.5 mmol MgSO_4 l^{-1} , 1 mmol dithiothreitol l^{-1} and 20 mmol Tris-HCl buffer l^{-1} (pH 7.9). To examine the effects of exogenous proteases or protease inhibitors, various concentrations of trypsin, chymotrypsin, papain or carboxypeptidase B at 40°C, and aprotinin or PMSF at 30°C were added to the reactivation medium. Phosphodiesterase, EGTA or Ca^{2+} were also added to trypsin-treated spermatozoa. The suspension of demembranated spermatozoa was viewed in a microscope slide chamber (Sekisui Chemical Co., Ltd, Tokyo, UR-157 type) on a thermostatically controlled warm plate, and the motility of spermatozoa was recorded by videomicroscopy at 30°C or 40°C (Katz and Overstreet, 1981).

Measurement of crude dynein ATPase activity

Crude dynein extract from fowl spermatozoa was obtained by the method described by Ashizawa and Hori (1990). ATPase activities were assayed by the reduction of ATP concentrations determined by firefly bioluminescence (Ashizawa and Hori, 1990). Protein concentrations of crude dynein extract were determined according to published methods (Bradford, 1976), with BSA as the standard. The rate of ATPase activity was expressed in terms of nmol ATP hydrolysed mg^{-1} protein min^{-1} .

Phosphorylation of endogenous proteins and electrophoresis

Phosphorylation reaction and electrophoresis on polyacrylamide gels of demembranated sperm proteins were carried out according to the methods described by Ashizawa *et al.* (1992). Briefly, demembranated spermatozoa with extraction-activation medium containing 0.1% Triton X-100, 0.2 mol sucrose l^{-1} , 25 mmol potassium glutamate l^{-1} , 1 mmol MgSO_4 l^{-1} , 1 mmol dithiothreitol l^{-1} , 20 mmol Tris-HCl buffer l^{-1} (pH 7.9), 0.1 mmol ATP l^{-1} and approximately 7000 c.p.m.

[γ - ^{32}P]ATP pmol^{-1} were incubated for 2 min at 30°C or 40°C. Aprotinin or trypsin, when incorporated, was present at a final concentration of 0.24 TIU (trypsin inhibitor unit) ml^{-1} or 50 ng ml^{-1} , respectively. The phosphorylation reaction was terminated by the addition of equal volumes of concentrated (twofold) Laemmli (1970) sample buffer and boiling for 5 min. Samples containing protein from approximately 1.3×10^6 spermatozoa were loaded on to 10% SDS-polyacrylamide slab gels, and electrophoresed. Autoradiography was performed at -80°C for 2–4 days exposure to X-ray film with an intensifying screen (Lightning plus, Du Pont, Wilmington, DE).

Statistical analysis

Statistical comparisons were performed using Student's *t* test.

Results

Effects of proteases and protease inhibitors on the motility of demembranated fowl spermatozoa

Reactivation of fowl spermatozoa without the addition of trypsin was negligible at 40°C. In contrast, the presence of 10–100 ng trypsin ml^{-1} permitted reactivation of sperm motility at 40°C, and optimum reactivation occurred at 50 ng trypsin ml^{-1} . At higher concentrations, fewer spermatozoa were reactivated (Fig. 1a). Chymotrypsin also stimulated the motility of demembranated spermatozoa at 40°C, and maximum motility was obtained at 600 ng chymotrypsin ml^{-1} (Fig. 1b). However, no stimulation of motility was observed following the addition of papain (Fig. 1a) or carboxypeptidase B (Fig. 1b), within the ranges 0–2 $\mu\text{g ml}^{-1}$ and 0–500 $\mu\text{g ml}^{-1}$, respectively, and disintegration of sperm tails was observed at higher concentrations. The addition of papain and carboxypeptidase B at lower concentrations (ng ml^{-1}) was also ineffective in initiating motility (data not shown).

However, the vigorous motility of demembranated spermatozoa at 30°C was inhibited in a dose-dependent manner by the addition of the protease inhibitors, aprotinin or PMSF (Fig. 2). However, the addition of 500 ng trypsin ml^{-1} released the inhibitory effect of aprotinin within 1 min (Fig. 3).

Neither the addition of phosphodiesterase to remove endogenous cAMP nor the addition of Ca^{2+} or EGTA to trypsin-treated spermatozoa had any significant effect on sperm motility at 40°C (Table 1).

Effects of proteases on dynein ATPase activity and the phosphorylation state of proteins of demembranated fowl spermatozoa

The flagellar ATPase activity of crude dynein extract without the addition of proteases was approximately 22 nmol ATP hydrolysed mg^{-1} protein min^{-1} . This activity was not stimulated by the addition of proteases, including trypsin and chymotrypsin (Table 2).

A marked difference in the phosphorylation status of a 43 kDa protein was obtained in demembranated spermatozoa at 30°C and at 40°C. This protein was slightly phosphorylated

Fig. 1. The motility of demembrated fowl spermatozoa in the reactivation medium at 40°C after addition of various concentrations of (a) (●) trypsin or (○) papain and (b) chymotrypsin (●) or carboxypeptidase B (○). Results are the mean ($\pm$ SEM) from five samples of spermatozoa.

Fig. 2. The motility of demembrated fowl spermatozoa in the reactivation medium at 30°C after addition of various concentrations of aprotinin or phenylmethylsulfonyl fluoride (PMSF). Results are the mean ($\pm$ SEM) from five samples of spermatozoa. TIU: trypsin inhibitor units.

at 40°C, but strongly phosphorylated at 30°C, confirming the previous observation of Ashizawa *et al.* (1992). However, neither aprotinin at 30°C nor trypsin at 40°C affected the phosphorylated state of this or other phosphorylated proteins (Fig. 4).

Fig. 3. The motility of demembrated fowl spermatozoa in the reactivation medium at 30°C following addition of (●) 0.24 trypsin inhibitor units aprotinin ml⁻¹ or (○) 500 ng trypsin ml⁻¹. Results are the mean ($\pm$ SEM) from five samples of spermatozoa.

Table 1. The effects of phosphodiesterase, EGTA and Ca²⁺ on the motility of trypsin-treated demembrated fowl spermatozoa at 40°C

Treatments	Motility (%)
None (control)	4.8 $\pm$ 1.9 ^a
Trypsin (50 ng ml ⁻¹)	62.1 $\pm$ 4.2 ^b
Trypsin (50 ng ml ⁻¹) + phosphodiesterase (2 U ml ⁻¹)	70.5 $\pm$ 5.3 ^b
EGTA (2 mmol l ⁻¹)	4.1 $\pm$ 1.8 ^a
Ca ²⁺ (0.1 μ mol l ⁻¹)	2.7 $\pm$ 1.3 ^a
Trypsin (50 ng ml ⁻¹) + EGTA (2 mmol l ⁻¹)	72.1 $\pm$ 3.5 ^b
Trypsin (50 ng ml ⁻¹) + Ca ²⁺ (0.1 μ mol l ⁻¹)	70.4 $\pm$ 2.6 ^b

Each value represents the mean ($\pm$ SEM) of five samples of spermatozoa. Values with different superscripts differ significantly ($P < 0.01$) from each other.

Table 2. The effects of proteases on the ATPase activity of crude dynein extract of fowl spermatozoa at 40°C

Protease	ATPase activity (nmol ATP hydrolysed mg ⁻¹ protein min ⁻¹)
None (control)	21.8 $\pm$ 0.3 ^a
Trypsin (50 ng ml ⁻¹)	20.5 $\pm$ 0.7 ^a
Chymotrypsin (600 ng ml ⁻¹)	20.9 $\pm$ 0.9 ^a
Papain (600 ng ml ⁻¹)	20.9 $\pm$ 0.7 ^a
Carboxypeptidase B (40 μ g ml ⁻¹)	21.0 $\pm$ 0.3 ^a

Each value represents the mean ($\pm$ SEM) of five samples of spermatozoa. Values with different superscripts differ significantly ($P < 0.01$) from each other.

Discussion

Protease activities were mainly thought to be involved in the catabolism of proteins, but recent evidence suggests that they

Fig. 4. 10% SDS-PAGE profile of demembrated fowl sperm proteins stained with Coomassie blue (lane 5) and corresponding autoradiography of phosphorylated proteins (lanes 1–4). Triton X-100 extracted and reactivated samples were incubated at 30°C or 40°C for 2 min. Lane 1: 30°C in the absence of aprotinin; lane 2: 30°C in the presence of 0.24 trypsin inhibitor units aprotinin ml⁻¹; lane 3: 40°C in the absence of trypsin; lane 4: 40°C in the presence of 50 ng trypsin ml⁻¹.

also have a short-term regulatory function, for example in the control of sperm motility (Gagnon and de Lamirande, 1987). The present study showed that temperature-dependent immobilization of demembrated fowl spermatozoa at 40°C was reversed by the addition of trypsin or chymotrypsin. Furthermore, sperm motility at 30°C was inhibited by aprotinin or PMSF, but could be restored by the subsequent addition of trypsin. It is therefore proposed that an endogenous, presumably serine, protease is present in the fowl sperm axoneme, and plays an important role in regulating sperm movement. The immobilization of fowl spermatozoa at 40°C might be due in part to the reduction of activity of this protease, although the effects of temperature on the activity of proteases of fowl spermatozoa were not measured in this experiment.

Although this is the first report of the presence of such protease activity in avian spermatozoa, similar sperm proteases appear to be preserved throughout evolution from sea urchins to humans (de Lamirande *et al.*, 1990), as protease inhibitors, including aprotinin, inhibited the motility of demembrated spermatozoa in sea urchins, fish and mammals (de Lamirande *et al.*, 1983; de Lamirande and Gagnon, 1986; Cosson and Gagnon, 1988; Inaba and Morisawa, 1991).

The target and precise mechanisms of action of proteases in the regulation of sperm motility remain to be elucidated. The inhibition of motility initiation by high concentrations of ATP was abolished by the inclusion of trypsin in demembrated

hamster spermatozoa, and it is assumed that trypsin may involve the action of cAMP (Yeung, 1986), since a trypsin-like protease derived from bovine spermatozoa can stimulate adenylyl cyclase (Johnson *et al.*, 1985). In the present study, however, no decrease in motility was observed by the addition of phosphodiesterase to remove endogenous cAMP from trypsin-treated spermatozoa. Moreover, trypsin-stimulated sperm motility was not affected by the removal or addition of Ca²⁺. From these results, it may be proposed that the proteases act directly on the axoneme in a cAMP- and Ca²⁺-independent manner. However, the results reported here also showed that ATPase activity of a dynein extract of fowl spermatozoa was not stimulated by the addition of proteases tested, suggesting that these proteases might not act directly on dynein ATPase. Similarly, trypsin did not stimulate the steady-state activity of the axonemal dynein ATPase of sea urchin spermatozoa (Kamimura *et al.*, 1985) and aprotinin, at 50-fold the concentration needed to block sperm motility, caused only a 30% inhibition of dynein ATPase isolated from bull spermatozoa (Gagnon and de Lamirande, 1987).

Recently, we suggested that cAMP-independent phosphorylation of a 43 kDa axonemal protein is likely to be a regulatory step in the maintenance of fowl sperm motility (Ashizawa *et al.*, 1992). However, the phosphorylation state of this protein did not change in the presence of proteases or their inhibitors, which supports the hypothesis that the site of action of endogenous protease may lie between the 43 kDa protein and the dynein-tubulin sliding system. This hypothesis agrees with the suggestion that other sperm proteases may act near the site where dynein arms interact with microtubules, without affecting the dynein ATPase (de Lamirande *et al.*, 1990).

This study was supported by a grant from the Ministry of Education, Science and Culture, Japan. G. J. Wishart was supported by the Royal Society of Edinburgh.

References

- Ashizawa K and Hori M (1990) Activation of dynein adenosine triphosphatase and flagellar motility of demembrated spermatozoa by monovalent salts at 40°C in the domestic fowl, *Gallus domesticus* *Comparative Biochemistry and Physiology* **97A** 325–328
- Ashizawa K and Nishiyama H (1978) Effects of temperature on the vigour of motility, oxygen consumption and duration of motility of fowl spermatozoa under aerobic conditions *Japanese Poultry Science* **15** 264–266
- Ashizawa K and Okauchi K (1984) Stimulation of sperm motility and oxygen consumption of fowl spermatozoa by a low molecular weight fraction of seminal plasma *Journal of Reproduction and Fertility* **71** 593–598
- Ashizawa K and Wishart GJ (1987) Resolution of the sperm motility-stimulating principle of fowl seminal plasma into Ca²⁺ and an unidentified low molecular weight factor *Journal of Reproduction and Fertility* **81** 495–499
- Ashizawa K, Maeda S and Okauchi K (1989a) The mechanisms of reversible immobilization of fowl spermatozoa at body temperature *Journal of Reproduction and Fertility* **86** 271–276
- Ashizawa K, Suzuki Y and Okauchi K (1989b) Flagellar movement in demembrated preparations of ejaculated fowl spermatozoa *Journal of Reproduction and Fertility* **86** 263–270
- Ashizawa K, Katayama S and Tsuzuki Y (1992) Regulation of flagellar motility by temperature-dependent phosphorylation of a 43 kDa axonemal protein in fowl spermatozoa *Biochemical and Biophysical Research Communications* **185** 740–745
- Bogdonoff PD Jr and Shaffner CS (1954) The effect of pH on *in vitro* survival, metabolic activity, and fertilizing capacity of chicken semen *Poultry Science* **33** 665–669

- Bradford MM** (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding *Analytical Biochemistry* **72** 248–254
- Cosson MP and Gagnon C** (1988) Protease inhibitor and substrates block motility and microtubule sliding of sea urchin and carp spermatozoa *Cell Motility and the Cytoskeleton* **10** 518–527
- de Lamirande E and Gagnon C** (1986) Effects of protease inhibitors and substrates on motility of mammalian spermatozoa *Journal of Cell Biology* **102** 1378–1383
- de Lamirande E, Bardin CW and Gagnon C** (1983) Aprotinin and a seminal plasma factor inhibit the motility of demembranated reactivated rabbit spermatozoa *Biology of Reproduction* **28** 788–796
- de Lamirande E, Cosson MP and Gagnon C** (1990) Protease involvement in sperm motility. In *Controls of Sperm Motility: Biological and Clinical Aspects* pp 241–250 Ed. C Gagnon. CRC Press, Boca Raton
- Gagnon C and de Lamirande E** (1987) Protease involvement in sperm motility. In *New Horizons in Sperm Cell Research* pp 205–214 Ed. H Mohri. Japan Scientific Societies Press, Tokyo
- Inaba K and Morisawa M** (1991) A chymotrypsin-like protease involved in motility of sperm in salmonid fish *Biomedical Research* **12** 435–437
- Johnson RA, Jakobs KH and Schultz G** (1985) Extraction of the adenylate cyclase-activating factor of bovine sperm and its identification as a trypsin-like protease *Journal of Biological Chemistry* **260** 114–121
- Kamimura S, Yano M and Shimizu H** (1985) ATP hydrolysis coupled to microtubule sliding in sea-urchin sperm flagella *Journal of Biochemistry* **97** 1509–1515
- Katz DF and Overstreet JW** (1981) Sperm motility assessment by videomicrography *Fertility and Sterility* **35** 188–193
- Laemmli UK** (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4 *Nature* **227** 680–685
- Lindemann CB and Kanous KS** (1989) Regulation of mammalian sperm motility *Archives of Andrology* **23** 1–22
- Majumder GC, Dey CS, Haldar S and Barua M** (1990) Biochemical parameters of initiation and regulation of sperm motility *Archives of Andrology* **24** 287–303
- Munro SS** (1938) Fowl sperm immobilization by a temperature-media interaction and its biological significance *Quarterly Journal of Experimental Physiology* **27** 281–287
- Nevo AC and Schindler H** (1968) Reversible inactivation of cock spermatozoa by temperature. *Proceedings of the Sixth International Congress of Animal Reproduction and Artificial Insemination, Paris, II*, 1637–1640
- Takeda A** (1982) Studies on reversible inactivation of cock spermatozoa by temperature. I. Effects of several factors on reversible inactivation *Japanese Poultry Science* **19** 26–36
- Tash JS and Means AR** (1983) Cyclic adenosine 3',5' monophosphate, calcium and protein phosphorylation in flagellar motility *Biology of Reproduction* **28** 75–104
- Thomson MF and Wishart GJ** (1989) Elucidation of the mechanism responsible for the temperature-dependent reversible inactivation of the motility of fowl spermatozoa *British Poultry Science* **30** 687–692
- Thomson MF and Wishart GJ** (1991) Temperature-mediated regulation of calcium flux and motility in fowl spermatozoa *Journal of Reproduction and Fertility* **93** 385–391
- Wishart GJ and Ashizawa K** (1987) Regulation of motility of fowl spermatozoa by calcium and cAMP *Journal of Reproduction and Fertility* **80** 607–611
- Yeung CH** (1986) Temporary inhibition of the initiation of motility of demembranated hamster sperm by high concentrations of ATP *International Journal of Andrology* **9** 359–370